



unidentified compounds. 2-Heptyl-1,3-dioxolane or 1-octanal which arises from hexyl group migration was not formed in the reaction products. Formation of 2-hexyl-1,4-dioxane (**5**) shows that the reaction corresponding to Scheme 3b is partly operative in this case. A similar reaction in 1,3-propanediol gave no compound corresponding to **5**.

Although thallium(III) was almost completely reduced to thallium(I) in these reactions, the yields of the products were generally low. This may be due to the consumption of thallium(III) for the oxidation of diols as was noted by Kabbe in the case of some 1,2-diols.<sup>5)</sup> We also confirmed that when thallium(III) acetate (10 mmol) is heated in 1,2-ethanediol (7.5 ml) at 82 °C for 4 h, 65% of thallium(III) is reduced to thallium(I) (by iodometry).

### Experimental

Commercial organic and inorganic materials were used. The IR and NMR spectra were recorded with a Hitachi EPI-S2 and a Varian EM-360 apparatus, respectively. Gas chromatography was carried out on Shimadzu 4BMPF and 5APTF apparatus using EGSS-X(15%)-Chromosorb-W (1 m and 3 m) and PEG-6000(25%)-Chromosorb-W (3 m) columns ( $N_2$  as a carrier gas).

**Oxidation of Olefin with  $Tl(OAc)_3$  in Diol.** A typical experimental procedure is as follows. A mixture of styrene (4.1 g, 40 mmol),  $Tl(OAc)_3$  (7.7 g, 20 mmol), and 1,2-ethanediol (15 ml) was heated at 80 °C for 4 h under stirring. After the usual work-up, GLC analysis of benzene extract revealed the presence of 7.2 mmol (36% yield based on  $Tl^{3+}$  charged) of 2-benzyl-1,3-dioxolane (**2a**) and small amounts of phenylacetaldehyde and acetophenone as the products, butyrophenone being used as an internal standard. Distillation gave 0.5 g of pure **2a**. Thallium(III) was almost completely reduced to thallium(I). Authentic samples of **2a–e**, **4**, and **6** were prepared by the acetalization or ketalization of phenylacetaldehyde, 2-phenylpropionaldehyde, acetophenone and 2-octanone in the corresponding diols<sup>6)</sup> and their structures were confirmed by NMR; **2a** bp 80–93 °C/2 Torr, **2b** bp 81 °C/2 Torr, **2c** bp 93–95 °C/4 Torr, **2d** bp 100–104.5 °C/4 Torr, **2e** bp 106–110 °C/4 Torr, **4** bp 83–84 °C/15 Torr, **6** bp 78–79 °C/15 Torr. **5** was prepared by the alkaline dehydrobromination of 1-bromo-2-(2-hydroxyethoxy)octane which was synthesized from 1-octene, 1,2-ethanediol, and *N*-bromosuccinimide.<sup>7)</sup> **8** was prepared by the reaction of 1-bromo-2-(2-hydroxyethoxy)octane with KOAc in DMSO; bp 120–123 °C/5 Torr, NMR ( $CCl_4$ )  $\delta$  0.67–1.07 (m, 3H), 1.07–1.73 (m, 10H), 1.98 (s, 3H), 2.3 (br. 1H), 3.23–3.67 (m, 5H), and 3.67–4.23 (m, 2H). No 2-acetoxy-1-(2-hydroxyethoxy)octane (positional isomer of **8**) was detected by NMR in the oxidation products of 1-octene.

**Isolation of **3**.** A mixture of styrene (4.1 g, 40 mmol),  $Tl(OAc)_3$  (7.7 g, 20 mmol), and 1,2-ethanediol (15 ml) was stirred at 15–25 °C for 20 h. To this was added 500 ml of petroleum ether and the mixture was then left to stand for 3 days in a refrigerator to give a white crystal of **3** as a precipitate. **3** was collected by filtration [4.2 g, 43% yield; mp 133–135 °C(dec)]. The treatment of organic filtrate as described above revealed the presence of 28% yield of **2a**. **3** is soluble in pyridine, DMSO, DMF, hot  $CHCl_3$ , and hot ethanol, but insoluble in water, THF, ether, benzene,  $CCl_4$ , acetonitrile, and acetic acid. IR (hexachlorobutadiene and paraffin mulls): 3250(s), 2950(m), 2920(m), 2860(m), 1625(sh), 1610(s), 1530(s), 1430(s), 1360(s), 1320(s), 1110(m), 1090(m), 1067(m), 974(m), 762(m), 695(m), and 685(m)  $cm^{-1}$ . NMR (pyridine- $d_5$ ):  $\delta$  2.09 (s, 6H), 3.10 (d, 1H,  $J_{Ti-H} = 732$  Hz), 3.14 (d, 1H,  $J_{Ti-H} = 875$  Hz), 3.53 (t, 2H), 3.76 (t, 2H), 5.01 (d, 1H,  $J_{Ti-H} = 739$  Hz), 5.76 (s, 1H), and 6.9–7.6 (m, 5H). Found: C, 34.48; H, 4.11; Tl, 41.28%. Calcd for  $C_{14}H_{18}O_6Tl$ : C, 34.48; H, 3.93; Tl, 41.91%.

**Reaction of **3** with Palladium(II) Salt.** **3** (0.98 g, 2 mmol) was added to a hot mixture (75–80 °C) of  $PdCl_2$  (0.36 g, 2.03 mmol), NaOAc (0.33 g, 4.02 mmol), and 1,2-ethanediol (10 ml) and the resulting mixture was stirred for 20 min. A black precipitate formed was filtered off. Saturated aq. NaCl was added to the filtrate which was then extracted with benzene. GLC analysis of the benzene extract revealed the presence of acetophenone (0.19 mmol) and 2-methyl-2-phenyl-1,3-dioxolane (**4**, 1.81 mmol) together with a trace amount of **2a**, *p*-methylacetophenone being used as an internal standard.

### References

- 1) Presented at the 33rd Annual Meeting of the Chemical Society of Japan, Fukuoka, October 19, 1975.
- 2) a) R. Jira and W. Freiesleben, "Organometallic Reactions," Vol. 3, ed. by E. I. Becker and M. Tsutsui, Wiley-Interscience, New York, N. Y. (1972), pp. 157–158; b) D. F. Hunt and G. T. Rodeheaver, *Tetrahedron Lett.*, **1972**, 3595.
- 3) a) S. Uemura, K. Zushi, M. Okano, and K. Ichikawa, *J. Chem. Soc., Chem. Commun.*, **1972**, 234; b) S. Uemura, A. Tabata, and M. Okano, *J. Chem. Soc., Chem. Commun.*, **1972**, 517; c) S. Uemura, K. Zushi, A. Tabata, A. Toshimitsu, and M. Okano, *Bull. Chem. Soc. Jpn.*, **47**, 920 (1974); d) S. Uemura, A. Toshimitsu, M. Okano, and K. Ichikawa, *Bull. Chem. Soc. Jpn.*, **48**, 1925 (1975).
- 4) A. Lethbridge, R. O. C. Norman, and C. B. Thomas, *J. Chem. Soc., Perkin Trans. 1*, **1974**, 1929.
- 5) H. J. Kabbe, *Justus Liebigs Ann. Chem.*, **656**, 204 (1962).
- 6) S. R. Sandler and W. Karo, "Organic Functional Group Preparations," Part III, Academic Press, New York, N. Y. (1972), Chap. 1.
- 7) A. Iovchev, H. Reinheckel, L. Stoilov, and K. Haage, *Monatsh. Chem.*, **102**, 114 (1971).