

Cyclopropanes from Allylic Halides. I. Synthesis of Dimethyl 3-(2-Methyl-1-propenyl)-2,2-dimethylcyclopropane-1,1-dicarboxylate and Pyrocin as Precursors of Chrysanthemic Acid


Sigeru TORII, Hideo TANAKA, and Yosio NAGAI

Department of Industrial Chemistry, School of Engineering, Okayama University, Okayama 700


(Received May 9, 1977)

Synopsis. Dimethyl 3-(2-methyl-1-propenyl)-2,2-dimethylcyclopropane-1,1-dicarboxylate and pyrocin as precursors of chrysanthemic acid were synthesized by the reaction of (2-halo-2-methylpropylidene)malonate with 2-methyl-1-propenylmagnesium bromide.

Chrysanthemic acid (**1**), an important moiety of pyrethrins and a naturally occurring pesticide, has been a subject of interest as regards its synthesis.¹⁾ In contrast to the nucleophilic substitution reaction of allylic halides taking place at the α - or γ -position, nucleophilic attack with cyanide at the β -carbon of allylic halides **2**





bearing alkoxy carbonyl groups in the γ -position followed by cyclization, giving cyclopropanecarboxylates **3**, was observed.²⁾ The diverse nature of allylic halides prompted us to examine the reaction of dimethyl (2-halo-2-methylpropylidene)malonate (**4**) with various nucleophiles.




The present paper describes the results obtained in the reaction of **4** with 2-methyl-1-propenylmagnesium bromide³⁾ as a nucleophile as well as the synthesis of pyrocin (**7b**)⁴⁾ using the same starting materials.

(2-Halo-2-methylpropylidene)malonate (**4**) was treated with 2-methyl-1-propenylmagnesium bromide in tetrahydrofuran under the conditions given in Table 1. Chloride **4a** was allowed to react with 2-methyl-1-propenylmagnesium bromide at 24 °C for 45 h to give the desired **5** in 46% yield as well as 11% of **6** and the recovered **4a** (30%) (run 1), whereas the reaction at 45 °C for 18 h resulted in a mixture of 55% of **5**, 19% of **6**, and 14% of **7a** (run 2). When the same reaction was carried out in the presence of catalytic amounts of copper(I) chloride,⁵⁾ most of **4a** was consumed within 45 min, giving 51% of **5** and 37% of **6** along with minor product **7a** (4%) (run 3). The presence of a good leaving group in the allylic halide **4** is not sufficient for the reaction, since the same reaction with **4b** (runs 4




and **5**) carried out at 24 °C afforded the dehalogenated product **6** principally⁶⁾ and **5** showed only 25—32% yields as compared to 46—55% for **5** from **4a**.

The ambident ion **8** is believed to be formed by an unusual nucleophilic reaction of 2-methyl-1-propenide ion on the β -position of **4**. The formation of the cyclopropane ring of **5** can be rationalized by S_N2 type nucleophilic attack of the carbanion of **8** to the γ -position (Path a), while the formation of the parent compound **7a** could be attributed to nucleophilic attack by the enolate ion at the γ -position (Path b).



When **4b** was heated to 170 °C for 2 h under nitrogen, the 2-butenolide **9** was obtained in 76% yield. Subsequently, Michael addition of **9** with 2-methyl-1-propenylmagnesium bromide gave **7a** in 85% yield.⁷⁾ Demeth oxycarbonylation of **7a** in wet dimethyl sulfoxide-sodium chloride at 170 °C afforded pyrocin (**7b**) smoothly.



Experimental

All boiling points and melting points were uncorrected, the boiling points indicated being air-bath temperatures. IR spectra were recorded on a Japan Spectroscopic Co., Ltd., IRA-I infrared recording spectrophotometer with a grating.

TABLE 1. REACTION OF **4a** AND **4b** WITH 2-METHYL-1-PROPENYLMAGNESIUM BROMIDE

Run	Substrate (mmol)	$\text{CH}_2=\text{CHMgBr}$ mmol	Additive	Temp °C	Time h	Products, % ^{a)}			
						5	6	7a	4
1	4a (0.6)	1.4	none	24	45	46	11	—	30
2	4a (4.5)	9.0	none	45	18	55	19	14	—
3	4a (0.6)	2.0	CuCl ^{b)}	45	0.75	51	37	4	5
4	4b (0.7)	0.8	CuCl ^{b)}	24	0.75	25	47	5	7
5	4b (0.6)	1.0	none	24	12	32	56	12	—

a) Isolated Yields. b) Copper(I) chloride (10 mg) was added.

NMR spectra were recorded at 60 MHz on a Hitachi R-24 spectrometer with an internal standard of tetramethylsilane. Elemental analyses were carried out in this laboratory.

Dimethyl (2-Chloro-2-methylpropylidene)malonate (4a). A mixture of dimethyl (2-methylpropylidene)malonate⁸⁾ (3.0 g, 16.1 mmol), *t*-butyl hypochlorite⁹⁾ (2.0 g, 18.4 mmol), and azobisisobutyronitrile (AIBN) (30 mg) in CCl₄ (3 ml) was heated under reflux for 2 h. After removal of the volatile substance, CCl₄ (3 ml), *t*-butyl hypochlorite (2.0 g, 18.4 mmol) and AIBN (30 mg) were added to the residue. The resulting mixture was refluxed for 2 h. The operation was repeated 5 times and the resulting oil was distilled to give **4a** (1.78 g, 50%): bp 75–77 °C/3 Torr; IR (neat) 1740 (C=O), 1651 cm⁻¹ (C=C); NMR (CDCl₃) δ 1.76 (s, 6, CH₃CCl), 3.77 (s, 6, CH₃), 6.93 (s, 1, HC=C). Found: C, 49.17; H, 6.01%. Calcd for C₉H₁₃O₄: C, 48.99; H, 5.94%.

Dimethyl (2-Bromo-2-methylpropylidene)malonate (4b). **4b** was prepared by bromination of dimethyl (2-methylpropylidene)malonate with *N*-bromosuccinimide, bp 92–95 °C/18 Torr, according to the reported procedure.¹⁰⁾

Dimethyl 2,2-Dimethyl-3-(2-methyl-1-propenyl)cyclopropane-1,1-dicarboxylate (5). The reaction conditions and results are given in Table 1. A typical experimental procedure (run 2) is as follows: To a THF solution (3 ml) of **4a** (1.0 g, 4.5 mmol) was added dropwise a THF solution (7 ml) of 2-methyl-1-propenylmagnesium bromide,³⁾ prepared from 2-methyl-1-propenyl bromide (1.22 g, 9.0 mmol) and magnesium (0.22 g, 9.0 mmol) at room temperature. After being stirred for 30 min at this temperature and for 18 h at 45 °C, the mixture was quenched with aqueous 10% NH₄Cl, extracted with ether, washed with brine and dried (Na₂SO₄). Removal of the solvent followed by column chromatography (SiO₂, THF–hexane, 1/20) gave **5** (597 mg, 55%), **6** (160 mg, 19%), and **7a** (143 mg, 14%).

Compound 5: Bp 81–84 °C/10 Torr; IR (neat) 1730 cm⁻¹ (C=O); NMR (CDCl₃) δ 1.23 (s, 3, CH₃), 1.27 (s, 3, CH₃), 1.74 (br s, 6, CH₃C=C), 2.44 (d, 1, *J*=8 Hz, CH), 3.68 (s, 3, CH₃O), 3.72 (s, 3, CH₃O), 5.02 (diffused d, 1, *J*=8 Hz, HC=C). Found: C, 64.99; H, 8.59%. Calcd for C₁₃H₂₀O₄: C, 64.98; H, 8.39%.

Compound 6: Bp 73–76 °C/8 Torr; IR (neat) 1735 cm⁻¹ (C=O); NMR (CDCl₃) δ 1.67 (diffused s, 3, CH₃C=C), 1.77 (diffused s, 3, CH₃C=C), 3.71 (s, 6, CH₃O), 4.24 (d, 1, *J*=9 Hz, CHCO), 5.43 (diffused d, 1, *J*=9 Hz, HC=C). Found: C, 58.29; H, 7.47%. Calcd for C₉H₁₄O₄: C, 58.05; H, 7.58%.

Compound 7a: Mp 81–81.5 °C (hexane–ether, 10/1); IR (Nujol) 1770 (lactone C=O), 1734 (C=O), 1675 cm⁻¹ (C=C); NMR (CDCl₃) δ 1.27 (s, 3, CH₃), 1.44 (s, 3, CH₃), 1.73 (m, 6, CH₃C=C), 3.48 (d, 1, *J*=10 Hz, CHC=O), 3.56 (dd, 1, *J*=5 and 10 Hz, CHC=C), 3.76 (s, 3, CH₃O), 4.98 (diffused d, 1, *J*=5 Hz, HC=C). Found: C, 63.82; H, 8.05%. Calcd for C₁₂H₁₈O₄: C, 63.70; H, 8.02%.

4,4-Dimethyl-2-methoxycarbonyl-2-butenolide (9). The diester **4b** (940 mg, 3.55 mmol) was heated in a Claisen

flask to 170 °C for 2 h under N₂ and then distilled to give **9** (456 mg, 76%): bp 90–95 °C/5 Torr; mp 72–73 °C (hexane); IR (Nujol) 1758 (lactone C=O), 1721 (C=O), 1640 cm⁻¹ (C=C); NMR (CDCl₃) δ 1.54 (s, 6, CH₃), 3.87 (s, 3, CH₃O), 8.08 (s, 1, HC=C). Found: C, 56.43; H, 5.79%. Calcd for C₈H₁₀O₄: C, 56.47; H, 5.92%.

4,4-Dimethyl-3-(2-methyl-1-propenyl)-2-(methoxycarbonyl)butanolide (7a). To a THF solution (3 ml) of **9** (100 mg, 0.95 mmol) was added dropwise a THF solution (2 ml) of 2-methyl-1-propenylmagnesium bromide, prepared from 2-methyl-1-propenyl bromide (135 mg, 1.0 mmol) and magnesium (24 mg, 1.0 mmol) at room temperature. After being stirred for 16 h at this temperature, the mixture was quenched with aqueous 10% NH₄Cl, extracted with ether, washed with brine and dried (Na₂SO₄). Removal of the solvents followed by chromatography (SiO₂, hexane–ether, 20/1) gave **7a** (113 mg, 85%): mp 81–81.5 °C (hexane–ether, 10/1). The IR and NMR spectra were identical with those described above.

Pyrocin (7b). A solution of **7a** (40 mg, 0.18 mmol) in DMSO (1.24 ml) containing NaCl (5 mg) and a few drops of water was heated in a sealed tube at 170 °C for 1 h. The mixture was diluted with water and extracted with ether. The extract was washed with brine, dried (Na₂SO₄), and concentrated *in vacuo*. A short-path distillation of the residue gave **7b** (21 mg, 71%): bp 80–85 °C/3 Torr (lit.⁴⁾ bp 130–137 °C/17 Torr). The IR and NMR spectra were identical with those reported.⁴⁾

References

- 1) M. J. Devos, L. Hevesi, P. Bayet, and A. Krief, *Tetrahedron Lett.*, **1976**, 3911 and references cited therein.
- 2) P. Kolsaker and H. J. Storesund, *J. Chem. Soc., Chem Commun.*, **1972**, 375; similar reactions of **2** (X=Cl, Br; Y=H, Ph; R=Me and/or H) with sodium methoxide and/or phenylmagnesium bromide, giving **3**, have been reported: R. Rambaud, *C. R. Acad. Sci.*, **228**, 1595 (1949); R. S. Ratney and English, Jr., *J. Org. Chem.*, **25**, 2213 (1960).
- 3) H. Normant, "Advances in Organic Chemistry," Vol II, Intersciences Publishers Inc., New York, N. Y. (1960), p. 1.
- 4) K. Ueda and M. Mastui, *Agr. Biol. Chem.*, **34**, 1115 (1970) and references cited therein.
- 5) M. S. Kharash and P. O. Tawney, *J. Am. Chem. Soc.*, **63**, 2308 (1941).
- 6) H. O. House, "Modern Synthetic Reactions," 2nd ed, W. A. Benjamin Inc., Menlo Park, California (1972), p. 219.
- 7) A. Takada, T. Sakai, S. Shinohara, and T. Tsuboi, *Bull. Chem. Soc. Jpn.*, **50**, 1133 (1977).
- 8) A. C. Cope, C. M. Hofmann, C. Wyckoff, and E. Hardenbergh, *J. Am. Chem. Soc.*, **63**, 3452 (1941).
- 9) M. J. Mintz and C. Walling, *Org. Synth.*, **49**, 9 (1969).
- 10) W. Herz, *J. Am. Chem. Soc.*, **78**, 1485 (1956).