

Photochemical Reactions of Aromatic Compounds. XXIX.¹⁾ Photochemical and Thermal Cycloreversions of 1,2,2a,8b-Tetrahydrocyclobuta[*a*]naphthalenes


Chyongjin PAC,* Kazuhiko MIZUNO,** and Hiroshi SAKURAI

The Institute of Scientific and Industrial Research, Osaka University, Suita, Osaka 565

(Received June 18, 1977)

Synopsis. Photochemical and thermal cycloreversions of 1,2,2a,8b-tetrahydrocyclobuta[*a*]naphthalenes were investigated. The cycloreversion in the excited singlet state was completely stereospecific; a concerted mechanism was assigned. The benzophenone-sensitized and thermal decomposition gave a mixture of *cis*- and *trans*-olefins, but favored stereoretention. The mechanisms are discussed in terms of singlet and triplet 1,4-biradicals.

The [2+2] cycloreversions of cyclobutane compounds have been of theoretical and mechanical interest in relation to the Woodward-Hoffmann rule²⁾ and 1,4-biradical intermediates.³⁾ We wish to report here on the stereochemical features of photochemical and thermal cycloreversions of 1,2,2a,8b-tetrahydrocyclobuta[*a*]naphthalenes **1a—d**.



The direct or benzophenone-sensitized photolysis of **1a—d** quantitatively gave **2a—d**, **3a—d**, and **4**. The results are listed in Table 1. The olefin formation by

the direct photolysis of **1a, b** occurred with >95% stereospecificities, whereas the cycloreversion in the presence of ferrocene or in neat 1,3-pentadiene was completely stereospecific. Since ferrocene and 1,3-pentadiene are well-known to be good triplet quenchers, the formation of small amounts of **3a, b** in the absence of such compounds probably occurs by means of a triplet mechanism. Similarly, the direct photolysis of **1c** or **1d** in the presence of ferrocene gave only **2c** or **2d** each. In the excited singlet state, therefore, the cycloreversions proceed *via* a $^1\text{S}_0 + ^1\text{S}_0$ concerted mechanism.

On the other hand, the benzophenone-sensitized decomposition of **1a, b** occurred with only a partial stereoretention. In recovered **1a, b**, the epimers could not be detected. The results suggest that triplet biradicals, $^3\text{4c}$ and $^3\text{5t}$, undergo the C₁—C₂ bond rotation in competition with the intersystem crossing to singlet biradicals (Scheme 1).⁴⁾


The thermal decomposition of **1a—d** at 300±10 °C gave **2a—d**, **3a—d**, and **4** in 80% yields, with high stereospecificities. It should be noted that the stereoretention of the *cis* configuration in **1a, c**→**2a, c** predominated over that of the *trans* configuration in **1b, d**→**2b, d**;⁶⁾ usually the retention of the *trans* configuration predominates over that of the *cis* configuration in thermolyses of cyclobutane^{5a,5c,7)} and oxetane⁸⁾ compounds. The C₂—C₃ bond cleavage of singlet biradicals is spin-allowed and favored by the aromatization of the dihydronaphthyl residue. Moreover, the sterically crowded nature of $^1\text{4c}$ reinforces the bond cleavage which overcomes the C₁—C₂ bond rotation, whereas $^1\text{5t}$ undergoes the bond rotation to a significant extent

TABLE 1. PHOTOCHEMICAL AND THERMAL CYCLOREVERSIONS OF **1a—d**^{a)}

		1a		1b		1c		1d	
		2a^{b)}	3a^{c)}	2b^{c)}	3b^{b)}	2c^{d)}	3c^{e)}	2d^{e)}	3d^{d)}
Photolysis	Direct	A ^{f, g)}	95±1	5±1	98±1	2±1	—	—	—
		C ^{f, h)}	95±1	5±1	99	1	—	—	—
		P ⁱ⁾	>99	<1	100	0	—	—	—
		F ^{j)}	100	0	100	0	100	0	100
	Sens ^{k)}	A ^{f, g)}	55±1	45±1	78±1	22±1	—	—	—
Thermolysis ^{l)}	300 °C		95±2	5±2	76±2	24±2	90±3	10±3	80±3
	350 °C		90	10	76	24	—	—	—
	480 °C		88	12	75	25	—	—	—

a) Averaged for each three runs. b) *cis*-1-Phenoxypropene. c) *trans*-1-Phenoxypropene. d) *cis*-1-Methoxypropene. e) *trans*-1-Methoxypropene. f) Extrapolated at zero-time irradiation. g) Acetonitrile solution. h) Cyclohexane solution. i) In neat 1,3-pentadiene; conversion, <10%. j) Acetonitrile solution in the presence of ferrocene; conversion, <10%. k) Sensitized by benzophenone. l) The thermolyses were carried out in the inlet part of a GC-2C machine.

** Present address: Department of Applied Chemistry, College of Engineering, University of Osaka Prefecture, Mozu-Umemachi, Sakai 591.



Scheme 1.

because of the smaller steric repulsion. Alternatively, a biradicaloid transition-state mechanism^{3b,3f} offers an attractive speculation; a concerted mechanism could contribute more to the cycloreversion of **1a, c** than to that of **1b,d**.^{8b,9}

Experimental

Materials. *cis*- and *trans*-endo-1-Phenoxy-2-methyl-1,2,2a,8b-tetrahydrocyclobuta[*a*]naphthalene-8b-carbonitriles **1a** (mp 138.5–139.5 °C) and **1b** (mp 97.5–98.5 °C) were obtained by the photocycloaddition of *cis*- and *trans*-1-phenoxypropenes to 1-naphthonitrile¹⁰ respectively. Similarly, **1c** (mp 65–66 °C) and **1d** (mp 127.5–129 °C) were prepared according to the method reported previously.¹¹ These materials, benzophenone, and ferrocene were recrystallized from methanol. The 1,3-pentadiene was distilled from sodium under a nitrogen atmosphere.

Photolyses. Solutions of **1a–d** (2 mg/cm³) in cyclohexane, acetonitrile, or 1,3-pentadiene were prepared. Solutions of **1a–d** (2 mg/cm³) and ferrocene (2 mg/cm³) or benzophenone (1 mg/cm³) in acetonitrile were also prepared. Each solution (2 cm³) was placed in a Pyrex tube, bubbled with a pure nitrogen stream, and irradiated with a high-pressure mercury arc at 20 ± 2 °C, using a merry-go-round apparatus. In benzophenone-sensitized runs, a hexane solution of naphthalene (13 g/dm³, 1 cm path length) was used as the light filter. During the course of the irradiation, the photolysates were analyzed by VPC at 5-min intervals.

Thermolyses. Pyrex tubes, in which **1a** or **1b** (10–20 mg) was placed, were degassed (<10^{−3} mmHg), sealed, and heated on a metal bath maintained at 300 ± 10 °C for 2–3 h. The decomposition was completed. 1-Naphthonitrile and a mixture of *cis*- and *trans*-1-phenoxypropenes were formed in ca. 90 and 80% yields respectively. More conveniently, the thermolyses were carried out by introducing 0.5–1.0 μl portions of acetonitrile solutions of **1a–d** into a Shimadzu GC-2C gas chromatograph, the inlet part of which was maintained at a constant temperature (300–480 °C).

References


- 1) Part XXVIII: C. Pac, A. Nakasone, and H. Sakurai,

J. Am. Chem. Soc., **99**, 5806 (1977).

- 2) R. B. Woodward and R. Hoffmann, "The Conservation of Orbital Symmetry," Verlag Chemie, GmbH, Weinheim/Bergstr. (1970).

- 3) (a) L. Salem and C. Rowland, *Angew. Chem., Int. Ed. Engl.*, **11**, 92 (1972); (b) R. Hoffmann, S. Swaminathan, B. G. Odell, and R. Gleiter, *J. Am. Chem. Soc.*, **92**, 7091 (1970); (c) L. M. Stephenson and T. A. Gibson, *ibid.*, **94**, 4599 (1972); (d) M. J. S. Dewar, S. Kirschner, H. W. Kollmar, and L. E. Wade, *ibid.*, **96**, 5242, 5246 (1974); (e) R. G. Bergman, in "Free Radicals," ed by J. K. Kochi, John Wiley & Sons, Vol. 1, New York (1973), p. 191; (f) G. Jones, II, *J. Chem. Educ.*, **51**, 175 (1974); (g) L. M. Stephenson and J. I. Brauman, *J. Am. Chem. Soc.*, **93**, 1988 (1971).

- 4) The **6** biradicals could be alternative intermediates, but are energetically less favorable than **4** and **5**.



- 5) (a) H. R. Gerberich and W. D. Walters, *J. Am. Chem. Soc.*, **83**, 4884 (1961); (b) L. A. Paquette and G. L. Thompson, *ibid.*, **94**, 7127 (1972); (c) L. M. Stephenson and T. A. Gibson, *ibid.*, **96**, 5624 (1974).

- 6) Under the conditions of the thermolyses, the isomerization of **2a–d** and **3a–d** is negligible.

- 7) (a) A. T. Cocks, H. M. Frey, and I. D. R. Stevens, *J. Chem. Soc., Chem. Commun.*, **1969**, 458; (b) G. Jones, II and M. H. Williamson, Jr., *J. Am. Chem. Soc.*, **96**, 5617 (1974).

- 8) (a) H. A. J. Carles, *Tetrahedron Lett.*, **1974**, 3425; (b) G. Jones, II and J. C. Staires, *ibid.*, **1974**, 2099.

- 9) G. Jones, II and V. L. Chow, *J. Org. Chem.*, **39**, 1447 (1974).

- 10) K. Mizuno, C. Pac, and H. Sakurai, *J. Chem. Soc., Chem. Commun.*, **1974**, 648.

- 11) K. Mizuno, C. Pac, and H. Sakurai, *J. Org. Chem.*, **42**, 3313 (1977).