

The Reductions of Carbonyl Compounds with Sodium 1-Benzyl-3-carbamoyl-1,4-dihydropyridine-4-sulfinate

Hiroo INOUE,* Ichiro SONODA, and Eiji IMOTO

Department of Applied Chemistry, College of Engineering, University of Osaka Prefecture, Sakai, Osaka 591


(Received September 8, 1978)

Synopsis. It is found that sodium 1-benzyl-3-carbamoyl-1,4-dihydropyridine-4-sulfinate reduces α -keto carbonyl compounds to α -hydroxy carbonyl compounds with the assistance of MgCl_2 and reacts with 9-fluorenone to give an adduct.

Although sodium 1-benzyl-3-carbamoyl-1,4-dihydropyridine-4-sulfinate ($\text{BNA-SO}_2\text{Na}$) is of interest as a reducing agent, not a great deal is yet known about the chemical reactivity of $\text{BNA-SO}_2\text{Na}$. Recently we reported that $\text{BNA-SO}_2\text{Na}$ can undergo reductions of α -halo ketones to the parent ketones¹⁾ and that of acridine to 9,9'-bi(9,10-dihydroacridine)²⁾ in a protic solvent. In analogy with sodium hydroxymethanesulfinate³⁾ and thiourea dioxide,⁴⁾ the carbon-sulfur bond of $\text{BNA-SO}_2\text{Na}$ is cleaved in the reduction process. Our interest in the reactivity of $\text{BNA-SO}_2\text{Na}$ prompted us to examine the capability of $\text{BNA-SO}_2\text{Na}$ for the reductions of the carbonyl compounds. We now wish to report several results giving information on the reactivity of $\text{BNA-SO}_2\text{Na}$ caused by its unique structure.

Results and Discussion

Reactions of $\text{BNA-SO}_2\text{Na}$ with α -Keto Carbonyl Compounds. $\text{BNA-SO}_2\text{Na}$ reacted under nitrogen with benzil (**1**) in 80 vol % aqueous methanol at 25 °C for 12 min to give α -hydroxydeoxybenzoin (**2**) and a thermally unstable adduct (**3**) in isolated yields of 80 and 19% respectively. In this reaction, $\text{BNA-SO}_2\text{Na}$ was converted to 1-benzyl-3-carbamoylpyridinium salt⁵⁾ in a 92% yield, based on **1**. The amount of the salt produced indicates that $\text{BNA-SO}_2\text{Na}$ undergoes a two-electron reduction of **1**. When the reaction mixture was allowed to stand at 25 °C for a prolonged reaction



time (20 h), the yield of **2** (88%) increased with a decrease in that of **3** (5%). On the other hand, the reaction of **1** with $\text{BNA-SO}_2\text{Na}$ in a 0.6 M sodium hydroxide solution (80 vol % aqueous methanol) gave **3** as the only product in a 92% yield.


The addition of MgCl_2 to the reaction system containing **1** and $\text{BNA-SO}_2\text{Na}$ (the molar ratio of MgCl_2 : $\text{BNA-SO}_2\text{Na}$: **1** = 5:2:1) resulted in the selective formation of **2** in a high yield, as Table 1 shows. Furthermore, **3** reacted with MgCl_2 to give **2** in a high yield (87% at 25 °C for 17 h). These facts indicate that **3** is converted to **2** with the assistance of MgCl_2 . The results of the reductions of the other α -keto carbonyl compounds by the $\text{BNA-SO}_2\text{Na}$ - MgCl_2 system are summarized in Table 1. In the absence of MgCl_2 , the yields of the α -hydroxy carbonyl compounds became much lower because of the formation of adducts and unproved by-products. Thus, the $\text{BNA-SO}_2\text{Na}$ - MgCl_2 system is effective as a reducing system for the conversion of α -keto carbonyl compounds to α -hydroxy carbonyl compounds.

TABLE 1. THE YIELDS OF α -HYDROXY CARBONYL COMPOUNDS IN THE REDUCTION OF α -KETO CARBONYL COMPOUNDS WITH THE $\text{BNA-SO}_2\text{Na}$ - MgCl_2 SYSTEM^{a)}

Substance	Time, h	Product (%) ^{b)}
PhCOCOPh	1	PhCOCH(OH)Ph (92)
PhCOCOOEt	17	PhCH(OH)COOEt (70)
PhCOCHO	18	PhCOCH_2OH (55)
$\text{C}_3\text{H}_7\text{COCOC}_3\text{H}_7$	1	$\text{C}_3\text{H}_7\text{COCH(OH)C}_3\text{H}_7$ (61)
	2	 (49)

a) Molar ratio of MgCl_2 : $\text{BNA-SO}_2\text{Na}$: substance = 5:2:1. Concentration of the substance: 0.06 mol/l. 80 vol % aqueous methanol. 25 °C. b) Isolated yield.

Reactions of $\text{BNA-SO}_2\text{Na}$ with Diaryl Ketones. The reaction of $\text{BNA-SO}_2\text{Na}$ with 9-fluorenone (**4**) in 80 vol % aqueous methanol was carried out under conditions similar to those in the case of **1**. With a $\text{BNA-SO}_2\text{Na}$: **4** molar ratio of 2:1, an adduct (**5**) and



9-fluorenone (**6**) were obtained in 30 and 1% yields respectively, with a 62% recovery of **4**. [9,9'-Bi-9H-fluorene]-9,9'-diol (**7**) was not obtained at all. $\text{BNA-SO}_2\text{Na}$ was converted to 1-benzyl-3-carbamoylpyridinium salt in a 72% yield, based on **4**. The yield of the salt was approximately comparable to the recovery

- 1) H. Inoue, N. Inoguchi, and E. Imoto, *Bull. Chem. Soc. Jpn.*, **50**, 197 (1977).
- 2) H. Inoue, I. Sonoda, N. Inoguchi, and E. Imoto, *Bull. Chem. Soc. Jpn.*, **51**, 3097 (1978).
- 3) R. Kerber and W. Gestrich, *Chem. Ber.*, **106**, 798 (1973).
- 4) K. Nakagawa and K. Minami, *Tetrahedron Lett.*, **1972**, 343; J. E. Herz and L. A. Marquez, *J. Chem. Soc., Perkin Trans. 1*, **1973**, 2633.
- 5) This salt was isolated as 1-benzyl-3-carbamoylpyridinium chloride.
- 6) It has been reported recently that benzophenone is reduced to benzhydrol by sodium dithionite in an alkaline medium at 90 °C: J. G. de Vries, T. J. van Bergen, and R. M. Kellogg, *Synthesis*, **1977**, 246.