

Cyclization of Isothiosemicarbazones. IV.¹⁾ Synthesis of the [1,2,4]Triazolo[1,5-*c*]pyrimidine Ring System

Chiji YAMAZAKI

Department of Chemistry, School of Hygienic Sciences, Kitasato University, Kitasato, Sagami-hara, Kanagawa 228

(Received October 29, 1980)

Condensation of isothiosemicarbazones with ethoxymethylenemalononitrile gave 2,3-dihydro[1,2,4]triazolo[1,5-*c*]pyrimidines in moderate to high yields. The 2,3-dihydro compounds were readily oxidized in dimethyl sulfoxide to give the corresponding [1,2,4]triazolo[1,5-*c*]pyrimidines.


Isothiosemicarbazones (**1**) are polyfunctional nucleophiles for the reaction with α -halo carbonyl compounds leading to the formation of nitrogen-containing heterocycles.^{1,2)} The internal nitrogen (N-2) of **1** preferentially attacks the halogen-bearing carbon of the halo compound to displace the halide and is considered to be a softer nucleophilic center³⁾ than the terminal nitrogen (N-4). On the other hand, N-4 invariably becomes attached to the carbonyl carbon whether it is unsubstituted²⁾ or monosubstituted.¹⁾ In the present work, an attempt was made to initiate the cyclization of **1** at N-4 by the reaction with ethoxymethylenemalononitrile (**2**) and to examine the formation of a six-membered heterocycle. If **1** reacts with **2** at N-4, 3,4-disubstituted isothiosemicarbazone (**3**) might initially be formed, undergoing intramolecular cycloaddition to give a 1-(alkylidene- or benzylideneamino)-5-cyano-6-imino-2-mercapto-1,6-dihydropyrimidine derivative (**4**) in a similar way to that for pyrimidine formation from *S*-alkylisothioureas.⁴⁾

It was found that the reaction of **1** with **2** leads directly to the formation of 2,3-dihydro[1,2,4]triazolo[1,5-*c*]pyrimidines (**5**) and that, when R³ is hydrogen, the dihydro compound (**5**) easily undergoes oxidation to [1,2,4]triazolo[1,5-*c*]pyrimidines (**6**) (Scheme 1). Among many condensed pyrimidine derivatives, few examples are known for the preparation of [1,2,4]-

triazolo[1,5-*c*]pyrimidines,⁵⁾ no reports having been published on a one-step synthesis of the bicyclic pyrimidines from an open-chain, flexible molecule. This paper deals with the preparation and structure of a new series of 2,5-disubstituted and 2,2,5-trisubstituted 8-cyano-2,3-dihydro[1,2,4]triazolo[1,5-*c*]pyrimidines (**5a—n**) and 2,5-disubstituted 8-cyano[1,2,4]triazolo[1,5-*c*]pyrimidines (**6a—i**).

Results and Discussion

The reaction was performed by allowing a solution of **1** and **2** in a 1 : 1.15 molar ratio in benzene to stand at room temperature (Procedure A). Except for **5i** and **5n**, most of **5** crystallized out of the reaction mixture in a substantially pure form, yields depending in part on the reaction period. For example, **5a** was obtained in 97% yield after the reaction mixture had been left to stand for one week, but the yield decreased to 82 and 56% with the elapse of 18 and 3 h, respectively (Table 1). Prolonged periods of time, however, may cause discoloration of the product, analytically pure compounds being obtained within no more than 1 d. The reaction can also be performed by heating a mixture of **1** and **2** (1 : 1.15) at 95–100 °C in the absence of solvent (Procedure B). The melted mixture rapidly solidifies within a few minutes, giving **5** in 55–86% yields after



Scheme 1. The reaction of isothiosemicarbazones with ethoxymethylenemalononitrile.

TABLE 1. 2,5-DI- AND 2,2,5-TRISUBSTITUTED 8-CYANO-2,3-DIHYDRO[1,2,4]TRIAZOLO[1,5-*c*]PYRIMIDINES

Compd No.	Yield/%	Procedure	Mp/°C	Formula	Found (Calcd) (%)		
					C	H	N
5a	56—97	A	181—182(dec) ^{a)}	C ₁₃ H ₁₁ N ₅ S	57.93 (57.98)	4.19 4.12	26.33 26.01)
5b	76—82	A	162—175 ^{b,c)}	C ₁₃ H ₁₀ ClN ₅ S	51.38 (51.40)	3.28 3.32	23.26 23.06)
5c	92	A	159—159.5 ^{a)}	C ₁₃ H ₁₀ ClN ₅ S	51.43 (51.40)	3.30 3.32	23.23 23.06)
5d	86	B	191 ^{d)}	C ₁₃ H ₉ Cl ₂ N ₅ S	46.18 (46.17)	2.65 2.68	20.98 20.71)
5e	86	A	162—163 ^{b)}	C ₁₄ H ₁₃ N ₅ OS	56.32 (56.18)	4.29 4.38	23.58 23.40)
5f	64 ^{e)} —87	A	193—194 ^{a,f)}	C ₁₄ H ₁₃ N ₅ OS	56.16 (56.18)	4.35 4.38	23.44 23.40)
5g	63—70	A	141—170 ^{a,c)}	C ₁₆ H ₁₇ N ₅ OS	58.73 (58.72)	5.23 5.20	21.46 21.41)
5h	71—80	A	151 ^{b)}	C ₁₆ H ₁₅ N ₅ OS	59.04 (59.07)	4.62 4.65	21.61 21.53)
5i	80	B	174—175 ^{d)}	C ₁₅ H ₁₁ Cl ₂ N ₅ S	49.48 (49.46)	3.00 3.04	19.49 19.23)
5j	70	A,B	160.5—161.5 ^{b)}	C ₁₉ H ₁₅ N ₅ S	65.97 (66.07)	4.53 4.38	20.51 20.28)
5k	77	g)	180.5—181.5 ^{b)}	C ₉ H ₁₁ N ₅ S	48.91 (48.86)	4.99 5.01	31.74 31.66)
5l	70	i)	157(dec) ^{j)}	C ₁₄ H ₁₃ N ₅ S	59.39 (59.35)	4.66 4.63	24.99 24.72)
5m	60	A	176 (dec) ^{k)}	C ₁₆ H ₁₆ ClN ₅ OS	52.83 (52.82)	4.98 4.99	19.27 19.25)
5n	86	B	200.5—201.5(dec) ^{h)}	C ₁₉ H ₁₅ N ₅ S	65.90 (66.07)	4.37 4.38	20.39 20.28)

a) An analytically pure compound obtained without recrystallization. b) Light yellow needles from EtOH-MeCN (1 : 1). c) Homogeneous on TLC (Kieselgel 60 F₂₅₄ plate, CHCl₃ containing 5% by volume of MeOH), partial oxidation during the course of melting point measurement presumably responsible for the wide melting range. d) Yellow prisms from EtOH-MeCN (1 : 1). e) MeCN used in place of benzene. f) Substantially converted into **6f** on recrystallizing from EtOH-MeCN (1 : 1). g) Refluxed in benzene for 30 min. h) Light yellow needles from 80% aqueous EtOH. i) Refluxed in EtOH for 30 min. j) An analytically pure compound obtained by drying *in vacuo* the initially formed product solvated with one molecule of EtOH. k) Yellow needles solvated with one molecule of EtOH.

TABLE 2. 2,5-DISUBSTITUTED 8-CYANO[1,2,4]TRIAZOLO[1,5-*c*]PYRIMIDINES

Compd No.	Yield/%	Method	Mp/°C	Formula	Found (Calcd) (%)		
					C	H	N
6a	57	A	256 ^{a)}	C ₁₃ H ₉ N ₅ S	58.44 (58.42)	3.42 3.39	26.46 26.21)
6b	50	B	226—226.5 ^{a)}	C ₁₃ H ₈ ClN ₅ S	51.70 (51.74)	2.61 2.67	23.32 23.21)
6c	61	A	213—214 ^{b)}	C ₁₃ H ₈ ClN ₅ S	51.72 (51.74)	2.71 2.67	23.31 23.21)
6d	82	B	261—262 ^{c)}	C ₁₃ H ₇ Cl ₂ N ₅ S	46.35 (46.49)	2.13 2.10	21.03 20.83)
6e	50	A	192—193 ^{b)}	C ₁₄ H ₁₁ N ₅ OS	56.62 (56.55)	3.60 3.73	23.72 23.56)
6f	53	A	238—239 ^{d)}	C ₁₄ H ₁₁ N ₅ OS	56.32 (56.55)	3.64 3.73	23.88 23.56)
6g	50	B	191.5—192 ^{e)}	C ₁₆ H ₁₅ N ₅ OS	58.82 (59.07)	4.62 4.65	21.42 21.53)
6i	56	C	140.5—142.5 ^{f)}	C ₁₅ H ₉ Cl ₂ N ₅ S	49.65 (49.73)	2.58 2.50	19.19 19.34)

a) Colorless needles from EtOH-pyridine (1 : 1). b) Colorless needles from benzene-EtOH (1 : 1). c) Pale yellow needles from EtOH-pyridine (1 : 1). d) Pale yellow needles from EtOH-MeCN (1 : 1). e) Colorless needles from 80% EtOH. f) Colorless prisms from benzene-EtOH (1 : 1).

TABLE 3. PARTIAL SPECTRAL DATA ON 2,5-DISUBSTITUTED AND 2,2,5-TRISUBSTITUTED 8-CYANO-2,3-DIHYDRO[1,2,4]TRIAZOLO[1,5-c]PYRIMIDINES

Compd No.	IR (KBr) $\tilde{\nu}_{\text{CN}}/\text{cm}^{-1}$	Mass spectra, m/e (rel int)			^1H NMR spectra δ/ppm (from TMS in DMSO- d_6)				
		M^+	$\text{M}^+ - \text{R}^2$	$\text{M}^+ - \text{R}^1$	SCH_2	SCH_3	H-2 (J/Hz)	H-3 (J/Hz)	H-7
5b	2220	303(17)	302(28)	192(100)	—	2.50	6.52(9.3)	7.49(9.3)	8.05
5c	2220	303(30)	302(56)	192(100)	—	2.53	6.28(9.7)	7.30(9.7)	8.04
5e	2230	299(35)	298(100)	192(88)	—	2.53	6.42(9.1)	7.20(9.1)	8.06
5f	2225	299(55)	298(100)	192(69)	—	2.51	6.20(10.0)	7.08(10.0)	8.04
5g	2215	327(66)	326(100)	220(50)	3.13 ^{a)}	—	6.19(9.5)	7.07(9.5)	8.01
5h	2230	325(15)	324(37) ^{b)}	218(34)	3.85 ^{c)}	—	6.20(9.8)	7.10(9.8)	8.04
5i	2220	263(9)	362(7)	218(100)	3.90 ^{c)}	—	7.15(9.4)	7.85(9.4)	8.13
5j	2215	345(15)	344(11) ^{d)}	268(11)	4.49	—	6.27(9.5)	7.23(9.5)	8.10
5k	2225	221(10)	206(100)	—	—	2.50	—	6.47	8.00
5k-d_6^{e)}		227(6)	209(100)	—	—	2.50	—	6.46	7.97
5l	2210	283(5)	268(100)	206(41)	—	2.50	—	6.90	8.02
5l-d_3^{f)}		286(2)	268(100)	209(16)	—	2.51	—	6.90	8.02
5m	2220	317(4)	302(100)	206(42)	—	2.52	—	7.22	7.71
5n	2210	345(5)	268(100)	—	—	2.55	—	7.18	8.07

a) Triplet, $J=6.9$ Hz. b) Base peak allyl cation (m/e 41). c) Doublet, $J=6.2$ Hz. d) Base peak tropylium ion (m/e 91). e) $\text{R}^1=\text{R}^2=\text{CD}_3$. f) $\text{R}^2=\text{CD}_3$.


TABLE 4. PARTIAL SPECTRAL DATA ON 2,5-DISUBSTITUTED 8-CYANO[1,2,4]TRIAZOLO[1,5-c]PYRIMIDINES

Compd No.	IR (KBr) $\tilde{\nu}_{\text{CN}}/\text{cm}^{-1}$	Mass spectra, m/e (rel int)			^1H NMR spectra δ/ppm (from TMS in CF_3COOD)		
		M^+	$\text{M}^+ - \text{R}^1\text{CN}$	$\text{R}^1\text{C}\equiv\text{NH}(\text{D})$	SCH_2	SCH_3	H-7
6c	2225	301(24)	164(69)	138(9)	—	3.01	9.00
6d	2240	335(83)	164(100)	172(24)	—	3.00	8.98
6e	2240	297(85)	164(49)	134(6)	—	3.01	9.07
6e-d_3^{a)}		300(61)	167(56)	135(9)	—	—	9.06
6f	2230	297(100)	164(21)	134(29)	—	3.01	9.03
6f-d_3^{a)}		300(100)	167(27)	135(27)	—	—	9.03
6g	2240	325(100)	192(6)	134(31)	3.64 ^{b)}	—	9.00

a) $\text{R}^3=\text{CD}_3$. b) Triplet, $J=7.1$ Hz.


washing with appropriate solvents. Procedure B was satisfactory for **1i** and **1n** from which no **5** was obtained by Procedure A. With ketone isothiosemicarbazones **1k** and **1l**, refluxing in benzene or EtOH gave good results. The reaction of aliphatic aldehyde isothiosemicarbazones (**1**: $\text{R}^1=\text{Et}$ or $n\text{-Pr}$, $\text{R}^2=\text{H}$, $\text{R}^3=\text{Me}$ or $p\text{-ClC}_6\text{H}_4\text{CH}_2$) with **2** under these conditions gave a complex mixture from which no expected product was isolated. Attempts to cyclize benzaldehyde 4-methyl- or 4-phenylthiosemicarbazone were unsuccessful with total recovery of the starting thiosemicarbazone.

In view of the high yields of hindered compounds (**5d**, **5i** and **5n**), no steric factor may be involved in the 2,3-dihydro-1,2,4-triazole ring formation. Although no intermediate has been detected on the NMR time-scale in the reaction carried out in C_6D_6 , $\text{C}_5\text{D}_5\text{N}$ or dimethyl- d_6 sulfoxide (DMSO- d_6), the reaction should proceed through a 3,4-disubstituted isothiosemicarbazone (**3**). In line with the proposed intermediacy, the reaction of **1a** with ethyl ethoxymethylenecyanoacetate gave the open-chain product **7**, the structure of which was established by elemental analysis and spectral data, particularly by the large coupling constant between N^4H and the methine proton ($J=13.2$ Hz).⁶⁾ If a



nucleophilic attack of N-2 in **3** on the cyano carbon had occurred, **4** would have been formed which might consecutively undergo intramolecular addition of the 6-imino group to the azomethine double bond at the 1-position giving **5**, although the ring closure of **4** to **5** is disfavored according to the rules proposed by Baldwin.⁷⁾ A mechanistic study on the ring formation is now being continued with related compounds including **7**.

Intramolecular hydrogen bonding between the N^4H and the carbonyl oxygen in **7** might stabilize the



8: $\text{R}^4=\text{CN}$

9: $\text{R}^4=\text{COOEt}$

conformation **8** and prevent N-2 from approaching the cyano group. The resulting insusceptibility to cyclization makes it possible to isolate **7** at the open chain stage. In the infrared spectrum of **7**, N⁴-H and the carbonyl stretching frequencies (ν NH 3195 and ν C=O 1690 cm⁻¹) observed at 4×10^{-3} mol dm⁻³ in carbon tetrachloride were comparable with those of **9** (ν NH 3195 and ν C=O 1697 cm⁻¹) obtained at the same concentration in carbon tetrachloride. The diester **9**, however, exhibited an additional carbonyl band at a higher frequency (ν C=O 1730 cm⁻¹). This band can unambiguously be assigned to the free stretching vibration of the remaining ethoxycarbonyl group which is not involved in the internal bonding. The anomalously downfield resonance (lower than δ 12 ppm) of the N⁴H protons of **7** and **9** in chloroform-*d* might also account for the hydrogen bond.⁸⁾

Oxidation of **5** (R²=H) to the corresponding [1,2,4]-triazolo[1,5-*c*]pyrimidines (**6**) was performed simply by leaving a solution of **5** in DMSO in an open vessel to stand at ambient temperature (Method A), a relatively insoluble **6** crystallizing out of the solution. In another procedure, the reaction between **1** and **2** was conducted in the absence of solvent, and the crude **5**, without isolation, was dissolved in DMSO to give **6** in moderate overall yields (Method B). The oxidation was also carried out with iron(III) chloride in aqueous acetic acid with poor yields of **6** (Method C), although **6i** was obtained only by this procedure in satisfactory purity. Some of **5** in which R¹ is unsubstituted or para-substituted phenyl group were particularly susceptible to oxidation and could not be recrystallized without contamination with the oxidized product.

The structures of the compounds **5** and **6** have been established on the basis of spectral measurements and elemental analyses. Partial spectral data are given in Tables 3 and 4 together with those for certain deuterated compounds. In mass spectrometry, molecular ions were obtained for all the compounds **5a—n**, with the intensity depending upon steric crowding at the 2-position (2—65%). Abundant ions M⁺-R¹ and M⁺-R² observed in the spectra of **5** characterized the 2,3-dihydro-1,2,4-triazole structure. The fragmentation pathways were confirmed by the mass spectra of deuterated compounds of **5a**, **5k**, and **5l**, in which R² is D, CD₃, and CD₃, respectively. The dehydrogenated compounds **6** showed a much more intense peak for M⁺ ion than that of **5**, indicating their extended conjugation systems (24—100%). A characteristic fragment ion R¹C≡NH⁺ (6—31%) observed in the spectra of **6** should be formed by hydrogen transfer from R³ group to N-3 probably through a six-membered transition state as evidenced by the spectra of trideuterated compounds of **6b**, **6e**, and **6f** (R³=CD₃), confirming the proposed arrangements of **5** and **6**. In NMR spectroscopy, two protons H-2 and H-3 on the dihydro-1,2,4-triazole ring of **5** (R²=H) appear as two AB-type doublets with coupling constants of 9.1—10.0 Hz. The peak assignment is based on the observation that the upfield resonance collapses to a singlet while the downfield one disappears on addition of methanol-*d*₁. The spectrum of 2-deuterated **5a** (R²=D) lacks the upfield resonance,

exhibiting the H-3 proton as a singlet. The chemical shift values of the H-2 proton in **5** are *ca.* 2.0 ppm higher than those of the azomethine proton in the corresponding isothiosemicarbazones, reflecting rehybridization of the carbon atom (C-2) from sp² in **1** to sp³ in **5**. The 2-methyl protons of **5k—m** (R²=Me) resonate at 0.6—0.7 ppm higher than those of the corresponding **1k—m**, also indicating the rehybridization of C-2. In line with the rehybridization of C-2, the anisotropic deshielding (0.43 ppm in DMSO-*d*₆) of phenyl protons ortho to the azomethine double bond in **1a** disappears in **5a** which exhibits only a single signal for the phenyl protons at δ 7.36. The effect recommences in **6a** in which the ortho protons (H^o) of 2-phenyl group are deshielded by 0.49 ppm (trifluoroacetic acid-*d*) relative to the remaining aromatic protons (H^{m,p}) due to the resonance interaction with the heteroaromatic ring system.⁹⁾ Substantially constant values of the chemical shifts of H-7 proton in **5** or **6** when R¹-R³ were widely changed in structure are in line with the assigned structures.

Experimental

General. Melting points were taken in open glass capillaries and are uncorrected. Infrared spectra were recorded on a Hitachi EPI-G2 or 260-30 spectrophotometer, and calibrated by comparison with a standard polystyrene film sample. Proton nuclear magnetic resonance spectra were obtained with a Hitachi R-24 spectrometer at 60 MHz. Unless otherwise stated, chemical shifts are given in parts per million (δ scale) downfield from internal tetramethylsilane (TMS). Solvents used are DMSO-*d*₆ for 2,3-dihydro-1,2,4-triazolo compounds (**5**) and trifluoroacetic acid-*d* for 1,2,4-triazolo compounds (**6**). The mass spectra (75 eV) were recorded on a JMS-D100 mass spectrometer. Ethoxymethylenemalononitrile (**2**) (Aldrich Chemical Co. Inc.) was used after removal of insoluble substances in benzene at room temperature.

Isothiosemicarbazones. The compounds **1a—n** were prepared by the method reported.⁸⁾ *S*-(Methyl-*d*₃)isothiosemicarbazones were obtained by using methyl-*d*₃ iodide in place of methyl iodide in the usual procedure. Other deuterated isothiosemicarbazones were similarly prepared from the corresponding deuterated carbonyl compounds, acetone-*d*₆ (CD₃COCOD₃), acetophenone- α,α,α -*d*₃ (C₆H₅COCOD₃), and benzaldehyde-*formyl-d* (C₆H₅CDO). New compounds are: 2,6-dichlorobenzaldehyde *S*-allylisothiosemicarbazone (**1i**): pale yellow prisms (from *i*-Pr₂O), mp 99—101 °C; NMR (CDCl₃) δ =3.76 (2H, dt, *J*=6.5 and 0.8 Hz, SCH₂), 5.5—6.3 (5H, m, CH₂=CH and NH₂), 7.06—7.45 (3H, m, aromatic), 8.62 (1H, s, CH=N). Found: C, 45.54; H, 3.79; N, 14.83%. Calcd for C₁₁H₁₁Cl₂N₃S: C, 45.84; H, 3.85; N, 14.58%. *p*-Chloroacetophenone *S*-methylisothiosemicarbazone (**1m**): colorless needles (from aqueous EtOH), mp 231 °C (HBr salt); NMR (CDCl₃) (free base): δ =2.38 (3H, s, CCH₃), 2.47 (3H, s, SCH₃), 5.41 (2H, bs, NH₂), 7.30 (2H, d, *J*=8.3 Hz, aromatic), 7.75 (2H, d, *J*=8.3 Hz, aromatic). Found: C, 37.30; H, 3.96; N, 13.30%. Calcd for C₁₀H₁₃BrClN₃S: C, 37.22; H, 4.06; N, 13.02%. Benzophenone *S*-methylisothiosemicarbazone (**1n**): colorless fine needles (from EtOH), mp 135.5—136 °C; NMR (CDCl₃) δ =2.20 (3H, s, SCH₃), 5.35 (2H, bs, NH₂), 7.35 (5H, s, aromatic), 7.21—7.73 (5H, m, aromatic). Found: C, 66.95; H, 5.57; N, 15.91%. Calcd for C₁₅H₁₅N₃S: C, 66.90; H, 5.61; N, 15.61%.

8-Cyano-2,3-dihydro-5-methylthio-2-phenyl[1,2,4]triazolo[1,5-*c*]pyrimidine (5a) (A Typical Example of Procedure A). A solution

of **1a** (0.19 g, 1 mmol) and **2** (0.14 g, 1.15 mmol) in benzene (1 ml) was allowed to stand at room temperature. After *ca.* 30 min, yellow prisms began to separate and the reaction was allowed to proceed for 18 h. The crystals were collected by filtration, washed with benzene, and air-dried, giving 0.22 g (82%) of analytically pure **5a**; mp 181–182 °C (dec); IR (KBr) 2220 (CN) cm^{-1} ; NMR δ =2.53 (3H, s, SCH_3), 6.26 (1H, d, J =9.7 Hz, H-2), 7.24 (1H, d, J =9.7 Hz, H-3), 7.36 (5H, s, aromatic), 8.04 (1H, s, H-7); MS, m/e (rel intensity), 269 (29) (M^+), 268 (56) ($\text{M}^+ - \text{H}$), 192 (100) ($\text{M}^+ - \text{C}_6\text{H}_5$).

The 2-deuterio compound of **5a** was similarly obtained from benzaldehyde-*formyl-d* *S*-methylisothiosemicarbazone [$\text{Ph-GD=N-N=C(SMe)NH}_2$] as yellow prisms, mp 181 °C (dec); NMR δ =2.51 (3H, s, SCH_3), 7.20 (1H, s, H-3), 7.34 (5H, s, aromatic), 8.03 (1H, s, H-7); MS, m/e (rel intensity), 270 (25) (M^+), 268 (40) ($\text{M}^+ - \text{D}$), 193 (100) ($\text{M}^+ - \text{C}_6\text{H}_5$).

8-Cyano-2-(2,6-dichlorophenyl)-2,3-dihydro-5-methylthio[1,2,4]-triazolo[1,5-c]pyrimidine (5d) (A Typical Example of Procedure B). A mixture of **1d** [hydriodide, mp 200–200.5 °C (dec) (lit.¹⁰) mp 209–211 °C (dec)]; Found: C, 27.88; H, 2.60; N, 11.00%. Calcd for $\text{C}_9\text{H}_{10}\text{Cl}_2\text{IN}_3\text{S}$: C, 27.71; H, 2.58; N, 10.77%] (0.26 g, 1 mmol) and **2** (0.14 g, 1.15 mmol) was heated on a boiling water bath to melt the solids. The liquid formed, still of low viscosity, was thoroughly agitated in order to confirm complete homogeneity. The mixture rapidly solidified within 30 s and heating was continued for 5 min to allow the by-product EtOH to evaporate. After cooling, the solid was triturated with acetone, collected by filtration, washed with acetone, and air-dried to give 0.29 g (86%) of **5d** as yellow crystalline powder, mp 175–178 °C. This was recrystallized from an EtOH–MeCN mixture (1 : 1 by volume) giving sparkling yellow prisms, mp 191 °C; IR (KBr) 2220 (CN) cm^{-1} ; NMR δ =2.55 (3H, s, SCH_3), 7.14 (1H, d, J =9.7 Hz, H-2), 7.43 (3H, s, aromatic), 7.79 (1H, d, J =9.7 Hz, H-3), 8.09 (1H, s, H-7); MS m/e (rel intensity), 337 (8) (M^+), 336 (7) ($\text{M}^+ - \text{H}$), 192 (100) ($\text{M}^+ - \text{C}_6\text{H}_5\text{Cl}_2$).

When Procedure A was applied to the preparation of **5d**, a product, mp 189–190 °C, was obtained in 65% yield after standing for 6 h.

8-Cyano-5-methylthio-2-phenyl[1,2,4]triazolo[1,5-c]pyrimidine (6a) (A Typical Example of Method A; One-step Synthesis from Isothiosemicarbazones). A mixture of **1a** (0.1 g, 0.52 mmol) and **2** (0.07 g, 0.58 mmol) was heated in an open vessel on a boiling water bath with constant shaking, the ethanol formed being allowed to evaporate for 5 min. The resulting crystalline mass was dissolved still hot in DMSO (1 ml) and the solution was allowed to stand at room temperature for 1 d. The separated crystals were collected by filtration, washed with DMSO and water and then air-dried, giving analytically pure **6a** in 57% overall yield as colorless needles, mp 255.5–256 °C. Recrystallization from an EtOH–pyridine mixture (1 : 1 by volume) gave colorless needles, mp 256 °C; IR (KBr) 2220 (CN) cm^{-1} ; NMR δ =3.00 (3H, s, SCH_3), 7.73 (3H, m, $\text{H}^{m,p}$ of phenyl), 8.22 (2H, m, H^o of phenyl), 9.02 (1H, s, H-7); MS, m/e (rel intensity), 267 (100) (M^+), 164 (85) ($\text{M}^+ - \text{C}_6\text{H}_5$), 104 (31) ($\text{C}_6\text{H}_5\text{C}\equiv\text{NH}$), 103 (27).

2-(o-Chlorophenyl)-8-cyano-5-methylthio[1,2,4]triazolo[1,5-c]pyrimidine (6b) (A Typical Example of Method B). A solution of **5b** (0.1 g) in DMSO (1 ml) was allowed to stand at room temperature for 1 d and separated crystals were collected by filtration, washed with DMSO and water and then air-dried, giving **6b** in 50% yield as colorless needles, mp 226–226.5 °C. Recrystallization from an EtOH–pyridine mixture (1 : 1 by volume) did not change the mp or appearance. IR (KBr) 2230 (CN) cm^{-1} ; NMR δ =3.00 (3H, s, SCH_3), 7.50–8.07 (4H, m, aromatic), 9.04 (1H, s, H-7); MS, m/e (rel intensity),

301 (66) (M^+), 266 (53) ($\text{M}^+ - \text{Cl}$), 164 (24) ($\text{M}^+ - \text{ClC}_6\text{H}_4\text{-CN}$), 138 (29) ($\text{ClC}_6\text{H}_4\text{C}\equiv\text{NH}$), 137 (100).

The 5-methylthio-*d*₃ compound of **6b** was obtained from the corresponding trideuterated **5b** by Method B as colorless needles, mp 225–226 °C; NMR δ =7.48–8.07 (4H, m, aromatic), 9.04 (1H, s, H-7); MS, m/e (rel intensity), 304 (100) (M^+), 269 (84) ($\text{M}^+ - \text{Cl}$), 167 (95) ($\text{M}^+ - \text{ClC}_6\text{H}_4\text{CN}$), 139 (33) ($\text{ClC}_6\text{H}_4\text{C}\equiv\text{ND}$).

5-Allylthio-8-cyano-2-(2,6-dichlorophenyl)[1,2,4]triazolo[1,5-c]pyrimidine (6i) (Method C).

To a solution of **5i** (0.09 g, 0.25 mmol) in AcOH (2 ml) was added 0.5 ml of a solution containing 1 mmol/ml of iron(III) chloride in 60% aqueous AcOH and the mixture was allowed to stand at room temperature with occasional agitation for 10 d. After being diluted with water, separated crystals (0.07 g), mp 127–140 °C, were recrystallized from a benzene–EtOH mixture (1 : 1 by volume) to give 0.05 g (56%) of **6i** as colorless prisms, mp 140.5–142.5 °C; IR (KBr) 2230 (CN) cm^{-1} ; NMR δ =4.26 (2H, d, J =6.6 Hz, SCH_2), 5.26–6.34 (3H, m, CH=CH_2), 7.55 (3H, s, aromatic), 8.95 (1H, s, H-7); MS, m/e (rel intensity), 361 (61) (M^+), 190 (80) ($\text{M}^+ - \text{Cl}_2\text{C}_6\text{H}_3\text{CN}$), 172 (25) ($\text{Cl}_2\text{C}_6\text{H}_3\text{C}\equiv\text{NH}$), 121 (100), 41 (77) (allyl cation).

Benzaldehyde 3-Methyl-4-[2-cyano-2-(ethoxycarbonyl)vinyl]isothiosemicarbazone (7).

A mixture of **1a** (0.19 g, 1 mmol), ethyl ethoxymethylenecyanoacetate (0.19 g, 1.15 mmol), and benzene (0.5 ml) was heated at 70 °C for 1 h. On cooling, the separated solid was collected by filtration, washed with EtOH and air-dried, giving 0.27 g (84%) of **7** as pale yellow crystalline powder, mp 140–143 °C. Recrystallization twice from EtOH provided pale yellow needles, mp 145 °C; IR (CCl_4) 3195 (NH), 2220 (CN), 1690 (C=O) cm^{-1} ; NMR (CDCl_3) δ =1.39 (3H, t, J =7.2 Hz, CH_2CH_3), 2.60 (3H, s, SCH_3), 4.34 (2H, q, J =7.2 Hz, CH_2CH_3), 7.43 (3H, m, $\text{H}^{m,p}$ of phenyl), 7.69 (1H, d, J =13.2 Hz, NH=CH), 7.90 (2H, m, H^o of phenyl), 8.47 (1H, s, CH=N), 12.40 (1H, d, J =*ca.* 13 Hz, NH-CH).

Found: C, 56.97; H, 5.09; N, 17.52%; M^+ , 316. Calcd for $\text{C}_{15}\text{H}_{16}\text{N}_4\text{O}_2\text{S}$: C, 56.96; H, 5.10; N, 17.71%; M , 316.

Benzaldehyde 3-Methyl-4-[2,2-bis(ethoxycarbonyl)vinyl]isothiosemicarbazone (9).

A mixture of **1a** (0.19 g, 1 mmol) and diethyl ethoxymethylenemalonate (0.22 g, 1 mmol) in benzene (0.5 ml) was refluxed for 4 h, the solvent being evaporated. Recrystallization of the crystalline residue from EtOH gave **9** as pale yellow prisms (0.27 g, 74%), mp 99–100 °C; NMR (CDCl_3) δ =1.31 (3H, t, J =7.0 Hz, CH_2CH_3), 1.36 (3H, t, J =7.0 Hz, CH_2CH_3), 2.60 (3H, s, SCH_3), 4.18 (2H, q, J =7.0 Hz, CH_2CH_3), 4.29 (2H, q, J =7.0 Hz, CH_2CH_3), 7.41 (3H, m, $\text{H}^{m,p}$ of phenyl), 7.88 (2H, m, H^o of phenyl), 8.22 (1H, d, J =13.3 Hz, CH-NH), 8.42 (1H, s, CH=N), 12.28 (1H, d, J =13.3 Hz, CH-NH).

Found: C, 56.11; H, 5.83; N, 11.67%; M^+ , 363. Calcd for $\text{C}_{17}\text{H}_{21}\text{N}_5\text{O}_4\text{S}$: C, 56.19; H, 5.83; N, 11.57%; M , 363.

The author wishes to thank Miss Yoko Ishizuka for assistance in the preparation of compounds.

References

- 1) Part III: C. Yamazaki, *Bull. Chem. Soc. Jpn.*, **53**, 3289 (1980).
- 2) C. Yamazaki, *Bull. Chem. Soc. Jpn.*, **51**, 1846 (1978); *Tetrahedron Lett.*, **1978**, 1295.
- 3) Exclusive methylation on the internal nitrogen with methyl iodide has been observed for isothiosemicarbazones⁸ and also for benzaldehyde amidinohydrazone [W. G. Finnegan, R. A. Henry, and G. B. L. Smith, *J. Am. Chem. Soc.*, **74**, 2981

(1952)].

4) S. G. Cottis and H. Tieckelmann, *J. Org. Chem.*, **26**, 79 (1961).

5) Y. Tamura, J. Kim, and M. Ikeda, *J. Heterocycl. Chem.*, **12**, 107 (1975).

6) K. Saito, I. Hori, M. Igarashi, and H. Midorikawa, *Bull. Chem. Soc. Jpn.*, **47**, 476 (1974).

7) J. E. Baldwin, *J. Chem. Soc., Chem. Commun.*, **1976**, 734;

736.

8) C. Yamazaki, *Can. J. Chem.*, **53**, 610 (1975).

9) L. A. Lee and J. W. Wheeler, *J. Org. Chem.*, **37**, 348 (1972); R. N. Butler, *Can. J. Chem.*, **51**, 2315 (1973); L. G. Tensmeyer and C. Ainsworth, *J. Org. Chem.*, **31**, 1878 (1966).

10) W. J. Houlihan and R. E. Manning, Ger. Offen. 1 902449 (1969); *Chem. Abstr.*, **71**, 123963q (1969).
