

Syntheses of *N*-Substituted 3,3,4-Triaryl- and 3,3,4,4-Tetraphenyl-2-azetidinones

Kailash Nath MEHROTRA* and Surendra Bahadur SINGH

Department of Chemistry, Banaras Hindu University, Varanasi-221005, India

(Received June 28, 1980)

The reactions of 2-diazo-1,2-diphenylethanone with Schiff's bases $\text{Ar}-\text{CH}=\text{N}-\text{R}$ gave new substituted 2-azetidinones together with 1,1',4,4'-tetraphenyl-2,2'-azinodiethanone. 2-Diazo-1,2-diphenylethanone reacts with *N*-benzhydrylidene-*N'*-phenylurea to give 1-(phenylcarbamoyl)-3,3,4,4-tetraphenyl-2-azetidinone.

The cycloaddition of diphenylketene with anils is known to form 2-azetidinones.¹⁾ Diphenylketene, in these cases, has been generated either thermally by dehydrohalogenation of suitable acid chlorides with tertiary bases²⁾ or photochemically from diazo ketone.³⁾ On the other hand, diphenylketene has been known to add on carbon-oxygen double bond in α,β -unsaturated ketones to give β -lactones.⁴⁾ We now report the cycloaddition of diphenylketene, generated *in situ* by thermal decomposition of 2-diazo-1,2-diphenylethanone (**1**), with Schiff's bases **2a—g** leading to *N*-alkyl-2-azetidinones **3a—g** in fair yields. The reaction of diphenylketene with *N*-benzhydrylidene-*N'*-phenylurea (**5**), having three reactive sites ($\text{C}=\text{N}$, $\text{C}=\text{O}$, and NH), has shown that $\text{C}=\text{N}$ group is attacked by diphenylketene in preference to either $\text{C}=\text{O}$ group or NH group. The products, **3a—g**, show extraordinary stability amongst β -lactams⁵⁾ towards common degradative reagents.

Results and Discussion


An equimolecular mixture of 2-diazo-1,2-diphenylethanone (**1**) and *N*-benzylideneisopropylamine (**2a**: $\text{Ar}=\text{C}_6\text{H}_5$, $\text{R}=\text{CH}(\text{CH}_3)_2$) was heated to reflux in dry benzene for 12 h under stream of nitrogen. The reaction product was separated by fractional crystallization from hexane-ethanol (1 : 1) and vacuum distillation; it consisted of 1,1',4,4'-tetraphenyl-2,2'-azinodiethanone (**4**, 2%) and 1-isopropyl-3,3,4-triphenyl-2-azetidinone (**3a**, 85%). An authentic sample of product **4** was prepared according to method reported⁶⁾ for comparison (IR, NMR, and also melting point). The structural assignment of **3a** was made on the basis of its analytical and spectral data.

Similar treatment of Schiff's bases **2b—f** gave 2-azetidinones **3b—f**, identified on the basis of their analytical and spectral data, together with ketazine (**4**, 3—5%) in each case.

The racemic *N*-benzylidene- α -methylbenzylamine (**2g**) on similar treatment gave two products, a white crystalline solid **3g** (81%) and a yellow crystalline solid, identified as ketazine **4** (3%). The white crystalline solid, mp 94—95 °C, showed absorption band at 1740 ($\text{C}=\text{O}$, β -lactam²⁾) cm^{-1} in IR spectrum. The NMR spectrum of product exhibited two doublets at δ 1.47 and 1.92, two quartets at δ 4.32 and 5.01 and one singlet at δ 5.02 which may be due to presence of two enantiomeric forms of product **3g**. This would be expected as the starting Schiff's base **2g** is racemic. The elemental analyses and spectral data agree with assigned structure, ($\pm$) 1-(α -methylbenzyl)-3,3,4-tri-

phenyl-2-azetidinone (**3g**), to the product.

The products **3a—g** were found to be unaffected on treatment with ordinary alkali or acid, whereas, *N*-aryl-2-azetidinones reported earlier⁵⁾ have been known to undergo hydrolysis. The formation of products can be explained as in Scheme 1.



Scheme 1.

Thermal decomposition of **1** may lead to benzoylphenylcarbene which may combine with **1** to form ketazine **4**. The benzoylphenylcarbene has been known to undergo Wolff rearrangement to give diphenylketene⁷⁾ which on reaction with $\text{C}=\text{N}$ bond of Schiff's bases **2a—g** may lead to 2-azetidinones **3a—g** through the collapse of a possible zwitterionic intermediate. A similar intermediate has been proposed in the reaction of diphenylketene and *N*-benzylideneaniline giving *N*-phenylimidate.⁸⁾

When *N*-benzhydrylidene-*N'*-phenylurea (**5**) was allowed to react with **1**, in an analogous manner, a white crystalline solid (70%) was obtained. The substituted urea **5** contains three possible reactive sites *viz.* $\text{C}=\text{N}$, $\text{C}=\text{O}$, and NH groups. It is known that interaction of ketene with $\text{C}=\text{N}$, $\text{C}=\text{O}$, and NH groups leads to the formation of β -lactam,⁸⁾ β -lactone⁹⁾ (which on decarboxylation gives corresponding olefin) and acetyl derivative¹⁰⁾ of NH group, respectively. The presence of an absorption bands at 1760 cm^{-1} , characteristic of a β -lactam¹¹⁾ and 1710 ($\text{C}=\text{O}$, $\text{CONH}^{12)$) cm^{-1} , absence of an absorption band between 1700

† 1 mmHg $\approx$ 133.322 Pa.

References

- 1) A. K. Mukerjee and R. C. Srivastava, *Synthesis*, **1973**, 327, and references cited therein.
 - 2) A. K. Bose and I. Kugajevsky, *Tetrahedron*, **23**, 957 (1967).
 - 3) W. Kirmse and L. Horner, *Chem. Ber.*, **89**, 2759 (1956).
 - 4) H. Staudinger, *Chem. Ber.*, **41**, 1355 (1908).
 - 5) H. T. Clarke, J. R. Johnson, and R. Robinson, "The Chemistry of Penicillin," Princeton University Press (1949), p. 945.
 - 6) T. Curtius and R. Kastner, *J. Prakt. Chem.*, **83**, 215 (1911); *Chem. Abstr.*, **5**, 2649 (1911).
 - 7) V. Franzen, *Justus Liebigs Ann. Chem.*, **614**, 31 (1958).
 - 8) H. B. Kagan and J. L. Luche, *Tetrahedron Lett.*, **1968**, 3093.
 - 9) L. L. Muller and J. Hamer, "1,2-Cycloaddition Reactions," Interscience Publ., New York (1967), p. 139.
 - 10) W. Jugelt and D. Schmidt, *Tetrahedron*, **25**, 969 (1969).
 - 11) G. Wittig and A. Hesse, *Justus Liebigs Ann. Chem.*, **1976**, 500.
 - 12) C. D. Nenitzescu and E. Solomonica, *Org. Synth.*, Coll. Vol. II, 496 (1950).
 - 13) A. W. Ingersoll, *Org. Synth.*, Coll. Vol. II, 503 (1950).
 - 14) A. Michaelis, *Chem. Ber.*, **26**, 2169 (1893).
 - 15) G. E. P. Smith, Jr., and F. W. Bergstrom, *J. Am. Chem. Soc.*, **56**, 2095 (1934).
-