

Friedel-Crafts Acylation with 2-Methyl- or 2-Benzylbutanedioic Anhydride

Iwao HASHIMOTO

Department of Industrial Chemistry, Wakayama Technical College, Noshima, Nada-cho, Gobo 649-15

(Received January 8, 1981)

Synopsis. The AlCl_3 -catalyzed acylation of benzene with 2-methylbutanedioic anhydride afforded a mixture of 3-benzoyl-2-methylpropanoic acid and 3-benzoylbutanoic acid. The intramolecular acylation of 2-benzylbutanedioic anhydride in the presence of AlCl_3 gave a mixture of 4-oxo-1,2,3,4-tetrahydro-2-naphthoic acid and 3-oxo-2-indanacetic acid. The results were discussed in terms of the solvent effect on the acylations.

We have previously reported^{1,2)} that the AlCl_3 -catalyzed acylations of benzene with 2-phenyl-butanedioic or -pentanedioic anhydride proceeded by means of competitive inter- and intramolecular acylations, where an electron-attracting phenyl group in the anhydrides had a strong influence on the direction of each acylation. A more detailed mechanism of the acylation with unsymmetrical dibasic acid anhydrides may be get by introducing an electron-donating methyl group into the dibasic acid anhydride and by treating individually the inter- and intramolecular acylations. The present paper deals with the acylation of benzene with 2-methylbutanedioic anhydride (**1**) in the presence of AlCl_3 as an example of the intermolecular acylation, and with the AlCl_3 -catalyzed condensation of 2-benzylbutanedioic anhydride (**2**) as an intramolecular example.


TABLE 1. AlCl_3 -CATALYZED ACYLATION WITH 2-METHYL- OR 2-BENZYLBUTANEDIOIC ANHYDRIDE AT 30 °C FOR 5 h

Anhydride	Solvent (20 cm)	$[\text{AlCl}_3]$ mmol	$[\text{C}_6\text{H}_6]$ mmol	Product yields ^{a)} %	
1				3	4
—	—	20	250 ^{b)}	62	39
—	—	10	250 ^{b)}	28	14
(ClCH_2) ₂	20	10	10	61	34
(ClCH_2) ₂	10	10	10	14	6
$\text{C}_6\text{H}_5\text{NO}_2$	30	10 ^{c)}	10	21	2
$\text{C}_6\text{H}_5\text{NO}_2$	20	10 ^{c)}	10	12	1
2				5	6
—	—	20	250 ^{d)}	37	63
—	—	10	250 ^{d)}	28	37
—	— ^{e)}	250 ^{d)}	65	1	
—	— ^{f)}	250 ^{d)}	34	66	
(ClCH_2) ₂	20	10	10	34	59
(ClCH_2) ₂	10	10	10	26	35
(ClCH_2) ₂	20	0	10	32	60
$\text{C}_6\text{H}_5\text{NO}_2$	30	10	10	81	5

a) Calculated on the basis of the amount of anhydride used. b) At 40 °C for 2 h. c) For 24 h. d) At 40 °C. e) Two hundred mmole of concd H_2SO_4 were used as the catalyst. f) Twenty mmol of AlBr_3 were used as the catalyst.

Several investigators^{3,4)} have isolated only 2-aroil-2-methylpropanoic acid in the condensations of some aromatic compounds with **1** in the presence of AlCl_3 . However, when 10 mmol of **1** was condensed in the

presence of 20 mmol of AlCl_3 with benzene, which was used in a large excess as a reactant and as a solvent, a mixture of 3-benzoyl-2-methylpropanoic acid (**3**) and the isomeric 3-benzoylbutanoic acid (**4**) was obtained; the yield of **3** was about twice that of **4**. The predominant formation of **3** was also observed in the cases using a limited amount (10 mmol) of benzene in 1,2-dichloroethane or in nitrobenzene. The lower yield of the keto acids in nitrobenzene may be a complex formation between AlCl_3 and nitrobenzene. The small solvent effect on the acylation of benzene with **1** suggests that the actual acylating agent in this acylation is a type of oxonium compound (**3a** and **4a**); since they are less polarized than the acylium ions, the solvent effect on them appears smaller.




Scheme 1.

The electron-donating effect of a methyl group reduces the electrophilic reactivity of the $\text{C}=\text{O}$ located closely to the methyl group; hence, the production of **3** is more favored than that of **4**.

The acylation with **2** in the presence of two equivalents of AlCl_3 in a large excess of benzene proceeded only intramolecularly to yield two isomeric keto acids, 4-oxo-1,2,3,4-tetrahydro-2-naphthoic acid (**5**) and 3-oxo-2-indanacetic acid (**6**), in a predominant yield, although early workers⁵⁻⁸⁾ in an analogous case succeeded in isolating only **5**. Similar results were also observed in 1,2-dichloroethane, whereas an overwhelming yield of **5** was obtained in nitrobenzene. Since the polarity of the solvent appears to affect markedly the relative yields between **5** and **6**, a reaction path including either the oxonium compounds (**5a** and **6a**) or the acylium ions (**5b** and **6b**) as the actual acylating agents may be formulated for the intramolecular acylation of **2** in the presence of AlCl_3 .

The formation of **6a** is probably more favored than that of **5a** due to the electron-donating effect of a benzyl group,⁹⁾ although the electrophilic reactivity of $\text{C}=\text{O}$ of **5b** to a benzene nucleus may be greater than that of **6b**; a higher concentration of **6a** results in a predominant yield of **6**. On the other hand, in a polar



solvent such as nitrobenzene or in the presence of a large amount of concd H_2SO_4 , the acylium ions bearing a more positive charge than the oxonium compounds may be greatly stabilized by solvation; therefore, a more reactive **5b** should give **5** in an overwhelming yield.

The product ratio obtained above does not result from the thermodynamic equilibrium, since no isomerization between **5** and **6** was observed in the presence of two equivalents of AlCl_3 in 1,2-dichloroethane.

Experimental

Materials. 2-Methylbutanedioic anhydride: bp 238—240 °C. 2-Benzylbutanedioic anhydride was prepared by the method of Haworth *et al.*: mp 98.5 °C (lit.⁷) mp 95—97 °C).

Acylation Procedures. The general procedure was as previously described.¹⁾ A mixed solution of diethyl ether and methyl acetate was used to extract the keto acids.

Analyses of the Products. The acylation products of benzene with 2-methylbutanedioic anhydride, after being esterified with an ethereal solution of diazomethane, were analyzed by GLC employing a Yanagimoto G-180 F model on a 1.5 m $\times$ 3 mm column packed with Ucon Oil 50 LB 550 X (3 wt %) on Uniport KS of 60—80 mesh at 180 °C. GC analyses of the intramolecular acylation products of 2-benzylbutanedioic anhydride in the presence of AlCl_3 were

made by using a column packed with Apiezon Grease L (3 wt %) at 170 °C or Ucon Oil 50 LB 550 X (3 wt %) at 171 °C. The compounds, **3**, **4** and **5**, were synthesized according to the method described in the literature: 3-benzoyl-2-methylpropanoic acid (**3**): mp 142 °C (lit.¹⁰) mp 140.5 °C; 3-benzoylbutanoic acid (**4**): mp 55 °C (lit.¹¹) mp 59 °C; 4-oxo-1,2,3,4-tetrahydro-2-naphthoic acid (**5**): mp 149 °C (lit.⁵) mp 149 °C).

3-Oxo-2-indanacetic Acid (6). A solution of 2-benzylbutanedioic anhydride (1.90 g) in benzene (12.1 cm³) was treated with AlBr_3 (5.34 g) in benzene (10 cm³) at 30 °C for 5 h. The reaction mixture was then work up in a usual manner. The crude keto acid was repeatedly recrystallized from acetic acid and then methyl acetate–petroleum ether: mp 150.5—151 °C; IR (KBr disk), 1750 ($\text{C}=\text{O}$), 1669, 1600 cm^{-1} ; MS (methyl ester) m/e (%) 204 (M^+ , 30), 173, 172 ($\text{M}^+ - \text{OCH}_3$, 25), 145 ($\text{M}^+ - \text{COOCH}_3$, 75), 131 ($\text{M}^+ - \text{CH}_2\text{COOCH}_3$, 49), 116 (100); $^1\text{H-NMR}$ [$(\text{CD}_3)_2\text{CO}$] δ 7.4—7.7 (m, 4H, arom.), 3.5 (q, 1H, $J=8$), 3.0 (m, 2H), 2.7 (q, 2H, $J=8$); $^{13}\text{C-NMR}$ [$(\text{CD}_3)_2\text{CO}$] δ 209.0 (s, $\text{C}=\text{O}$), 175.5 (s, $-\text{COOH}$), 156.0 (s, C-9), 139.0 (s, C-4), 137.0, 129.5, 129.0 and 125.5 (d, C-arom.), 45.0 (d, C-2), 35.5 (t, CH_2-COOH), 34.0 (t, C-3); Found: C, 69.43; H, 5.31%. Calcd for $\text{C}_{11}\text{H}_{10}\text{O}_3$: C, 69.46; H, 5.30%.

The author wishes to thank Professor Yoshiro Ogata, Nagoya University, for his valuable discussions.

References

- 1) I. Hashimoto and R. Takatsuka, *Bull. Chem. Soc. Jpn.*, **50**, 2495 (1977).
- 2) I. Hashimoto, *Bull. Chem. Soc. Jpn.*, **52**, 251 (1979).
- 3) W. Cocker, A. K. Fateen, and C. Lipman, *J. Chem. Soc.*, **1951**, 926.
- 4) W. Cocker and D. H. Hayes, *J. Chem. Soc.*, **1951**, 844.
- 5) A. J. Attwood, A. Stevenson, and J. F. Thorpe, *J. Chem. Soc.*, **123**, 1764 (1923).
- 6) E. L. Speight, A. Stevenson, and J. F. Thorpe, *J. Chem. Soc.*, **125**, 2185 (1924).
- 7) R. D. Haworth, B. Jones, and Y. M. Way, *J. Chem. Soc.*, **1943**, 10.
- 8) J. V. Braum, *Ber.*, **61**, 2602 (1927).
- 9) J. E. Leffler and E. Grunwald, "Rates and Equilibria of Organic Reactions," Wiley, New York (1963), p. 222.
- 10) E. R. Alexander and A. Mudrak, *J. Chem. Soc.*, **72**, 3195 (1950).
- 11) L. Higginbotham, A. Lapworth, and C. Simpson, *J. Chem. Soc.*, **125**, 2342 (1924).