

Synthesis of Dihydrobenzofuran Derivatives from Substituted *p*-Benzoquinones

Tadashi KOZUKA

Department of Chemistry, Faculty of Science, Kyoto University, Sakyo-ku, Kyoto 606

(Received October 30, 1981)


A series of *p*-benzoquinones having an ester functional group in the α position of alkyl side chains was prepared. Irradiation of these quinones, except for methyl 3-methyl-2-(4-methyl-3,6-dioxo-1,4-cyclohexadienyl)-butyrate (**3e**), rapidly gave methyl 3-substituted 2,3-dihydro-5-hydroxybenzofuran-2-carboxylate in fairly good yields, and it was found that rearrangement of the side chain occurred concomitantly. Photolysis of quinone **3e** afforded methyl 2-(2,5-dihydroxy-4-methylphenyl)-3-methyl-3-butenate as the major product. An intermediate was postulated to account for the isolated photoproduct.

2,3-Dihydrobenzofurans are of interest as a potential synthetic intermediate of a variety of medicinally important natural products¹⁾ in addition to their own pharmacological activity.

In this report, syntheses of a variety of methyl 2,3-dihydro-5-hydroxybenzofuran-2-carboxylate derivatives by way of photochemical reactions are described. One of the possible approaches to the synthesis of this system is photocyclization of substituted *p*-quinones.^{2,3)} The present author has developed a simple general route to the dihydrobenzofuran derivatives starting from alkyl substituted *p*-benzoquinones. The alkyl group has a methoxycarbonyl grouping in its α position.⁴⁾ It is anticipated that the β -hydrogen atom of the alkyl side chain will occupy a favorable position for the γ -hydrogen abstraction by a photoexcited quinonoid carbonyl group, because of the dipole-dipole repulsion between quinonoid carbonyl and methoxycarbonyl groups. The most interesting aspect of the present results is that the photochemical reaction of these quinones occurs with rearrangement to give methyl 3-substituted, 2,3-dihydro-5-hydroxybenzofuran-2-carboxylates.

Results and Discussion

Preparation of Quinones. All quinones **3a—n** were prepared in excellent yields by the synthetic route



(a) $\text{Tl}(\text{NO}_3)_3 \cdot 3\text{H}_2\text{O}$, $\text{CH}(\text{OCH}_3)_3$, MeOH ; (b) $\text{CAN/aq CH}_3\text{CN}$; (c) $h\nu$, benzene; (d) $\text{CAN/aq CH}_3\text{CN}$

$\text{R}_1 = \text{H, Me}$

$\text{R}_2 = \text{H, Me, Et, } n\text{-Pr, CO}_2\text{Me, (CH}_2)_2\text{CO}_2\text{Me, (CH}_2)_3\text{-NPhth,* (CH}_2)_3\text{NHCOCF}_3, \text{CH}_2\text{OCO}_2\text{C}_6\text{H}_5$

$\text{R}_3 = \text{H, Me}$

$\text{X} = \text{H, Me, Br, OMe}$

* NPhth = phthalimido

Scheme 1.

shown in Scheme 1. That is, oxythallation of the ketone **1** with thallium(III) nitrate (TTN)⁵⁾ and trimethyl orthoformate (TMOF) in methanol followed by oxidation of the ester **2** with ceric ammonium nitrate (CAN)^{6a)} in 70% aqueous acetonitrile to give the quinone **3**.

Irradiation of Quinones. On irradiation of a solution of the quinone **3** (2 mmol) in dry benzene with a high pressure Hg lamp (300 W) after bubbling nitrogen, the quinone solution rapidly turned from yellow to colorless. The photochemical reaction was complete within several hours (0.5—5 h) at 0—10 °C and gave methyl 3-substituted 2,3-dihydro-5-hydroxybenzofuran-2-carboxylate **4** in relatively good yields after purification by column chromatography. The results of photolyses of the quinones **3a—n** are summarized in Table 1.

The structures of the irradiation products **4a—n** were established by their IR, ¹H NMR data, and chemical reactions. In particular, the structure of **4a** was confirmed as follows. The ¹H NMR spectrum of **4a** consisted of a characteristic AB pattern due to geminal protons at C-3 ($J_{AB} = 15 \text{ Hz}$),⁷⁾ indicating the presence of a C-2 methoxycarbonyl-substituted 2,3-dihydrobenzofuran ring (C-2 methoxycarbonyl, IR: 1740 cm^{-1} , CO). Such an ABX pattern of C-2 substituted 2,3-dihydrobenzofuran was similar to that of ethyl 2,3-

TABLE 1. FORMATION OF PHOTOREARRANGED DIHYDROBENZOFURAN **4** FROM QUINONE **3**

Quinone	X	R ₁	R ₂	R ₃	Product	Yield/% ^{a)}
3a	Me	H	H	H	4a	41
3b	Me	H	Me	H	4b	65
3c	Me	Me	H	H	4c	45
3d	Me	H	Et	H	4d	68
3e	Me	H	Me	Me	4e ^{b)}	40
3f	Me	H	<i>n</i> -Pr	H	4f	60
3g	Me	H	CO ₂ Me	H	4g	56
3h	Me	H	(CH ₂) ₂ CO ₂ Me	H	4h	51
3i	Me	H	(CH ₂) ₃ NPhth	H	4i	75
3j	Me	H	(CH ₂) ₃ NHCOCF ₃	H	4j	69
3k	Me	H	CH ₂ OCO ₂ C ₆ H ₅	H	4k	39
3l	H	H	Me	H	4l	60
3m	Br	H	Me	H	4m	31
3n	OMe	H	Me	H	4n	42

a) Reported yield are for isolated, purified products.

b) No dihydrobenzofuran deriv was isolated. Product was methyl 2-(2,5-dihydroxy-4-methylphenyl)-3-methyl-3-butenate.

dihydro-5-hydroxybenzofuran-2-carboxylate.⁸⁾

For the photoproduct **4c**, the ¹H NMR spectrum showed a coupling ($J_{AB}=15$ Hz),⁷⁾ which suggested that the substitution pattern was similar to that of **4a**. Furthermore, the structures of **4a** and **4c** were also supported by oxidation with ceric ammonium nitrate^{6b)} to the corresponding quinones **5a** and **5c**. ¹H NMR spectra of **5a** and **5c** showed the presence of methylene protons directly attaching to the quinone ring. The methylene protons were long-range coupled with the quinone ring proton ($J=1.2$ Hz).⁹⁾

The photoproduct **4g** was compatible with the structure of dimethyl 2,3-dihydro-5-hydroxy-6-methylbenzofuran-2,3-dicarboxylate on the basis of the spectral data outlined below. Its IR spectrum showed three characteristic bands at 3340 (phenolic OH), 1735 (ester CO), and 1695 cm⁻¹ (ester CO). Its ¹H NMR spectrum showed two doublets at δ 4.41 ($J=6$ Hz) and δ 5.56 ($J=6$ Hz) due to the protons C₃-H and C₂-H, respectively, and a broad singlet at δ 5.32 due to the phenolic hydroxyl proton.


Irradiation of the quinones **3b**, **3d**, **3f**, **3l**, **3m**, and **3n** afforded predominantly the corresponding dihydrobenzofurans **4b**, **4d**, **4f**, **4l**, **4m**, and **4n**, respectively, and a small amount of isomers. The major photoproducts in the reactions were isolated by chromatography and/or fractional crystallization. The minor products were detected by ¹H NMR and GC, but not isolated. The ¹H NMR spectra of the above six major products showed similar chemical shifts and coupling constants for the proton signals at C-3 and C-2 on the dihydrobenzofuran ring as shown in Table 2. The regiochemistry for the substituents around C-2 and C-3 of these photoproducts cannot be determined directly by comparison with the chemical shifts of C-2 and C-3 protons of other products, i.e., **4a**, **4c**, and **4g** as was already described. In general, chemical shifts of C-2 and C-3 protons on 2,3-disubstituted 2,3-dihydrobenzofurans are variable according to the substituent groups and stereospecific environment.¹⁰⁾ Therefore, **4b**, **4d**, and **4f** were converted by oxidation with CAN to

quinones **5b**, **5d**, and **5f**, respectively. The structures of these three quinones were assigned on the basis of the spectroscopic data. For example, quinone **5f** showed three characteristic bands at 3470 (OH), 1720 (ester CO), and 1650 cm⁻¹ (quinone CO) in its IR spectrum. Its ¹H NMR spectrum consisted of the signals at δ 1.64 (m, 4H), at δ 2.87 (d, $J=7$ Hz, 1H), at δ 3.31 (m, 1H), and at δ 4.31 (dd, $J=3.5$ Hz, and 7 Hz, 1H). On adding D₂O the signal at δ 2.87 disappeared and the double doublets signal at δ 4.31 was simplified to a doublet ($J=3.5$ Hz). Irradiation of the methylene proton at δ 1.64 changed the signal at δ 3.31 to a sharp doublet ($J=3.5$ Hz), and the signal at δ 4.31 coupled with the methine proton (δ 3.31) to a triplet ($J=7$ Hz) and the hydroxyl proton signal at δ 2.87 to a singlet. These results suggested that the quinone **5f** included the partial structure $-\text{CH}_2\text{CHCH}(\text{OH})\text{CO}-$ in the side chains. Further support for the structure **5f** was provided by the reduction of the quinone **5f** with sodium dithionite followed by acetylation of the obtained hydroquinone with acetic anhydride-pyridine to the corresponding triacetate **7**. However, the methine proton signal adjacent to the acetoxyl group showed a downfield shift at δ 5.18 (d, $J=5$ Hz) due to a deshielding effect of the acetoxyl carbonyl. The above results indicated that these original photoproducts **4b**, **4d**, **4f**, **4l**, **4m**, and **4n** contain an ester group at C-2 position and other substituent group (R₂ and R₃) at C-3 position, respectively.

Irradiation of other substituted quinones **3h**, **3i**, **3j**, and **3k** also gave corresponding photocyclization products **4h**, **4i**, **4j**, and **4k**.

The ¹H NMR spectrum of photoproduct **4k** showed the presence of a methine proton signal at δ 5.02 (d, $J=5.1$ Hz), probably at C-2 position. This lower field shift can be explained in terms of the deshielding effect of the carbonate carbonyl group.

The structures of the major and the minor components were correlated with each other by their chemical reactions and by comparing their ¹H NMR spectra. Acetylation of crude photoproducts **4b**, **4d**, **4f**, and



Scheme 2.

TABLE 2. ¹H NMR SPECTRA OF DIHYDROBENZOFURANS **4** AND QUINONES **5**

X	R ₁	R ₂	Dihydrobenzofuran 4						Quinone 5		
			<i>trans</i> -Isomer			<i>cis</i> -Isomer			C _α -H	C _β -H	OH
			C ₂ -H	C ₃ -H	R ₂ =Me	C ₂ -H	R ₂ =Me				
a	Me	H	5.12(dd) <i>J</i> =7 Hz <i>J</i> =10 Hz	3.14 } (m) 3.62 A	—	—	—	4.13(dd) <i>J</i> =5 Hz <i>J</i> =8.5 Hz	D	B	
b	Me	Me	4.63(d) <i>J</i> =7 Hz	3.40 } (m) 3.68	1.40(d) <i>J</i> =7 Hz	5.15(d) <i>J</i> =9 Hz	1.16(d)	4.30(bm)	3.46(dq) <i>J</i> =4 Hz <i>J</i> =7 Hz	3.00(bs)	
c	Me	Me	—	3.00(d) <i>J</i> =15 Hz 3.50 <i>J</i> =15 Hz	—	—	—	—	2.70(d) <i>J</i> =13.5 Hz 2.97(d) <i>J</i> =13.5 Hz	3.34(s)	
d	Me	Et	4.74(d) <i>J</i> =5 Hz	3.42(m)	—	5.14(d) <i>J</i> =8.5 Hz	—	4.36 <i>J</i> =3.5 Hz <i>J</i> =7.6 Hz	3.23(dt) <i>J</i> =3.5 Hz <i>J</i> =7 Hz	2.98(d) <i>J</i> =7.6 Hz	
f	Me	<i>n</i> -Pr	4.76(d) <i>J</i> =5 Hz	3.46(m)	—	5.12(d) <i>J</i> =8.5 Hz	—	4.31(dd) <i>J</i> =3.5 Hz <i>J</i> =7 Hz	3.31(m) <i>J</i> =3.5 Hz <i>J</i> =7 Hz	2.87(d) <i>J</i> =7 Hz	
g	Me	CO ₂ Me	5.56(d) <i>J</i> =6 Hz	4.41(d) <i>J</i> =6 Hz	—	C	—	4.86(d) <i>J</i> =4.2 Hz	4.32(d) <i>J</i> =4.2 Hz	3.75(bs)	
h	Me	(CH ₂) ₂ CO ₂ Me	4.77(d) <i>J</i> =6 Hz	3.44—	—	C	—	4.35(bs)	3.38(m)	3.07(bd) <i>J</i> =8 Hz	
i	Me	(CH ₂) ₃ NPh _t	4.76(d) <i>J</i> =5 Hz	B	—	C	—	—	—	—	
j	Me	(CH ₂) ₃ NHCOCF ₃	4.71(d) <i>J</i> =5 Hz	B	—	5.15(d) <i>J</i> =9 Hz	—	—	—	—	
k	Me	H	CH ₂ OCO ₂ C ₆ H ₅	5.02(d) <i>J</i> =5.1 Hz	3.90(m)	—	C	—	—	—	
l	H	H	Me	4.64(d) <i>J</i> =7 Hz	3.53(m)	1.38(d) <i>J</i> =7 Hz	5.16(d) <i>J</i> =9 Hz	1.13(d) <i>J</i> =7 Hz	—	—	
m	Br	H	Me	4.65(d) <i>J</i> =7 Hz	3.53(m)	1.40(d) <i>J</i> =7 Hz	5.15(d) <i>J</i> =9 Hz	1.16(d) <i>J</i> =7 Hz	—	—	
n	OMe	H	Me	4.66(d) <i>J</i> =7 Hz	3.56(m)	1.41(d) <i>J</i> =7 Hz	5.17(d) <i>J</i> =8.5 Hz	1.15(d) <i>J</i> =7 Hz	—	—	

a) The *cis*-isomer, which was a minor component of the mixture of products, was detectable only by the ¹H NMR spectrum. (A) AB type: *J*=15 Hz, (B) Chemical shifts were not determined because of overlapping of the signal with those of other protons. (C) Not detected. (D) δ 2.69(dd), *J*=8.5 and 15 Hz, 1H, δ 2.98 (dd, *J*=5 and 15 Hz, 1H).


4i containing minor components with acetic anhydride in pyridine solution gave only the single monoacetate **6b**, **6d**, **6f**, and **6i**, respectively. The minor components are therefore presumed to be the stereoisomers of the thermodynamically stable major products. In the ¹H NMR spectra, the coupling constants due to C-2 and C-3 protons of 2,3-disubstituted 2,3-dihydrobenzofuran are *J*_{cis} > *J*_{trans}¹¹⁾ as predicted by the Karplus equation.¹²⁾ The most striking differences between the ¹H NMR spectra of the *cis* and *trans* isomers of 2,3-disubstituted 2,3-dihydrobenzofuran, other than the values of *J*_{2,3} are the different values for the chemical shift of the C-2 proton. The ¹H NMR signal of the C-2 proton of the *trans* isomers is located upfield at about δ 0.3 compared

to the corresponding proton of the *cis* isomers. The major product **4** (δ 4.63–4.77, *J*=5–7 Hz) is concluded to be *trans*, and the minor to be *cis* (δ 5.14–5.17, *J*=8.5–9 Hz) as shown in Table 2.

On the other hand, the photolysis of quinone **3e** did not afford a dihydrobenzofuran derivative, but methyl 2-(2,5-dihydroxy-4-methylphenyl)-3-methyl-3-butenolate **4e** as the major product.

The proposed mechanism of these photochemical reactions is given in Scheme 2. γ-Hydrogen abstraction by photoexcited carbonyl of the quinone **3** forms the biradical intermediate **A**¹³⁾ and subsequently yields the spirocyclopropyl ketone **B-1**³⁾ and/or spirocyclopropyl zwitter-ion intermediate **B-2**³⁾ then the cyclopropane

ring opens to afford the observed product **4**. However it is difficult to anticipate a ring-opened zwitter-ion intermediate,¹⁴⁾ because the three-membered ring may cleave to give a positive charge at the α position to the methoxycarbonyl group. From the present results preferential cleavage of bond **a** can be recognized **B-1** to give phenyl migrated products **4** with exception of only one case **3e** ($R_2=R_3=Me$). In the latter case cleavage of bond **b** may be the preferential course to give **4e**, in which no phenyl migration occurs. Thus, among the alkyl substituted *p*-benzoquinones examined, the trend of phenyl migration initiated by photochemical γ -hydrogen abstraction reaction may be explained in terms of the stability of the radical intermediates involved. The present results indicate the stability of the radical to be



Of course, this trend should be applied under the very limited conditions, for a contribution of a dipolar intermediate, such as **B-2**, could not be excluded in the present reactions.

Experimental

Melting points and boiling points were uncorrected. Bulb to bulb distillations were carried out on a Kugelrohr-type apparatus. IR spectra were measured on a JASCO IRA-1 spectrophotometer. ¹H NMR spectra were measured on a JEOL PS-100 instrument with tetramethylsilane as an internal standard and the chemical shifts are reported in δ values. Mass spectra were obtained with a Hitachi M-52 mass spectrometers (20 eV). Elemental analyses were performed by the Kyoto University Microanalytical Laboratories. Column chromatography was performed using Wako reagent grade silica gel (200 mesh). The irradiation was carried out in a Pyrex vessel at 0–10 °C, using an Eikosha 300-W high pressure Hg lamp.

Materials. 3-(Methoxycarbonyl)propionyl chloride¹⁵⁾ and 5-(methoxycarbonyl)pentanoyl chloride¹⁶⁾ were prepared from the corresponding methyl hydrogen succinate and methyl hydrogen adipate, respectively. 6-Phthalimidohehexanoyl chloride¹⁷⁾ and 6-(trifluoroacetamido)hexanoyl chloride¹⁸⁾ were prepared from 6-aminohexanoic acid. Other acid chlorides were commercially available.

Preparation of Starting Materials. All of the compounds **1a–n**, except for **1k**, were prepared by the Friedel-Crafts acylation according to the previous reported methods.^{5a)}

Physical Properties of the Starting Materials. 2',5'-Dimethoxy-4'-methylpropiophenone **1a**: 96%; mp 78 °C,^{5a)} (lit,¹⁹⁾ mp 76–77 °C).

2',5'-Dimethoxy-4'-methylbutyrophenone **1b**: 96%; bp 129–130 °C at 3 Torr (1 Torr=133.322 Pa); mp 46–46.5 °C (hexane-ether); IR (KBr): 1660 cm⁻¹; ¹H NMR (CDCl₃) δ : 0.93 (t, 3H), 1.67 (m, 2H), 2.23 (s, 3H), 2.92 (t, 2H), 3.80 (s, 3H), 3.84 (s, 3H), 6.74 (s, 1H), and 7.20 (s, 1H); Found: C, 69.99; H, 8.21%. Calcd for C₁₃H₁₈O₃: C, 70.24; H, 8.16%.

2',5'-Dimethoxy-2,4'-dimethylpropiophenone **1c**: 99%; bp 125–127 °C at 3 Torr; mp 38–39 °C (petroleum ether); IR (neat): 1675 cm⁻¹; ¹H NMR (CDCl₃) δ : 1.14 (d, 6H), 2.25 (s, 3H), 3.57 (m, 1H), 3.82 (s, 3H), 3.86 (s, 3H), 6.78 (s, 1H), and 7.14 (s, 1H); Found: C, 69.96; H, 8.20%. Calcd for

C₁₃H₁₈O₃: C, 70.24; H, 8.16%.

2',5'-Dimethoxy-4'-methylvalerophenone **1d**: 95%; bp 140–143 °C at 3 Torr; mp 33.5–34 °C (petroleum ether); IR (KBr): 1670 cm⁻¹; ¹H NMR (CDCl₃) δ : 1.92 (t, 3H), 1.20–1.90 (m, 4H), 2.25 (s, 3H), 2.97 (t, 2H), 3.80 (s, 3H), 3.85 (s, 3H), 6.77 (s, 1H), and 7.21 (s, 1H); Found: C, 71.21; H, 8.75%. Calcd for C₁₄H₂₀O₃: C, 71.16; H, 8.53%.

2',5'-Dimethoxy-3,4'-dimethylbutyrophenone **1e**: 88%; bp 140–143 °C at 3 Torr; mp 48–48.5 °C (petroleum ether); IR (KBr): 1600 cm⁻¹; ¹H NMR (CDCl₃) δ : 0.97 (d, 6H), 2.08–2.41 (m, 1H), 2.26 (s, 3H), 2.84 (d, 2H), 3.81 (s, 3H), 3.85 (s, 3H), 6.76 (s, 1H), and 7.19 (s, 1H); Found: C, 71.42; H, 8.69%. Calcd for C₁₄H₂₀O₃: C, 71.16; H, 8.53%.

2',5'-Dimethoxy-4'-methylhexanophenone **1f**: 85%; bp 183 °C at 3 Torr; IR (neat): 1665 cm⁻¹; ¹H NMR (CDCl₃) δ : 0.90 (t, 3H), 1.22–1.82 (m, 6H), 2.26 (s, 3H), 2.98 (t, 2H), 3.84 (s, 3H), 3.88 (s, 3H), 6.81 (s, 1H), and 7.27 (s, 1H); Found: C, 72.25; H, 8.87%. Calcd for C₁₅H₂₂O₃: C, 71.97; H, 8.86%.

Methyl 3-(2,5-Dimethoxy-4-methylbenzoyl)propionate **1g**: 90%; mp 89–90 °C (petroleum ether-dichloromethane); IR (KBr): 1735 and 1660 cm⁻¹; ¹H NMR (CDCl₃) δ : 2.26 (s, 3H), 2.69 (t, 2H), 3.32 (t, 2H), 3.70 (s, 3H), 3.81 (s, 3H), 3.88 (s, 3H), 6.77 (s, 1H), and 7.30 (s, 1H); Found: C, 63.33; H, 6.86%. Calcd for C₁₄H₁₈O₅: C, 63.14; H, 6.81%.

Methyl 5-(2,5-Dimethoxy-4-methylbenzoyl)pentanoate **1h**: 73%; mp 80–81 °C (petroleum ether-dichloromethane); IR (KBr): 1730 and 1660 cm⁻¹; ¹H NMR (CDCl₃) δ : 1.64–1.86 (m, 4H), 2.26 (s, 3H), 2.36 (m, 2H), 3.00 (m, 2H), 3.67 (s, 3H), 3.82 (s, 3H), 3.87 (s, 3H), 6.78 (s, 1H), and 7.23 (s, 1H); Found: C, 65.33; H, 7.58%. Calcd for C₁₆H₂₂O₅: C, 65.29; H, 7.53%.

2',5'-Dimethoxy-4'-methyl-6-(phthalimido)hexanophenone **1i**: 71%; mp 156–157 °C (methanol-dichloromethane); IR (KBr): 1775, 1715, and 1660 cm⁻¹; ¹H NMR (CDCl₃) δ : 1.31–1.87 (m, 6H), 2.25 (s, 3H), 2.97 (t, 2H), 3.69 (t, 2H), 3.81 (s, 3H), 3.86 (s, 3H), 6.75 (s, 1H), 7.21 (s, 1H), and 7.63–7.77 (m, 4H); Found: C, 69.92; H, 6.39, N, 3.54%. Calcd for C₂₃H₂₅O₅N: C, 69.85; H, 6.37; N, 3.54%.

2',5'-Dimethoxy-4'-methyl-6-(trifluoroacetamido)hexanophenone **1j**: 56%; mp 87.5–89 °C (petroleum ether-dichloromethane); IR (KBr): 3300, 1730, and 1640 cm⁻¹; ¹H NMR (CDCl₃) δ : 1.20–1.84 (m, 6H), 2.25 (s, 3H), 3.00 (t, 2H), 3.39 (q, 2H), 3.82 (s, 3H), 3.86 (s, 3H), 6.80 (s, 1H), 6.90 (bs, 1H), and 7.24 (s, 1H); Found: C, 56.43; H, 6.12; N, 4.00%. Calcd for C₁₇H₂₂O₄NF₃: C, 56.50; H, 6.14; N, 3.88%.

2',5'-Dimethoxy-4'-methyl-4-(phenoxycarbonyloxy)butyrophenone **1k**: was prepared from **1g** in the following three steps. Reduction of the **1g** with excess lithium aluminum hydride in THF followed by reaction of the obtained 1-(2,5-dimethoxy-4-methylphenyl)-1,4-butanediol (1 mol equiv.) in dichloromethane with phenyl chloroformate (1.2 mol equiv.) in the presence of pyridine at 0 °C afforded 1-(2,5-dimethoxy-4-methylphenyl)-4-(phenoxycarbonyloxy)-1-butanol, which on oxidation with excess Cr-Pyr complex in dichloromethane yield **1k**, total yield: 30%; IR (neat): 1760 and 1660 cm⁻¹; ¹H NMR (CDCl₃) δ : 2.00–2.36 (m, 2H), 2.21 (s, 3H), 3.12 (t, $J=7$ Hz, 2H), 3.81 (s, 3H), 3.87 (s, 3H), 4.31 (d, $J=6.5$ Hz, 2H), 6.78 (s, 1H), and 7.12–7.46 (m, 6H); Found: C, 66.81; H, 6.30%. Calcd for C₂₀H₂₂O₆: C, 67.02; H, 6.19%.

2',5'-Dimethoxybutyrophenone **1l**: 90%; bp 121–123 °C at 3 Torr, (lit,²⁰⁾ bp 172–175 °C at 25 Torr).

2',4',5'-Trimethoxybutyrophenone **1n**: 88%; mp 79 °C (petroleum ether-dichloromethane), (lit,^{21,22)} mp 77 °C, 78.5–80 °C).

The following esters **2a–n**, except for **2m**, were prepared according to the reported methods.⁵⁾

Methyl 2-(2,5-Dimethoxy-4-methylphenyl)propionate **2a**: was

prepared from **1a** with $\text{Ti}(\text{NO}_3)_3$ -TMOF-MeOH as described previously^{5a)} (90%): bp 162–165 °C at 4 Torr (bulb to bulb).

Methyl 2-(2,5-Dimethoxy-4-methylphenyl)butyrate 2b: was prepared from **1b** with $\text{Ti}(\text{NO}_3)_3$ -TMOF-MeOH following by the general procedure (96%): bp 150–152 °C at 3 Torr (bulb to bulb); IR (neat): 1735 cm^{-1} ; ^1H NMR (CDCl_3) δ : 0.87 (t, 3H), 1.56–2.20 (m, 2H), 2.20 (s, 3H), 3.64 (s, 3H), 3.77 (s, 6H), 3.90 (t, 1H), 6.68 (s, 1H), and 6.74 (s, 1H); Found: C, 66.49; H, 8.17%. Calcd for $\text{C}_{14}\text{H}_{20}\text{O}_4$: C, 66.64; H, 7.99%.

Methyl 2-Methyl-2-(2,5-dimethoxy-4-methylphenyl)propionate 2c: was prepared from **1c** with $\text{Ti}(\text{NO}_3)_3$ -TMOF-MeOH following by the same general procedure (95%): bp 154–156 °C at 4 Torr (bulb to bulb); IR (neat): 1740 cm^{-1} ; ^1H NMR (CDCl_3) δ : 1.49 (s, 6H), 2.18 (s, 3H), 3.60 (s, 3H), 3.68 (s, 3H), 3.78 (s, 3H), 6.63 (s, 1H), and 6.74 (s, 1H); Found: C, 66.50; H, 8.27%. Calcd for $\text{C}_{14}\text{H}_{20}\text{O}_4$: C, 66.64; H, 7.99%.

Methyl 2-(2,5-Dimethoxy-4-methoxy-4-methylphenyl)pentanoate 2d: was prepared from **1d** with $\text{Ti}(\text{NO}_3)_3$ -TMOF-MeOH following by the general reaction procedure (98%): bp 158–159 °C at 3 Torr (bulb to bulb); IR (neat): 1730 cm^{-1} ; ^1H NMR (CDCl_3) δ : 0.90 (t, 3H), 1.09–2.10 (m, 4H), 2.19 (s, 3H), 3.63 (s, 3H), 3.77 (s, 3H), 4.00 (t, 1H), 6.67 (s, 1H), and 6.76 (s, 1H); Found: C, 67.79; H, 8.49%. Calcd for $\text{C}_{15}\text{H}_{22}\text{O}_4$: C, 67.64; H, 8.33%.

Methyl 2-(2,5-Dimethoxy-4-methylphenyl)-3-methylbutyrate 2e: was prepared from **1e** with $\text{Ti}(\text{NO}_3)_3$ -TMOF-MeOH following by the usual manner (81%): bp 133–135 °C at 3 Torr (bulb to bulb); mp 78–79 °C (petroleum ether); IR (KBr): 1735 cm^{-1} ; ^1H NMR (CDCl_3) δ : 0.71 (d, 3H), 1.03 (d, 3H), 2.00–2.26 (m, 1H), 2.20 (s, 3H), 3.65 (s, 3H), 3.79 (s, 6H), 3.84 (d, 1H), 6.71 (s, 1H), and 6.91 (s, 1H); Found: C, 67.69; H, 8.41%. Calcd for $\text{C}_{15}\text{H}_{22}\text{O}_4$: C, 67.64; H, 8.33%.

Methyl 2-(2,5-Dimethoxy-4-methylphenyl)hexanoate 2f: was prepared from **1f** with $\text{Ti}(\text{NO}_3)_3$ -TMOF-MeOH following by the general method (90%): bp 135–140 °C at 0.1 Torr (bulb to bulb); IR (neat): 1735 cm^{-1} ; ^1H NMR (CDCl_3) δ : 0.86 (t, 3H), 1.00–2.19 (m, 6H), 2.19 (s, 3H), 3.64 (s, 3H), 3.78 (s, 6H), 3.98 (t, 1H), 6.68 (s, 1H), and 6.75 (s, 1H); Found: C, 68.30; H, 8.77%. Calcd for $\text{C}_{16}\text{H}_{24}\text{O}_4$: C, 68.54; H, 8.63%.

Dimethyl 2-(2,5-Dimethoxy-4-methylphenyl)succinate 2g: was prepared from **1g** with $\text{Ti}(\text{NO}_3)_3$ -TMOF-MeOH following by the general procedure (95%): mp 93–94 °C (petroleum ether–dichloromethane); IR (KBr): 1735 cm^{-1} ; ^1H NMR (CDCl_3) δ : 2.18 (s, 3H), 2.54 (dd, 1H), 3.14 (dd, 1H), 3.65 (s, 6H), 3.75 (s, 6H), 4.35 (dd, 1H), 6.62 (s, 1H), and 6.67 (s, 1H); Found: C, 60.57; H, 6.86%. Calcd for $\text{C}_{15}\text{H}_{20}\text{O}_6$: C, 60.80; H, 6.80%.

Dimethyl 2-(2,5-Dimethoxy-4-methylphenyl)hexanedioate 2h: was prepared from **1h** with $\text{Ti}(\text{NO}_3)_3$ -TMOF-MeOH by the same method (99%): bp 129–131 °C at 0.06 Torr (bulb to bulb); IR (neat): 1735 cm^{-1} ; ^1H NMR (CDCl_3) δ : 1.40–2.20 (m, 4H), 2.22 (s, 3H), 2.34 (t, 2H), 3.66 (s, 6H), 3.80 (s, 6H), 4.02 (t, 1H), 6.72 (s, 1H), and 6.75 (s, 1H); Found: C, 62.68; H, 7.40%. Calcd for $\text{C}_{17}\text{H}_{24}\text{O}_6$: C, 62.95; H, 7.46%.

Methyl 2-(2,5-Dimethoxy-4-methylphenyl)-6-(phthalimido)hexanoate 2i: was prepared from **1i** with $\text{Ti}(\text{NO}_3)_3$ -TMOF-MeOH by the usual method (84%): mp 96–97.5 °C (methanol); IR (KBr): 1775, 1720, and 1710 cm^{-1} ; ^1H NMR (CDCl_3) δ : 1.16–1.84 (m, 6H), 2.14 (s, 3H), 3.60 (s, 3H), 3.60 (m, 2H), 3.73 (s, 6H), 3.95 (t, 1H), 6.62 (s, 1H), 6.68 (s, 1H), and 7.58–7.84 (m, 4H); Found: C, 67.70; H, 6.39; N, 3.28%. Calcd for $\text{C}_{24}\text{H}_{27}\text{O}_6\text{N}$: C, 67.75; H, 6.40; N, 3.29%.

Methyl 2-(2,5-Dimethoxy-4-methylphenyl)-6-(trifluoroacetamido)hexanoate 2j: was prepared from **1j** with $\text{Ti}(\text{NO}_3)_3$ -TMOF-MeOH following by the general procedure (93%): mp 86.5–88 °C (petroleum ether–ether); IR (KBr): 3340, 1725, and 1710 cm^{-1} ; ^1H NMR (CDCl_3) δ : 1.20–2.20 (m, 6H), 2.20 (s,

3H), 3.34 (q, 2H), 3.66 (s, 3H), 3.79 (s, 6H), 6.56 (bs, 1H), and 6.71 (s, 2H); Found: C, 55.27; H, 6.30; N, 3.52%. Calcd for $\text{C}_{18}\text{H}_{24}\text{O}_6\text{NF}_3$: C, 55.24; H, 6.18; N, 3.58%.

Methyl 2-(2,5-Dimethoxy-4-methylphenyl)-4-(phenoxycarbonyloxy)butyrate 2k: was prepared from **1k** with $\text{Ti}(\text{NO}_3)_3$ -TMOF-MeOH following by the same procedure (70%): IR (neat): 1760 and 1735 cm^{-1} ; ^1H NMR (CDCl_3) δ : 1.92–2.71 (m, 2H), 2.21 (s, 3H), 3.67 (s, 3H), 3.77 (s, 6H), 4.07–4.33 (m, 3H), 6.73 (s, 2H), and 7.13–7.42 (m, 5H); Found: C, 65.13; H, 6.25%. Calcd for $\text{C}_{21}\text{H}_{24}\text{O}_7$: C, 64.93; H, 6.23%.

Methyl 2-(2,5-Dimethoxyphenyl)butyrate 2l: was prepared from **1l** with $\text{Ti}(\text{NO}_3)_3$ -TMOF-MeOH following by the general procedure (91%): bp 120–121 °C at 3 Torr (bulb to bulb); IR (neat): 1730 cm^{-1} ; ^1H NMR (CDCl_3) δ : 0.88 (t, 3H), 1.57–2.19 (m, 2H), 3.64 (s, 3H), 3.75 (s, 3H), 3.77 (s, 3H), 3.89 (t, 1H), and 6.77–6.83 (m, 3H); Found: C, 65.81; H, 7.80%. Calcd for $\text{C}_{13}\text{H}_{18}\text{O}_4$: C, 65.53; H, 7.61%.

Methyl 2-(4-Bromo-2,5-dimethoxyphenyl)butyrate 2m: was prepared from **2l** with bromine in chloroform (96%): bp 136–140 °C at 0.05 Torr (bulb to bulb); IR (neat): 1730 cm^{-1} ; ^1H NMR (CDCl_3) δ : 0.88 (t, 3H), 1.56–2.18 (m, 2H), 3.66 (s, 3H), 3.84 (s, 3H), 3.89 (s, 3H), 3.89 (m, 1H), 6.86 (s, 1H), and 7.04 (s, 1H); Found: C, 49.24; H, 5.52; Br, 25.18%. Calcd for $\text{C}_{13}\text{H}_{17}\text{O}_4\text{Br}$: C, 49.22; H, 5.40; Br, 25.20%.

Methyl 2-(2,4,5-Trimethoxyphenyl)butyrate 2n: was prepared from **1n** with $\text{Ti}(\text{NO}_3)_3$ -TMOF-MeOH following by the general method (60%): mp 60–62 °C (petroleum ether); IR (KBr): 1735 cm^{-1} ; ^1H NMR (CDCl_3) δ : 0.87 (t, 3H), 1.60–2.12 (m, 2H), 3.64 (s, 3H), 3.80 (s, 3H), 3.82 (s, 3H), 3.88 (s, 3H and m, 1H), 6.49 (s, 1H), and 6.80 (s, 1H); Found: C, 62.91; H, 7.60%. Calcd for $\text{C}_{14}\text{H}_{20}\text{O}_5$: C, 62.67; H, 7.51%.

Preparation of Quinones 3. Quinones **3a–n** were prepared from the corresponding **2a–n** with ceric ammonium nitrate (CAN) in aq acetonitrile by the method of Jacob *et al.*^{5a)}

Methyl 2-(4-Methyl-3,6-dioxo-1,4-cyclohexadienyl)propionate 3a: (CAN oxidation of **2a**): 88%; mp 59–60 °C.^{5a)}

Methyl 2-(4-Methyl-3,6-dioxo-1,4-cyclohexadienyl)butyrate 3b: (CAN oxidation of **2b**): 98%; bp 130–131 °C at 3 Torr (bulb to bulb); IR (neat): 1735 and 1655 cm^{-1} ; ^1H NMR (CDCl_3) δ : 0.92 (t, $J=7$ Hz, 3H), 1.60–2.00 (m, 2H), 2.04 (d, $J=1.6$ Hz, 3H), 3.70 (s, 3H), 3.70–3.98 (m, 1H), 6.64 (q, $J=1.6$ Hz, 1H), and 6.70 (s, 1H); Found: C, 64.98; H, 6.60%. Calcd for $\text{C}_{12}\text{H}_{14}\text{O}_4$: C, 64.85; H, 6.35%.

Methyl 2-Methyl-2-(4-methyl-3,6-dioxo-1,4-cyclohexadienyl)propionate 3c: (CAN oxidation of **2c**): 95%; mp 111–112 °C (petroleum ether); IR (KBr): 1740 and 1650 cm^{-1} ; ^1H NMR (CDCl_3) δ : 1.42 (s, 6H), 2.03 (d, $J=1.6$ Hz, 3H), 3.62 (s, 3H), 6.56 (q, $J=1.6$ Hz, 1H), and 6.59 (s, 1H); Found: C, 64.56; H, 6.35%. Calcd for $\text{C}_{12}\text{H}_{14}\text{O}_4$: C, 64.85; H, 6.35%.

Methyl 2-(4-Methyl-3,6-dioxo-1,4-cyclohexadienyl)pentanoate 3d: (CAN oxidation of **2d**): 96%; bp 143–145 °C at 3 Torr (bulb to bulb); IR (neat): 1735 and 1655 cm^{-1} ; ^1H NMR (CDCl_3) δ : 0.92 (t, $J=7$ Hz, 3H), 1.15–2.04 (m, 4H), 2.04 (d, $J=1.6$ Hz, 3H), 3.68 (s, 3H), 3.90 (t, $J=7$ Hz, 1H), 6.62 (q, $J=1.6$ Hz, 1H), and 6.69 (s, 1H); Found: C, 66.05; H, 6.89%. Calcd for $\text{C}_{13}\text{H}_{16}\text{O}_4$: C, 66.08; H, 6.83%.

Methyl 3-Methyl-2-(4-methyl-3,6-dioxo-1,4-cyclohexadienyl)-butyrate 3e: (CAN oxidation of **2e**): 96%; bp 140–143 °C at 3 Torr (bulb to bulb); IR (neat): 1735 and 1655 cm^{-1} ; ^1H NMR (CDCl_3) δ : 0.83 (d, $J=7$ Hz, 3H), 0.99 (d, $J=7$ Hz, 3H), 2.03 (d, $J=1.6$ Hz, 3H), 1.91–2.38 (m, 1H), 3.65 (s, 1H), 3.74 (t, $J=7$ Hz, 1H), 6.60 (q, $J=1.6$ Hz, 1H), and 6.84 (s, 1H); Found: C, 66.04; H, 6.96%. Calcd for $\text{C}_{13}\text{H}_{16}\text{O}_4$: C, 66.08; H, 6.83%.

Methyl 2-(4-Methyl-3,6-dioxo-1,4-cyclohexadienyl)hexanoate 3f: (CAN oxidation of **2f**): 90%; IR (neat): 1735 and 1655 cm^{-1} ;

^1H NMR (CDCl_3) δ : 0.88 (t, $J=7$ Hz, 3H), 1.00–2.04 (m, 6H), 3.68 (s, 3H), 3.82 (t, $J=7$ Hz, 1H), 6.60 (q, $J=1.6$ Hz, 1H), and 6.68 (s, 1H); Found: C, 66.95; H, 7.38%. Calcd for $\text{C}_{14}\text{H}_{18}\text{O}_4$: C, 67.18; H, 7.25%.

Dimethyl 2-(4-Methyl-3,6-dioxo-1,4-cyclohexadienyl)succinate 3g: (CAN oxidation of **2g**): 80%; mp 85–86 °C (methanol–petroleum ether); IR (KBr): 1735 and 1640 cm^{-1} ; ^1H NMR (CDCl_3) δ : 2.06 (d, $J=1.6$ Hz, 3H), 2.61 (dd, $J=6$ and 17 Hz, 1H), 3.06 (dd, $J=6$ and 17 Hz, 1H), 3.66 (s, 3H), 3.68 (s, 3H), 4.12 (t, $J=7$ Hz, 1H), and 6.63 (m, 2H); Found: C, 58.57; H, 5.36%. Calcd for $\text{C}_{13}\text{H}_{14}\text{O}_6$: C, 58.64; H, 5.30%.

Dimethyl 2-(4-Methyl-3,6-dioxo-1,4-cyclohexadienyl)hexanediate 3h: (CAN oxidation of **2h**): 88%; bp 169–172 °C at 0.03 Torr (bulb to bulb); IR (neat): 1735 and 1655 cm^{-1} ; ^1H NMR (CDCl_3) δ : 1.36–2.05 (m, 4H), 2.05 (d, $J=1.6$ Hz, 3H), 2.33 (t, $J=7.5$ Hz, 2H), 3.64 (s, 3H), 3.67 (s, 3H and m, 1H), 6.60 (q, $J=1.6$ Hz, 1H), and 6.64 (s, 1H); Found: C, 61.18; H, 6.30%. Calcd for $\text{C}_{15}\text{H}_{18}\text{O}_6$: C, 61.21; H, 6.17%.

Methyl 2-(4-Methyl-3,6-dioxo-1,4-cyclohexadienyl)-6-(phthalimido)hexanoate 3i: (CAN oxidation of **2i**): 88%; IR (neat): 1775, 1735, 1710, and 1655 cm^{-1} ; ^1H NMR (CDCl_3) δ : 1.18–1.88 (m, 6H), 2.03 (d, $J=1.6$ Hz, 3H), 3.61 (s, 3H), 3.56–3.7 (m, 2H), 3.87 (m, 1H), 6.58 (q, $J=1.6$ Hz, 1H), 6.64 (d, $J=1.2$ Hz, 1H), and 7.60–7.82 (m, 4H); Found: C, 66.77; H, 5.42; N, 3.58%. Calcd for $\text{C}_{22}\text{H}_{21}\text{O}_6\text{N}$: C, 66.82; H, 5.35; N, 3.54%.

Methyl 2-(4-Methyl-3,6-dioxo-1,4-cyclohexadienyl)-6-(trifluoroacetamido)hexanoate 3j: (CAN oxidation of **2j**): 80%; IR (neat): 3320, 1710, and 1650 cm^{-1} ; ^1H NMR (CDCl_3) δ : 1.00–2.00 (m, 6H), 2.04 (d, $J=1.6$ Hz, 3H), 3.35 (q, $J=7$ Hz, 2H), 3.68 (s, 3H), 3.76 (m, 1H), 6.63 (q, $J=1.6$ Hz, 1H), 6.67 (s, 1H), and 6.68 (bs, 1H); Found: C, 52.87; H, 5.13; N, 3.84%. Calcd for $\text{C}_{16}\text{H}_{18}\text{O}_5\text{NF}_3$: C, 53.18; H, 5.02; N, 3.88%.

Methyl 2-(4-Methyl-3,6-dioxo-1,4-cyclohexadienyl)-4-(phenoxy-carbonyloxy)butyrate 3k: (CAN oxidation of **2k**): 71%; IR (neat): 1750 and 1655 cm^{-1} ; ^1H NMR (CDCl_3) δ : 2.05 (d, $J=1.6$ Hz, 3H), 2.05–2.59 (m, 2H), 3.72 (s, 3H), 3.88 (t, $J=7.5$ Hz, 1H), 4.28 (t, $J=6.5$ Hz, 2H), 6.63 (q, $J=1.6$ Hz, 1H), 6.71 (s, 1H), and 7.11–7.46 (m, 5H); Found: C, 63.29; H, 5.19%. Calcd for $\text{C}_{19}\text{H}_{18}\text{O}_7$: C, 63.68; H, 5.06%.

Methyl 2-(3,6-Dioxo-1,4-cyclohexadienyl)butyrate 3l: (CAN oxidation of **2l**): 54%; bp 131–134 °C at 3 Torr (bulb to bulb); IR (neat): 1730 and 1655 cm^{-1} ; ^1H NMR (CDCl_3) δ : 0.95 (t, $J=7$ Hz, 3H), 1.56–2.13 (m, 2H), 3.68 (s, 3H and m, 1H), 6.69 (m, 1H), and 6.75 (s, 2H); Found: C, 63.17; H, 6.06%. Calcd for $\text{C}_{11}\text{H}_{12}\text{O}_4$: C, 63.45; H, 5.81%.

Methyl 2-(4-Bromo-3,6-dioxo-1,4-cyclohexadienyl)butyrate 3m: (CAN oxidation of **2m**): 68%; bp 140–145 °C at 0.05 Torr (bulb to bulb); IR (neat): 1735 and 1660 cm^{-1} ; ^1H NMR (CDCl_3) δ : 0.95 (t, $J=7$ Hz, 3H), 1.56–2.15 (m, 2H), 3.62 (t, $J=7.5$ Hz, 1H), 3.68 (s, 3H), 6.88 (s, 1H), and 7.27 (s, 1H); Found: C, 45.81; H, 3.87; Br, 28.07%. Calcd for $\text{C}_{11}\text{H}_{11}\text{O}_4\text{Br}$: C, 46.06; H, 3.86; Br, 27.83%.

Methyl 2-(4-Methoxy-3,6-dioxo-1,4-cyclohexadienyl)butyrate 3n: (CAN oxidation of **2n**): 60%; mp 70–72 °C (petroleum ether); IR (KBr): 1740 and 1665 cm^{-1} ; ^1H NMR (CDCl_3) δ : 0.94 (t, $J=7$ Hz, 3H), 1.55–2.13 (m, 2H), 3.69 (s, 3H and m, 1H), 3.83 (s, 3H), 5.95 (s, 1H), and 6.64 (s, 1H); Found: C, 60.29; H, 5.99%. Calcd for $\text{C}_{12}\text{H}_{14}\text{O}_5$: C, 60.50; H, 5.92%.

Irradiation of Quinones 3. Irradiation of 3a: A solution of **3a** (416 mg, 2 mmol) in dry benzene (200 ml) was purged with nitrogen for 20 min and irradiated for 1 h. Evaporation of the solvent and column chromatography of the residue on silica gel with 10% ether–benzene as eluant gave methyl 2,3-dihydro-5-hydroxy-6-methylbenzofuran-2-carboxylate **4a** (171

mg, 41%); mp 153–154 °C (dichloromethane–petroleum ether); IR (KBr): 3420 and 1740 cm^{-1} ; ^1H NMR (CDCl_3) δ : 2.20 (s, 3H), 3.26 (dd, $J=7$ and 15 Hz, 1H), 3.51 (dd, $J=10$ and 15 Hz, 1H), 3.80 (s, 3H), 4.50 (s, 1H), 5.12 (dd, $J=7$ and 10 Hz, 1H), 6.60 (s, 1H), and 6.74 (s, 1H); MS, m/e (rel intensity), 208 (M^+ , 65), 176 (42), 149 (100); Found: C, 63.18; H, 5.89%. Calcd for $\text{C}_{11}\text{H}_{12}\text{O}_4$: C, 63.45; H, 5.81%.

Irradiation of 3b: A solution of **3b** (444 mg, 2 mmol) in dry benzene (200 ml) was purged with nitrogen for 20 min and irradiated for 1 h. After evaporation of the solvent, the residue was purified by chromatography on silica gel with 10% ether–benzene to give methyl 2,3-dihydro-5-hydroxy-3,6-dimethylbenzofuran-2-carboxylate **4b** (289 mg, 65%); mp 99–101 °C (dichloromethane–petroleum ether); IR (KBr): 3440 and 1745 cm^{-1} ; ^1H NMR (CDCl_3) δ : 1.40 (d, $J=7$ Hz, 3H), 2.18 (s, 3H), 3.40–3.68 (m, 1H), 3.79 (s, 3H), 4.63 (d, $J=7$ Hz, 1H), 4.71 (bs, 1H), 6.56 (s, 1H), and 6.63 (s, 1H); MS, m/e (rel intensity), 222 (M^+ , 100), 191 (17), 190 (17), 163 (68), 162 (91); Found: C, 64.85; H, 6.34%. Calcd for $\text{C}_{12}\text{H}_{14}\text{O}_4$: C, 64.85; H, 6.35%.

Irradiation of 3c: A solution of **3c** (222 mg, 1 mmol) in dry benzene (100 ml) was purged with nitrogen for 20 min and irradiated for 1 h. The solvent was evaporated, and the residue was chromatographed on silica gel with 10% ether–benzene to give methyl 2,3-dihydro-5-hydroxy-2,6-dimethylbenzofuran-2-carboxylate **4c** (100 mg, 45%); bp 170–180 °C at 0.01 Torr (bulb to bulb); IR (neat): 3420 and 1735 cm^{-1} ; ^1H NMR (CDCl_3) δ : 1.64 (s, 3H), 2.16 (s, 3H), 3.00 (d, $J=15$ Hz, 1H), 3.50 (d, $J=15$ Hz, 1H), 3.75 (s, 3H), and 6.56 (s, 2H); MS, m/e , 222 (M^+); Found: C, 68.69; H, 6.21%. Calcd for $\text{C}_{12}\text{H}_{14}\text{O}_4$: C, 68.85; H, 6.35%.

Irradiation of 3d: A solution of **3d** (473 mg, 2 mmol) in dry benzene (200 ml) was purged with nitrogen for 15 min and irradiated for 1 h. After evaporation of the solvent, the residue was chromatographed on silica gel with 10% ether–benzene to give methyl 3-ethyl-2,3-dihydro-5-hydroxy-6-methylbenzofuran-2-carboxylate **4d** (322 mg, 68%); mp 101–102 °C (dichloromethane–petroleum ether); IR (KBr): 3440 and 1740 cm^{-1} ; ^1H NMR (CDCl_3) δ : 1.00 (t, $J=7$ Hz, 3H), 1.40–2.00 (m, 2H), 2.19 (s, 3H), 3.42 (m, 1H), 3.76 (s, 3H), 4.51 (s, 1H), 4.74 (d, $J=5$ Hz, 1H), 6.58 (s, 1H), and 6.63 (s, 1H); MS, m/e (rel intensity), 236 (M^+ , 100), 205 (31), 204 (55); Found: C, 66.36; H, 7.07%. Calcd for $\text{C}_{13}\text{H}_{16}\text{O}_4$: C, 66.08; H, 6.83%.

Irradiation of 3e: A solution of **3e** (473 mg, 2 mmol) in dry benzene (200 ml) was purged with nitrogen for 20 min and irradiated for 1 h. The solvent was removed, and the crude oil was chromatographed on silica gel with 20% ether–benzene to give methyl 4-methyl-2-(2,5-dihydroxy-4-methylphenyl)-3-butenolate **4e** (189 mg, 40%); mp 130–131 °C (dichloromethane–petroleum ether); IR (KBr): 3400, 3340, 1715, and 900 cm^{-1} ; ^1H NMR (CDCl_3) δ : 1.78 (s, 3H), 2.15 (s, 3H), 3.77 (s, 1H), 4.38 (bs, 1H), 4.84 (s, 1H), 5.02 (bs, 1H), 6.60 (s, 1H), and 6.64 (s, 1H); MS, m/e (rel intensity), 236 (M^+ , 3), 204 (57), 175 (100).

Irradiation of 3f: A solution of **3f** (500 mg, 2 mmol) in dry benzene (200 ml) was purged with nitrogen for 10 min and irradiated for 1 h. Removal of the solvent left a dark oil which was subjected to column chromatography on silica gel with 10% ether–benzene. The only component isolated was methyl 2,3-dihydro-5-hydroxy-6-methyl-3-propylbenzofuran-2-carboxylate **4f** (300 mg, 60%); IR (neat): 3440 and 1745 cm^{-1} ; ^1H NMR (CDCl_3) δ : 0.94 (t, $J=7$ Hz, 3H), 1.20–1.82 (m, 4H), 2.18 (s, 1H), 3.46 (m, 1H), 3.77 (s, 3H), 4.54 (bs, 1H), 4.76 (d, $J=5$ Hz, 1H), 6.57 (s, 1H), and 6.63 (s, 1H); Found: C, 67.14; H, 7.50%. Calcd for $\text{C}_{14}\text{H}_{18}\text{O}_4$: C, 67.18; H, 7.25%.

Irradiation of 3g: A solution of **3g** (532 mg, 2 mmol) in dry

benzene (200 ml) was purged with nitrogen for 20 min and irradiated for 2 h. The solvent was removed, and the residue was subjected to column chromatography on silica gel with 10% ether–chloroform. The product isolated was dimethyl 2,3-dihydro-5-hydroxy-6-methylbenzofuran-2,3-dicarboxylate **4g** (299 mg, 56%): 180–181 °C (dichloromethane–petroleum ether); IR (KBr): 3340, 1735, and 1695 cm^{-1} ; ^1H NMR (CDCl_3) δ : 2.17 (s, 3H), 3.77 (s, 3H), 3.80 (s, 3H), 4.41 (d, $J=6$ Hz, 1H), 5.32 (bs, 1H), 5.56 (d, $J=6$ Hz, 1H), 6.65 (s, 1H), and 6.78 (s, 1H); MS, m/e (rel intensity), 266 (M^+ , 8), 264 (100); Found: C, 58.35; H, 5.24%. Calcd for $\text{C}_{13}\text{H}_{14}\text{O}_6$: C, 58.64; H, 5.30%.

Irradiation of 3h: A solution of **3h** (589 mg, 2 mmol) in dry benzene (300 ml) was purged with nitrogen for 10 min and irradiated for 1 h. The solution was concentrated, and the crude residue was purified by column chromatography on silica gel with 10% ether–chloroform to give methyl 2,3-dihydro-5-hydroxy-3-[2-(methoxycarbonyl)ethyl]-6-methylbenzofuran-2-carboxylate **4h** (300 mg, 51%): mp 136–137 °C (dichloromethane–petroleum ether); IR (KBr): 3480, 1750, and 1730 cm^{-1} ; ^1H NMR (CDCl_3) δ : 2.02 (m, 2H), 2.20 (s, 3H), 2.46 (t, $J=7$ Hz, 2H), 3.44–3.69 (m, 1H), 3.69 (s, 3H), 3.78 (s, 3H), 4.77 (d, $J=6$ Hz, 1H), 4.88 (s, 1H), 6.60 (s, 1H), and 6.67 (s, 1H); MS, m/e (rel intensity), 294 (M^+ , 26), 174 (100); Found: C, 60.93; H, 6.13%. Calcd for $\text{C}_{18}\text{H}_{18}\text{O}_6$: C, 61.21; H, 6.17%.

Irradiation of 3i: A solution of **3i** (791 mg, 2 mmol) in dry benzene (300 ml) was purged with nitrogen for 15 min and irradiated for 1 h. The solvent was removed, and the residue was chromatographed on silica gel with 10% ether–chloroform to give methyl 2,3-dihydro-5-hydroxy-6-methyl-3-[3-(phthalimido)propyl]benzofuran-2-carboxylate **4i** (593 mg, 75%): mp 180–181 °C (dichloromethane–petroleum ether); IR (KBr): 3480, 1775, 1745, and 1705 cm^{-1} ; ^1H NMR (CDCl_3) δ : 1.40–2.00 (m, 4H), 2.17 (s, 3H), 3.30–4.00 (m, 3H), 3.76 (s, 3H), 4.76 (d, $J=5$ Hz, 1H), 5.56 (bs, 1H), 6.63 (s, 2H), and 7.60–7.90 (m, 4H); MS, m/e (rel intensity), 395 (M^+ , 61), 203 (100); Found: C, 66.50; H, 5.40; N, 3.56%. Calcd for $\text{C}_{22}\text{H}_{21}\text{O}_6\text{N}$: C, 66.82; H, 5.35; N, 3.54%.

Irradiation of 3j: A solution of **3j** (723 mg, 2 mmol) in dry benzene (300 ml) was purged with nitrogen for 10 min and irradiated for 1 h. Removal of the solvent left a crude oil which was purified by column chromatography on silica gel using 10% ether–chloroform. The only component isolated was identified as methyl 2,3-dihydro-5-hydroxy-6-methyl-3-[3-(trifluoroacetamido)propyl]benzofuran-2-carboxylate **4j** (499 mg, 69%): IR (neat): 3300 and 1740–1700 cm^{-1} ; ^1H NMR (CDCl_3) δ : 1.69 (m, 4H), 2.17 (s, 3H), 3.38 (m, 3H), 3.77 (s, 3H), 4.71 (d, $J=5$ Hz, 1H), 6.57 (s, 1H), 6.63 (s, 1H), and 6.95 (bs, 1H); MS, m/e , 361 (M^+); Found: C, 52.80; H, 5.16; N, 3.82%. Calcd for $\text{C}_{16}\text{H}_{18}\text{O}_6\text{NF}_3$: C, 53.18; H, 5.02; N, 3.88%.

Irradiation of 3k: A solution of **3k** (300 mg, 0.83 mmol) in dry benzene (150 ml) was purged with nitrogen for 5 min and irradiated for 1 h. The solution was concentrated. The residue was chromatographed on silica gel with 10% ether–chloroform to give methyl 2,3-dihydro-5-hydroxy-6-methyl-3-(phenoxycarbonyloxymethyl)benzofuran-2-carboxylate **4k** (117 mg, 39%): IR (neat): 3440 and 1750 cm^{-1} ; ^1H NMR (CDCl_3) δ : 2.17 (s, 3H), 3.79 (s, 3H), 3.90 (m, 1H), 4.35 (m, 2H), 5.02 (d, $J=5.1$ Hz, 1H), 5.55 (bs, 1H), 6.61 (s, 1H), 6.67 (s, 1H), and 7.08–7.43 (m, 5H); Found: C, 63.50; H, 5.20%. Calcd for $\text{C}_{19}\text{H}_{18}\text{O}_7$: C, 63.68; H, 5.06%.

Irradiation of 3l: A solution of **3l** (416 mg, 2 mmol) in dry benzene (200 ml) was purged with nitrogen for 20 min and irradiated for 1 h. Removal of the solvent left an oil which was subjected to chromatography on silica gel with 10% ether–benzene. The product obtained was methyl 2,3-

dihydro-5-hydroxy-3-methylbenzofuran-2-carboxylate **4l** (250 mg, 60%): IR (neat): 3400 and 1735 cm^{-1} ; ^1H NMR (CDCl_3) δ : 1.38 (d, $J=7$ Hz, 3H), 3.53 (m, 1H), 3.79 (s, 3H), 4.64 (d, $J=7$ Hz, 1H), and 6.60 (s, 3H and bs, 1H); Found: C, 63.19; H, 5.81%. Calcd for $\text{C}_{11}\text{H}_{12}\text{O}_4$: C, 63.45; H, 5.81%.

Irradiation of 3m: A solution of **3m** (574 mg, 2 mmol) in dry benzene (200 ml) was purged with nitrogen for 20 min and irradiated for 1 h. Evaporation of the solvent gave a dark oil. The oil was chromatographed on silica gel to give methyl 6-bromo-2,3-dihydro-5-hydroxy-3-methylbenzofuran-2-carboxylate **4m** (178 mg, 31%): IR (neat): 3420 and 1735 cm^{-1} ; ^1H NMR (CDCl_3) δ : 1.41 (d, $J=7$ Hz, 3H), 3.53 (m, 1H), 3.78 (s, 3H), 4.65 (d, $J=7$ Hz, 1H), 6.74 (s, 1H), and 6.90 (s, 1H); MS, m/e (rel intensity), 287 (M^+ , 67); Found: C, 45.79; H, 3.95%. Calcd for $\text{C}_{11}\text{H}_{11}\text{O}_4\text{Br}$: C, 46.01; H, 3.86%.

Irradiation of 3n: A solution of **3n** (476 mg, 2 mmol) in dry benzene (200 ml) was purged with nitrogen for 10 min and irradiated for 1 h. Evaporation of the solvent and chromatographic separation of the crude oil by column chromatography on silica gel with 20% ether–benzene gave two compounds: The first fractions gave methyl 5-hydroxy-6-methoxy-3-methylbenzofuran-2-carboxylate (76 mg, 16%): mp 179–180 °C (dichloromethane–petroleum ether); IR (KBr): 3510 and 1720 cm^{-1} ; ^1H NMR (CDCl_3) δ : 2.51 (s, 3H), 3.94 (s, 6H), 5.58 (s, 1H), 6.98 (s, 1H), and 7.04 (s, 1H); Found: C, 60.72; H, 5.24%. Calcd for $\text{C}_{12}\text{H}_{12}\text{O}_5$: C, 61.01; H, 5.12%. The second fractions gave methyl 2,3-dihydro-5-hydroxy-6-methoxy-3-methylbenzofuran-2-carboxylate **4n** (200 mg, 42%): IR (neat): 3500 and 1735 cm^{-1} ; ^1H NMR (CDCl_3) δ : 1.41 (d, $J=7$ Hz, 3H), 3.56 (m, 1H), 3.80 (s, 3H), 3.83 (s, 3H), 4.66 (d, $J=7$ Hz, 1H), 5.32 (s, 1H), 6.49 (s, 1H), and 6.68 (s, 1H); Found: C, 60.25; H, 5.87%. Calcd for $\text{C}_{12}\text{H}_{14}\text{O}_5$: C, 60.50; H, 5.92%.

Preparation of Quinones 5 by Oxidative Cleavage^{a,b} of Photo-products 4. **Methyl 2-Hydroxy-3-(4-methyl-3,6-dioxo-1,4-cyclohexadienyl)propionate 5a:** A solution of **4a** (208 mg, 1 mmol) in 70% aq acetonitrile (5 ml) was treated with ceric ammonium nitrate 1.21 g, 2.2 mmol in water (3 ml). The mixture was stirred at 0 °C for 30 min, poured into water (100 ml), and extracted with chloroform (2 $\times$ 20 ml). The extract was washed with water, dried, and evaporated to give **5a** (174 mg, 78%): IR (neat): 3500, 1735, and 1650 cm^{-1} ; ^1H NMR (CDCl_3) δ : 2.03 (d, $J=1.6$ Hz, 3H), 2.69 (dd, $J=8.5$ and 15 Hz, 1H), 2.98 (dd, $J=5$ and 15 Hz, 1H), 3.81 (s, 3H), 4.18 (dd, $J=5$ and 8.5 Hz, 1H), 6.61 (q, $J=1.6$ Hz, 1H), and 6.71 (s, 1H); MS, m/e (rel intensity), 224 (M^+ , 6), 206 (12), 164 (86), 136 (100); Found: C, 58.79; H, 5.50%. Calcd for $\text{C}_{11}\text{H}_{12}\text{O}_5$: C, 58.92; H, 5.40%.

Methyl 2-Hydroxy-3-(4-methyl-3,6-dioxo-1,4-cyclohexadienyl)butyrate 5b: A solution of **4b** in aq acetonitrile was treated with CAN by the general procedure to give **5b** (80%): mp 104.5–105.5 °C (dichloromethane–petroleum ether); IR (KBr): 3480, 1725, and 1650 cm^{-1} ; ^1H NMR (CDCl_3) δ : 1.28 (d, $J=7$ Hz, 3H), 2.04 (d, $J=1.6$ Hz, 3H), 3.00 (bs, 1H), 3.46 (dq, $J=7$ and 4 Hz, 1H), 3.76 (s, 3H), 4.30 (bs, 1H), 6.60 (q, $J=1.6$ Hz, 1H), and 6.71 (s, 1H); Found: C, 60.54; H, 6.12%. Calcd for $\text{C}_{12}\text{H}_{14}\text{O}_5$: C, 60.50; H, 5.92%.

Methyl 2-Hydroxy-2-methyl-3-(4-methyl-3,6-dioxo-1,4-cyclohexadienyl)propionate 5c: A solution of **4c** in aq acetonitrile was treated with CAN by the same method to give **5c** (67%): mp 78 °C (petroleum ether); IR (KBr): 3480, 1720, and 1655 cm^{-1} ; ^1H NMR (CDCl_3) δ : 1.47 (s, 3H), 2.04 (d, $J=1.6$ Hz, 3H), 2.70 (d, $J=13.5$ Hz, 1H), 2.97 (d, $J=13.5$ Hz, 1H), 3.34 (s, 1H), 3.78 (s, 3H), 6.59 (q, $J=1.6$ Hz, 1H), and 6.70 (s, 1H); Found: C, 60.74; H, 5.98%. Calcd for $\text{C}_{12}\text{H}_{14}\text{O}_5$: C, 60.50; H, 5.92%.

Methyl 2-Hydroxy-3-(4-methyl-3,6-dioxo-1,4-cyclohexadienyl)-

pentanoate 5d: A solution of **4d** in aq acetonitrile was treated with CAN by the usual method to give **5d** (97%): mp 97–98 °C (petroleum ether); IR (KBr): 3420, 1720, and 1650 cm^{-1} ; ^1H NMR (CDCl_3) δ : 0.94 (t, $J=7.5$ Hz, 3H), 1.57–1.94 (m, 2H), 2.04 (d, $J=1.6$ Hz, 3H), 2.98 (d, $J=7.6$ Hz, 1H), 3.27 (dt, $J=3.5$ and 7 Hz, 1H), 3.74 (s, 3H), 4.36 (dd, $J=3.5$ and 7.6 Hz, 1H), 6.58 (q, $J=1.6$ Hz, 1H), and 6.71 (s, 1H); Found: C, 61.64; H, 6.38%. Calcd for $\text{C}_{13}\text{H}_{16}\text{O}_5$: C, 61.89; H, 6.39%.

Methyl 2-Hydroxy-3-(4-methyl-3,6-dioxo-1,4-cyclohexadienyl)-hexanoate 5f: A solution of **4f** in aq acetonitrile was treated with CAN by the usual manner to give **5f** (80%): mp 75–76 °C; IR (KBr): 3470, 1720, 1650, and 1645 cm^{-1} ; ^1H NMR (CDCl_3) δ : 0.90 (t, $J=7$ Hz, 3H), 1.31 (m, 2H), 1.64 (m, 2H), 2.02 (d, $J=1.6$ Hz, 3H), 2.87 (d, $J=7$ Hz, 1H), 3.31 (m, 1H), 3.92 (s, 3H), 4.31 (dd, $J=3.5$ and 7 Hz, 1H), 6.55 (q, $J=1.6$ Hz, 1H), and 6.68 (s, 1H); Found: C, 62.87; H, 6.73%. Calcd for $\text{C}_{14}\text{H}_{18}\text{O}_5$: C, 63.14; H, 6.81%.

Dimethyl 2-Hydroxy-3-(4-methyl-3,6-dioxo-1,4-cyclohexadienyl)-succinate 5g: A solution of **4g** in aq acetonitrile was treated with CAN by the general procedure to give **5g** (50%): IR (neat): 3470, 1735, and 1650 cm^{-1} ; ^1H NMR (CDCl_3) δ : 2.03 (d, $J=1.6$ Hz, 3H), 3.75 (s, 6H and 1H), 4.32 (d, $J=4.2$ Hz, 1H), 4.86 (d, $J=4.2$ Hz, 1H), 6.62 (q, $J=1.6$ Hz, 1H), and 6.87 (s, 1H); Found: C, 55.03; H, 5.23%. Calcd for $\text{C}_{13}\text{H}_{14}\text{O}_7$: C, 55.32; H, 5.00%.

Dimethyl 2-Hydroxy-3-(4-methyl-3,6-dioxo-1,4-cyclohexadienyl)-hexanedioate 5h: A solution of **4h** in aq acetonitrile was treated with CAN according to the general method to give **5h** (98%): IR (neat): 3460, 1730, and 1650 cm^{-1} ; ^1H NMR (CDCl_3) δ : 2.04 (d, $J=1.6$ Hz, 3H), 2.00–2.42 (m, 4H), 3.07 (bd, $J=8$ Hz, 1H), 3.38 (m, 1H), 3.68 (s, 3H), 3.37 (s, 3H), 4.35 (bs, 1H), 6.59 (q, $J=1.6$ Hz, 1H), and 6.74 (s, 1H); Found: C, 58.30; H, 6.01%. Calcd for $\text{C}_{15}\text{H}_{18}\text{O}_7$: C, 58.06; H, 5.85%.

Acetylation of Photoproducts 4b, 4d, 4f, and 4i. **Methyl 5-Acetoxy-2,3-dihydro-3,6-dimethylbenzofuran-2-carboxylate 6b:** The compound **4b** (111 mg, 0.5 mmol) was acetylated with acetic anhydride (5 ml) in the presence of pyridine (1 ml) to give **6b** (80%): mp 86–87 °C (petroleum ether); IR (KBr): 1750 cm^{-1} ; ^1H NMR (CDCl_3) δ : 1.45 (d, $J=7$ Hz, 3H), 2.13 (s, 3H), 2.30 (s, 3H), 3.64 (m, 1H), 3.83 (s, 3H), 4.73 (d, $J=7.2$ Hz, 1H), 6.75 (s, 1H), and 6.78 (s, 1H). Found: C, 63.50; H, 6.04%. Calcd for $\text{C}_{14}\text{H}_{16}\text{O}_5$: C, 63.62; H, 6.10%.

Methyl 5-Acetoxy-3-ethyl-2,3-dihydro-6-methylbenzofuran-2-carboxylate 6d: Acetylation of **4d** as described above gave **6d** (85%): mp 98–98.5 °C (dichloromethane–petroleum ether); IR (KBr): 1760 and 1745 cm^{-1} ; ^1H NMR (CDCl_3) δ : 1.01 (t, $J=7.5$ Hz, 3H), 1.78 (m, 2H), 2.13 (s, 3H), 2.31 (s, 3H), 3.50 (dt, $J=5.5$ and 7 Hz, 1H), 3.80 (s, 3H), 4.84 (d, $J=5.5$ Hz, 1H), 6.77 (s, 1H), and 6.82 (s, 1H); Found: C, 64.79; H, 6.49%. Calcd for $\text{C}_{15}\text{H}_{18}\text{O}_5$: C, 64.73; H, 6.52%.

Methyl 5-Acetoxy-2,3-dihydro-6-methyl-3-propylbenzofuran-2-carboxylate 6f: The compound **4f** was treated with acetic anhydride–pyridine to give **6f** (82%): IR (neat): 1760 and 1735 cm^{-1} ; ^1H NMR (CDCl_3) δ : 0.96 (t, $J=7$ Hz, 3H), 1.06–1.84 (m, 4H), 2.08 (s, 3H), 2.21 (s, 3H), 3.46 (m, 1H), 3.72 (s, 3H), 4.67 (d, $J=5$ Hz, 1H), 6.59 (s, 1H), and 6.65 (s, 1H); MS, m/e (rel intensity), 292 (M^+ , 4), 250 (100). Found: C, 65.59; H, 6.97%. Calcd for $\text{C}_{16}\text{H}_{20}\text{O}_5$: C, 65.74; H, 6.90%.

Methyl 5-Acetoxy-2,3-dihydro-6-methyl-3-[3-(phthalimido)-propyl]benzofuran-2-carboxylate 6i: The title compound was prepared from **4i** by acetylation with acetic anhydride in pyridine (77%): mp 128–129 °C (methanol); IR (KBr): 1760 and 1710 cm^{-1} ; ^1H NMR (CDCl_3) δ : 1.40–2.00 (m, 4H), 2.10 (s, 3H), 2.28 (s, 3H), 3.47–4.00 (m, 3H), 3.76 (s, 3H), 4.81 (d, $J=5$ Hz, 1H), 6.72 (s, 1H), 6.80 (s, 1H), and 6.80 (s, 1H), and 7.60–7.90 (m, 4H); Found: C, 65.62; H, 5.24; N, 3.11%.

Calcd for $\text{C}_{24}\text{H}_{23}\text{O}_7\text{N}$: C, 65.89; H, 5.30; N, 3.20%.

Methyl 2-Acetoxy-3-(2,5-diacetoxy-4-methylphenyl)hexanoate 7: A solution of **5f** (133 mg, 0.5 mmol) in ether (50 ml) was reduced with excess aq sodium dithionite to give methyl 2-hydroxy-3-(2,5-dihydroxy-4-methylphenyl)hexanoate, which was acetylated with acetic anhydride (2 ml) in the presence of pyridine (0.1 ml) to afford **7** (118 mg, 60%): IR (neat): 1760 cm^{-1} ; ^1H NMR (CDCl_3) δ : 0.86 (t, $J=7$ Hz, 3H), 1.27 (m, 2H), 1.64 (m, 2H), 2.10 (s, 3H), 2.16 (s, 3H), 2.33 (s, 6H), 3.46 (m, 1H), 3.65 (s, 3H), 5.18 (d, $J=5$ Hz, 1H), 6.96 (s, 1H), and 7.09 (s, 1H); MS, m/e (rel intensity), 394 (M^+ , 1), 352 (38), and 310 (100); Found: C, 60.66; H, 6.87%. Calcd for $\text{C}_{20}\text{H}_{26}\text{O}_8$: C, 60.90; H, 6.64%.

The author is grateful to Professor Kazuhiro Maruyama for his invaluable suggestions to this work and his continuous encouragement.

References

- 1) A. G. Shultz, Y. K. Yee, and M. H. Berger, *J. Am. Chem. Soc.*, **99**, 8065 (1977) and references cited therein.
- 2) For reviews of quinones and their chemistry, see: "The Chemistry of the Quinonoid Compounds," ed by S. Patai, Wiley, New York (1974), Parts 1 and 2; J. M. Bruce, *Quart. Rev.*, **21**, 405 (1967).
- 3) a) C. M. Orland, Jr., H. Mark, A. K. Bose, and M. S. Manhas, *J. Org. Chem.*, **32**, 2512 (1968); b) J. M. Bruce and K. Dawes, *J. Chem. Soc., C*, **1970**, 645; c) J. M. Bruce and D. Creed *ibid.*, **1970**, 649; d) J. M. Bruce, D. Creed, and K. Dawes, *ibid.*, **1971**, 2244; e) J. M. Bruce, D. Creed, and K. Dawes, *ibid.*, **1971**, 3748; f) J. M. Bruce and A.-u.-h. Chaudhry *J. Chem. Soc., Perkin Trans. 1*, **1972**, 372.
- 4) In a preliminary report of this work, the author erroneously assigned the positions of substituent groups for dihydrobenzofuran structure of these several photoproducts, it should be corrected to the photorearranged dihydrobenzofuran structure **4**: K. Maruyama and T. Kozuka, *Chem. Lett.*, **1980**, 341.
- 5) a) K. Maruyama and T. Kozuka, *Bull. Chem. Soc. Jpn.*, **51**, 3586 (1978); b) E. C. Taylor, R. L. Robey, K.-T. Liu, B. Favre, H. T. Bozino, R. D. Conley, C.-S. Chiang, A. McKillop, and M. E. Ford, *J. Am. Chem. Soc.*, **98**, 3037 (1976); c) E. C. Taylor, C.-S. Chiang, A. McKillop, and J. F. White, *ibid.*, **98**, 750 (1976).
- 6) a) P. Jacob, III, P. S. Callery, A. T. Shulgin, and N. Catagnoli, Jr., *J. Org. Chem.*, **41**, 3627 (1976); b) R. L. Hannan, R. B. Barker, and H. Rapoport, *ibid.*, **44**, 2153 (1979).
- 7) Ref. 3d and references cited therein the 2- and 3-substituted 2,3-dihydrobenzofuran compounds have $J_{3A,3B} \approx ca. 15$ and $J_{2A,2B} \approx ca. 8$ Hz, respectively.
- 8) Ref 3e.
- 9) L. M. Jackman, "Applications of Nuclear Magnetic Resonance Spectroscopy in Organic Chemistry," 2nd ed, Pergamon Press, New York (1967), p. 316.
- 10) a) J. Gripenberg and T. Hase, *Acta Chem. Scand.*, **20**, 1561 (1966); b) C. H. Ludwig, B. J. Nist, and J. L. McCarthy, *J. Am. Chem. Soc.*, **86**, 1186 (1964); c) G. Aulin-Erdtman and Y. Tomita, *Acta Chem. Scand.*, **17**, 535 (1963).
- 11) a) D. P. Brust, D. S. Tarbell, S. M. Hecht, E. C. Hayward, and L. D. Colebrook, *J. Org. Chem.*, **31**, 2196 (1966); b) E. C. Hayward, D. S. Tarbell, and L. D. Colebrook, *ibid.*, **33**, 399 (1968); c) L. H. Zalkow and M. Ghosal, *ibid.*, **34**, 1646 (1969); d) M. P. Mertes and L. J. Powes, *ibid.*, **36**, 1805 (1971). The coupling constants for 2,3-dialkyl 2,3-dihydrobenzofuran were $J_{cis} = 7.0$ Hz and $J_{trans} = 4.8$ Hz.

- 12) M. Karplus, *J. Am. Chem. Soc.*, **85**, 2870 (1963).
 - 13) S. Farid, *J. Chem. Soc., Chem. Commun.*, **1971**, 73.
 - 14) S. Farid, *J. Chem. Soc., Chem. Commun.*, **1970**, 303.
 - 15) *Org. Synth.*, Coll. Vol. III, 169 (1955).
 - 16) *Org. Synth.*, Coll. Vol. IV, 556 (1963).
 - 17) S. Gabriel and J. Colman, *Chem. Ber.*, **41**, 2014 (1908).
 - 18) E. E. Schallenberg and M. Calvin, *J. Am. Chem. Soc.*, **77**, 2779 (1955).
 - 19) B.-T. Ho, L. W. Tansey, and W. M. McIssac, *J. Med. Chem.*, **13**, 1022 (1970).
 - 20) T. R. Ingle, N. L. Phalnikar, and B. V. Bhide, *J. Indian Chem. Soc.*, **26**, 569 (1949).
 - 21) E. Hardegger, H. P. Knowpfel, E. Widmer, H. Corrodi, Th. Schmidt, H. P. Knoepfel, W. Reder, H. J. Meyer, F. Kugler, and H. Gempeler, *Helv. Chem. Acta*, **47**, 1996 (1964).
 - 22) R. B. Moffett, A. R. Hanze, and P. H. Seay, *J. Med. Chem.*, **7**, 178 (1964).
-