

A Simple and Convenient Method for the Synthesis of Sulfones Using Polyethylene Glycols or Their Dialkyl Ethers as Solvents or Catalysts


Kazuaki SUKATA

Nihon Tokushu Kagaku Kogyo Co., Ltd., 3-36, Tohda-cho, Moriguchi, Osaka 570

(Received June 2, 1983)

Synopsis. Various alkyl *p*-tolyl sulfones were prepared in good yields under mild conditions in the presence of polyethylene glycols or their dialkyl ethers as solvents or catalysts.

It has been reported¹⁾ that the alkylation of sodium *p*-toluenesulfinate in methanol (MeOH) or *N,N*-dimethylformamide gives both the *p*-toluenesulfinic ester and the sulfone, and alkylation with hard alkylating agents gives predominantly the ester, whereas a soft alkylating agent gives predominantly the sulfone. This alkylation usually gives the sulfone only in moderate yields.¹⁾



Recently, Veenstra and Zwanenburg²⁾ have reported the reaction of tetrabutylammonium *p*-toluenesulfinate with alkyl halides as an improved method for the synthesis of sulfones. More recently, Manescalchi *et al.*³⁾ have reported a similar method using Amberlyst

A-26, a macroreticular anion-exchange resin. In these methods, however, the tedious preparation of tetrabutylammonium *p*-toluenesulfinate or the sulfinate form of Amberlyst A-26 is required before reaction with the alkyl halide.

In this paper, a simpler and more convenient method for the synthesis of sulfones using polyethylene glycols (PEG) or their diethyl ethers is reported.

The reactions were carried out with sodium *p*-toluenesulfinate monohydrate and various alkyl halides in PEG having an average molecular weight of 400 (PEG-400), PEG-400 diethyl ether (PEG-400-Et₂), or MeOH containing PEG-1000 or PEG-1000-Et₂ as a catalyst. Table 1 shows the results. In all cases, the sulfones were obtained in good to excellent yields. When PEG-400 was employed as a solvent, GLC showed that the product was almost entirely the sulfone because of the rapid hydrolysis of the ester. In the case of PEG-400-Et₂ or MeOH with a catalytic amount of PEG-1000 or PEG-1000-Et₂, however, GLC showed that a small amount of the ester remained under the experimental conditions used. These facts indicate that PEG is a good solvent for the alkylation of sulfinate anion to yield sulfones.

TABLE 1. REACTION OF C₇H₇SO₂Na·H₂O WITH ALKYL HALIDES^{a)}

Alkyl halide	Solvent	Catalyst ^{b)}	Time h	Temp °C	Yield ^{c)} %	Mp θ_m or Bp θ_b /°C (mmHg)	Lit Mp θ_m or Bp θ_b /°C (mmHg)
C ₂ H ₅ Br	PEG-400	—	3	60	89		
	PEG-400-Et ₂	—	4	60	89	53—55	55—56 ⁵⁾
	MeOH	A	7	reflux	80 ^{d)}	151—152(3)	
	MeOH	—	7	reflux	64 ^{e)}		
<i>n</i> -C ₄ H ₉ Br	PEG-400	—	3	60	90		
	PEG-400-Et ₂	—	9	60	82 ^{d)}	143—144(1)	175—177(4) ⁶⁾
	MeOH	A	7	60	83 ^{d)}		
	MeOH	—	7	60	61 ^{e)}		
<i>n</i> -C ₈ H ₁₇ Br	PEG-400	—	4	60	94		
	PEG-400-Et ₂	—	12	60	59 ^{d)}	40.0—40.5	^{f)}
	MeOH	A	7	60	80	150—151(0.12)	
	MeOH	—	7	60	56 ^{d)}		
CH ₂ =CHCH ₂ Br	PEG-400	—	1.5	60	95		
	PEG-400-Et ₂	—	3	60	95	51—52	52—53 ⁷⁾
	MeOH	A	3	60	89 ^{d)}	137(1)	
PhCH ₂ Cl ^{g)}	PEG-400	—	2	60	94		
	MeOH	A	4	60	90	144—145	144—145 ⁸⁾
ClCH ₂ CN	PEG-400	—	2	60	93		
	MeOH	B	4	60	83	147—148	145—146 ¹⁰⁾
BrCH ₂ Br	PEG-400	—	6	60	73		
	PEG-400	—	4	80	89	87—88	90—92 ⁹⁾

a) All reactions were carried out with C₇H₇SO₂Na·H₂O (0.05 mol) and alkyl halide (0.075 mol) in a solvent (20 ml) with or without a catalyst (0.005 mol). b) A: PEG-1000-Et₂, B: PEG-1000. c) Isolated yields. Unless otherwise noted, purities were >99% by GLC. d) Purities were 98% by GLC. e) Purities were 97% by GLC. f) See Experimental section. g) Forty milliliters of solvent were used.

Experimental

Materials. Methanol and PEG-400 (reagent grade, Kishida Kagaku Co., Ltd.) were dried with molecular sieves 4A. PEG-1000 (reagent grade, Kishida Kagaku Co., Ltd.) was used without further purification. PEG-400-Et₂ and PEG-1000-Et₂ were prepared according to the method previously reported.⁴ Sodium *p*-toluenesulfinate (C₇H₇SO₂Na n H₂O, $n \approx 1$) (reagent grade, Tokyo Kasei Kogyo Co., Ltd.) was used as obtained. The physical properties (bp or mp) of ethyl *p*-tolyl sulfone,⁵ butyl *p*-tolyl sulfone,⁶ allyl *p*-tolyl sulfone,^{2,7} benzyl *p*-tolyl sulfone,⁸ cyanomethyl *p*-tolyl sulfone,^{2,10} and bromomethyl *p*-tolyl sulfone^{2,9} were in good agreement with those reported in the literature (Table 1) and satisfactory IR spectra for these compounds were obtained.

Typical Procedure for the Reaction of Sodium *p*-Toluenesulfinate Monohydrate with Alkyl Halides.

Bromomethyl *p*-Tolyl Sulfone: A mixture of powdered sodium *p*-toluenesulfinate monohydrate (9.8 g, 0.05 mol) and dibromomethane (13.0 g, 0.075 mol) was stirred at 80°C for 4 h in PEG-400 (20 ml). After removal of the excess dibromomethane, the reaction mixture was poured into water (300 ml). The precipitated white solid was collected and recrystallized from ethanol-water, giving 11.1 g (89%) of bromomethyl *p*-tolyl sulfone, mp 87–88°C (lit.⁹ mp 90–92°C).

Butyl *p*-Tolyl Sulfone: A mixture of powdered sodium *p*-toluenesulfinate monohydrate (9.8 g, 0.05 mol), butyl bromide (10.3 g, 0.075 mol), MeOH (20 ml), and PEG-1000-Et₂ (5.3 g, 0.005 mol) was stirred at 60°C for 7 h. The reaction mixture was diluted with water (200 ml) and extracted with benzene (20 ml $\times$ 3). The benzene layers were combined, washed with water, and dried. After removal of the benzene, the residual oil was distilled under reduced pressure, giving 8.8 g (83%) of butyl *p*-tolyl sulfone, bp 137–142°C/1 mmHg (1 mmHg = 133.322 Pa) (lit.⁶ bp 175–177°C/4 mmHg). The purity was 98% by GLC.

Octyl *p*-Tolyl Sulfone: A mixture of powdered sodium *p*-toluenesulfinate monohydrate (9.8 g, 0.05 mol), octyl bromide (14.5 g, 0.075 mol), and PEG-400 (20 ml) was stirred at 60°C for 3 h. After the reaction mixture had been treated according to the method for butyl *p*-tolyl sulfone, the residual oil was distilled under reduced pressure, giving 12.6 g (94%) of octyl *p*-tolyl sulfone (>99% pure), bp 175–178°C/0.5 mmHg. Redistillation of the product gave pure octyl *p*-tolyl sulfone having bp 150–151°C/0.12 mmHg, mp 40.0–40.5°C; IR (neat) 1310, 1140 cm⁻¹ (–SO₂–). Found: C, 66.90; H, 9.07; S, 11.93%. Calcd for C₁₅H₂₄O₂S: C, 67.12; H, 9.01; S, 11.94%.

The author thanks Nihon Tokushu Kagaku Kogyo Co., Ltd. for permission to publish this article, and also Dr. Nobuaki Wada for useful suggestions during this work.

References

- 1) J. S. Meek and J. S. Fowler, *J. Org. Chem.*, **33**, 3422 (1968).
- 2) G. H. Veenstra and B. Zwanenburg, *Synthesis*, **1975**, 519.
- 3) F. Manescalchi, M. Orena, and D. Savoia, *Synthesis*, **1979**, 445.
- 4) K. Sukasa, *Yuki Gosei Kagaku Kyokai Shi*, **39**, 86 (1981).
- 5) R. Otto, *Ber.*, **13**, 1277 (1880).
- 6) H. Gilman and N. J. Beaver, *J. Am. Chem. Soc.*, **47**, 1450 (1925).
- 7) R. Otto, *Justus Liebigs Ann. Chem.*, **283**, 184 (1894).
- 8) R. Otto, *Ber.*, **13**, 1278 (1880).
- 9) R. Otto, *J. Prakt. Chem.*, [2], **40**, 544 (1889).
- 10) J. Tröger and W. Hille, *J. Prakt. Chem.* [2], **71**, 226 (1905).