

An Easy Preparation of Triphenylmethyl Carboxylates


Shun-ichi HASHIMOTO, Masahiko HAYASHI, and Ryoji NOYORI*

Department of Chemistry, Nagoya University, Chikusa, Nagoya 464

(Received December 27, 1983)

Synopsis. A new recipe for triphenylmethylation of carboxylic acids and phenol is disclosed.

Triphenylmethylation of carboxylic acids has been conventionally achieved by heating triphenylmethyl bromide and metal carboxylates in nonpolar solvents.¹⁾ Sensitive, polymerizable acrylic acids, however, can be esterified only using the expensive silver salts.²⁾ We found that condensation of economically accessible trimethylsilyl carboxylates (**1**) and triphenylmethyl fluoride (**2**) proceeds smoothly at room temperature in acetonitrile containing a catalytic amount of tetrafluorosilane (**3**). This facile procedure allows the easy synthesis of triphenylmethyl methacrylate (**4a**),³⁾ an important material for the preparation of the chiral liquid chromatographic stationary phases.^{4,5)}



a: R = CH₂=C(CH₃), **b:** R = *n*-C₉H₁₉, **c:** R = C₆H₅

In a like manner, the tetrafluorosilane-catalyzed reaction of phenyl trimethylsilyl ether and the fluoride **2** gave phenyl triphenylmethyl ether in a high yield.

Experimental

All melting points were uncorrected. IR spectra were recorded on a JASCO IRA-1 spectrometer. ¹H NMR spectra were measured with a JEOL JNM FX-90Q (90 MHz) spectrometer using tetramethylsilane as an internal standard. Mass spectra were determined with a Hitachi RMU-6C mass spectrometer, operating with an ionization energy of 70 eV. Tetrafluorosilane (**3**) was purchased from Matheson Co. Triphenylmethyl fluoride (**2**) was prepared by the reported method.⁶⁾ Acetonitrile was distilled from CaH₂ before use. All products are known compounds and gave consistent spectral data.

Triphenylmethyl Methacrylate (4a). To a solution of **1a** (1.25 g, 7.91 mmol) and **2** (2.07 g, 7.91 mmol) in dry acetonitrile (2 cm³) was added a 0.08 mol dm⁻³ solution of **3** in acetonitrile (4.9 cm³, 0.396 mmol) at 14 °C under argon atmosphere. After 6-h stirring at this temperature, the resulting yellow mixture was poured into a vigorously stirred, pH 7.4 phosphate buffer (80 cm³) containing KF (8.0 g) which was kept at 0 °C. The whole mixture was extracted with a 2:1 mixture of ether and hexane (50 cm³×2), and the combined organic layers were washed with aq NaHCO₃ and brine, dried over anhydrous Na₂SO₄, and evaporated *in vacuo*. The residual solid was recrystallized from hexane to give **4a** (2.42 g, 94% yield) as colorless prisms: Mp 98–99 °C (lit.⁷⁾ 101–102 °C; IR (CCl₄) 1725, 1490, 1140 cm⁻¹; ¹H NMR (CDCl₃) δ=1.99 (3H, dd, *J*=1.0, 1.4 Hz), 5.60 (1H, m), 6.23 (1H, m), 7.2–7.5 (15H, m); MS *m/z* 328 (M⁺), 260, 259, 243, 183, 165, 105, 85, 83, 77.

The above reaction was carried out in acetonitrile-*d*₃ for 6 h, and the reaction mixture was subjected to ¹H NMR analysis, which indicated the production of the theoretical amount of trimethylfluorosilane⁸⁾ (δ=0.21, d, *J*_{CH₃F}=7.5 Hz) in addition to **4a**.⁹⁾

Triphenylmethyl Decanoate (4b). A mixture of **1b** (232 mg, 0.95 mmol), **2** (249 mg, 0.95 mmol), and **3** (0.08 mol dm⁻³ in acetonitrile, 0.62 cm³, 0.05 mmol) in acetonitrile (2.5 cm³) was stirred at 16 °C for 6 h. The reaction mixture was worked up as described above, and the oily residue was chromatographed on silica gel, which had been treated with triethylamine (2%), using a 5:1 mixture of hexane and ether as an eluent. Pure **4b** was obtained as a colorless oil (361 mg, 88%); IR (CCl₄) 1760, 1500, 1150 cm⁻¹; ¹H NMR (CDCl₃) δ=0.87 (3H, br t, *J*=6.0 Hz), 1.07–1.73 (14H, m), 2.43 (2H, t, *J*=6.8 Hz), 7.1–7.4 (15H, m); MS *m/z* 414 (M⁺), 260, 259, 244, 243, 165, 78, 77.

Triphenylmethyl Benzoate (4c). A mixture of **1c** (372 mg, 1.92 mmol), **2** (503 mg, 1.92 mmol), and **3** (0.08 mol dm⁻³ in acetonitrile, 1.2 cm³, 0.1 mmol) in acetonitrile (4 cm³) was stirred at 16 °C for 4 h. Workup followed by recrystallization from a 10:1 mixture of hexane and benzene afforded **4c** (671 mg, 96%) as colorless plates: Mp 170–171 °C (lit.¹⁾ 168–170 °C; IR (CCl₄) 1740, 1600, 1500, 1270 cm⁻¹; ¹H NMR (CDCl₃) δ=7.1–7.6 (18H, m), 8.13 (2H, *J*=1.5, 7.8 Hz); MS *m/z* 364 (M⁺), 260, 259, 244, 243, 85, 83.

Phenyl Triphenylmethyl Ether. A mixture of phenyl trimethylsilyl ether (166 mg, 1.0 mmol), **2** (262 mg, 1.0 mmol), and **3** (0.08 mol dm⁻³ in acetonitrile, 0.63 cm³, 0.05 mmol) in acetonitrile (2 cm³) was stirred at 15 °C for 24 h. Aqueous workup followed by column chromatography on 2% triethylamine-treated silica gel using a 30:1 mixture of hexane and ether as an eluent, afforded the product (286 mg, 85%) as a colorless oil. This oil crystallized as colorless prisms from ethanol solution: Mp 103–104 °C (lit.¹⁰⁾ 103 °C; IR (CCl₄) 1600, 1500, 1450, 1220 cm⁻¹; ¹H NMR (CDCl₃) δ=6.6–7.1 (5H, m), 7.1–7.5 (15H, m); MS *m/z* 244, 243, 165, 94.

References

- 1) K. D. Berlin, L. H. Gower, J. W. White, D. E. Gibbs, and G. P. Sturm, *J. Org. Chem.*, **27**, 3595 (1962).
- 2) H. Yuki, K. Hatada, Y. Kikuchi, and T. Niinomi, *J. Polym. Sci. Part B*, **6**, 753 (1968).
- 3) For the preparation of **4a** from **1a** and triphenylmethyl trimethylsilyl ether, see: S. Murata and R. Noyori, *Tetrahedron Lett.*, **22**, 2107 (1981).
- 4) Y. Okamoto, K. Suzuki, K. Ohta, K. Hatada, and H. Yuki, *J. Am. Chem. Soc.*, **101**, 4763 (1979); H. Yuki, Y. Okamoto, and I. Okamoto, *ibid.*, **102**, 6356 (1980); Y. Okamoto, I. Okamoto, and H. Yuki, *Chem. Lett.*, **1981**, 835; Y. Okamoto, I. Okamoto, and H. Yuki, *J. Polym. Sci. Polym. Lett. Ed.*, **19**, 451 (1981); Y. Okamoto, S. Honda, I. Okamoto, H. Yuki, S. Murata, R. Noyori, and H. Takaya, *J. Am. Chem. Soc.*, **103**, 6971 (1981); R. Noyori, N. Sano, S. Murata, Y. Okamoto, H. Yuki, and T. Ito, *Tetrahedron Lett.*, **23**, 2969 (1982); Y. Kawada, H. Iwamura, Y. Okamoto, and H. Yuki, *ibid.*, **24**, 791 (1983).
- 5) A column of (+)-poly(triphenylmethyl methacrylate) coated on macroporous silica gel (Chiralpak OT) is available from Japan Spectroscopic Co. Ltd., Ishikawa-cho, Hachioji,

Tokyo.

- 6) G. A. Olah, J. T. Welch, Y. D. Vankar, M. Nojima, I. Kerekes, and J. A. Olah, *J. Org. Chem.*, **44**, 3872 (1979).
 - 7) N. A. Adrova and L. K. Prokhorova, *Vysokomol. Soedin.*, **3**, 1509 (1961); *Chem. Abstr.*, **56**, 10384 (1962).
 - 8) A. E. Newkirk, *J. Am. Chem. Soc.*, **68**, 2736 (1946).
 - 9) Other Lewis acids such as BF_3 , AlCl_3 (in CH_3CN), and TiCl_4 (in CH_2Cl_2) also acted as catalyst of this reaction.
 - 10) A. Baeyer, *Ber.*, **42**, 2625 (1909).
-