

2-Phenyl-1,2-oxaphospholane 1a: To a mixture of dichlorophenylphosphine (2.01 g, 11.2 mmol), pyridine (0.95 ml, 11.8 mmol), and 10 ml of THF was added 3-chloro-1-propanol (1.06 g, 11.2 mmol) in 4 ml of THF at 0 °C. After refluxing for 1 h, the precipitate of pyridinium hydrochloride was removed by filtration. The filtrate was added to 8 ml of THF containing small pieces of lithium metal (0.172 g 24.8 mg-atom) at -78 °C and stirred overnight gradually up to room temperature. THF was evaporated *in vacuo*, and 30 ml of diethyl ether was added to the residue to extract the product. The mixture was well stirred and filtered. After concentration of the filtrate, the residue was

subjected to distillation *in vacuo* to give 0.69 g of **1a** (37% yield), bp 62–65 °C (0.13 Torr)³. ³¹P{¹H}NMR(CDCl₃) +110.2 ppm (relative to 85% H₃PO₄ external standard); ¹H NMR δ(CDCl₃) 1.4–2.5 (m, 4H, PCH₂CH₂), 3.6–4.3 (m, 2H, OCH₂), and 7.1–7.4 ppm (m, 5H, C₆H₅); ¹³C{¹H}NMR(CDCl₃) 23.3 (s, PC₂H₂), 32.2 (d, J_{CCP}=20.9 Hz, PCH₂C₂H₂), 72.4 (d, J_{COF}=14.0 Hz, OCH₂), 126.9 (s, P-Ph; δ-C), 128.0 (d, J_{CCP}=14.8 Hz, P-Ph; β-C), 128.1 (s, P-Ph; γ-C), and 144.8 ppm (d, J_{CP}=36.7 Hz, P-Ph; α-C).

These procedures for the preparation of **1a** were similarly applied to those of **1b–d**.

2-Phenyl-1,2-oxaphosphorinane 1b: From dichlorophenylphosphine (2.19 g, 12.2 mmol), 4-chloro-1-butanol (1.33 g, 12.2 mmol), pyridine (1.04 ml, 12.8 mmol), and lithium metal (0.186 g 26.8 mg-atom), 0.39 g of **1b** was obtained; 18% yield, bp 76–79 °C (0.55 Torr). ³¹P{¹H}NMR(CDCl₃) +111.1 ppm; ¹H NMR δ(CDCl₃) 1.2–2.4 (m, 6H, PCH₂CH₂CH₂CH₂), 3.4–3.9 (m, 2H, OCH₂), and 7.1–7.5 (m, 5H, C₆H₅); ¹³C{¹H}NMR(CDCl₃) 19.3 (d, J_{CP}=3.5 Hz, PC₂H₂), 25.5 (d, J_{CCP}=22.6 Hz, PCH₂C₂H₂), 27.6 (d, J_{CCOP}=3.5 Hz, OCH₂C₂H₂), 63.4 (d, J_{COF}=7.8 Hz, OCH₂), 127.8 (s, P-Ph; δ-C), 128.6 (d, J_{CCCP}=2.6 Hz P-Ph; γ-C), 129.0 (d, J_{CCP}=14.1 Hz, P-Ph; β-C), and 140.7 (d, J_{CP}=31.4 Hz, P-Ph; α-C); *m/z* 180 (M⁺, 69%).

2-Phenyl-1,2-thiaphospholane 1c: From dichlorophenylphosphine (48 ml, 0.355 mol), 3-chloro-1-propanethiol (39.2 g, 0.355 mol), pyridine (30 ml, 0.371 mol), and lithium metal (4.93 g, 0.710 g-atom), 6.59 g of **1c** was obtained; 10% yield, bp 102–104 °C (0.50 Torr). ³¹P{¹H}NMR(CDCl₃) +22.3 ppm; ¹H NMR δ(CDCl₃) 1.1–3.1 (m, 6H, SCH₂CH₂CH₂P) and 6.8–7.6 (m, 5H, C₆H₅); ¹³C{¹H}NMR(CDCl₃) 28.9 (d, J_{CSP}=7.3 Hz, SC₂H₂), 35.2 (s, PC₂H₂), 36.0 (d, J_{CCP}=23.2 Hz, PCH₂C₂H₂), 128.1 (s, P-Ph; δ-C), 128.1 (d, J_{CCCP}=3.7 Hz, P-Ph; γ-C), 130.0 (d, J_{CCP}=17.1 Hz, P-Ph; β-C), and 140.4 (d, J_{CP}=40.3, P-Ph, α-C); *m/z* 182 (M⁺, 100%).

2-Phenyl-1,2-thiaphosphorinane 1d: From dichlorophenylphosphine (12.65 g, 0.071 mol), 4-chloro-1-butanethiol (9.52 g, 0.071 mol), pyridine (6.0 ml, 0.074 mol), and lithium metal (0.985 g, 0.142 g-atom), 0.659 g of **1d** was obtained; 5% yield, bp 99–103 °C (0.32 Torr). ³¹P{¹H}NMR(CDCl₃) +4.8 ppm; ¹H NMR δ(CDCl₃) 1.0–3.0 (m, 8H, SC₂H₂CH₂CH₂CH₂P) and 6.9–7.8 (m, 5H, C₆H₅); ¹³C{¹H}NMR(CDCl₃) 20.1 (s, PC₂H₂ or PCH₂CH₂C₂H₂), 23.5 (d,

J_{CCP}=22.6 Hz, PCH₂C₂H₂), 24.5 (d, J_{CSP}=10.5 Hz, SC₂H₂), 27.8 (s, PC₂H₂ or PCH₂CH₂C₂H₂), 127.2 (s, P-Ph; δ-C), 128.6 (d, J_{CCCP}=3.5 Hz, P-Ph; γ-C), 130.5 (d, J_{CCP}=14.8, P-Ph; β-C), and 137.1 ppm (d, J_{CP}=32.2 Hz, P-Ph; α-C); *m/z* 196 (M⁺, 100%).

References

- 1) S. Kobayashi, M. Suzuki, and T. Saegusa, *Polymer Bull.*, **4**, 315 (1981); *ibid.*, **8**, 417 (1982); *Macromolecules*, **17**, 107 (1984).
- 2) A trivial name of "deoxophostone" for **1a** was used by Grayson and Farley.³ A "phostone" is recognized as a derivative of lactone.⁴ We followed logically these naming for new phosphorus heterocycles **1c** and **1d** as "deoxothiolphostones". Then, the inter-relationships are schematically shown as follows:

thiolactone
thiolphostone
deoxothiolphostone
- 3) M. Grayson and C. E. Farley, *J. Chem. Soc., Chem. Commun.*, **1967**, 830. This literature reported bp of **1a** as 112 °C (0.5 Torr). However, we prepared **1a** according to the present method as well as to this literature's method, distilled it several times, and confirmed that the present bp value is correct.
- 4) J.-N. Collard and C. Benezra, *Tetrahedron Lett.*, **23**, 3725 (1982).
- 5) K. Issleib and H.-R. Roloff, *Chem. Ber.*, **98**, 2091 (1965); S. Antczak and S. Trippett, *J. Chem. Soc., Perkin Trans. 1*, **1978**, 1326; F. Mathey and F. Mercier, *J. Chem. Soc., Chem. Commun.*, **1980**, 191.
- 6) F. Mathey, M.-B. Comarmond, and D. Moras, *J. Chem. Soc., Chem. Commun.*, **1979**, 417.
- 7) S. Kobayashi, M. Suzuki, and T. Saegusa, *Macromolecules*, in press.
- 8) B. Sjöberg, *Chem. Ber.*, **74**, 64 (1941).
- 9) J. H. Chapman and L. N. Owen, *J. Chem. Soc.*, **1950**, 579.