

The Reaction of Cyclopropenethione with Carbonyl-stabilized Phosphonium Methylides to Yield 2*H*-Pyran-2-thione

Hiroshi YOSHIDA,* Jyun SHIMIZU, Tsuyoshi OGATA, and Kiyoshi MATSUMOTO†

Department of Applied Chemistry, Faculty of Engineering, Shizuoka University, Hamamatsu 432

†College of Liberal Arts and Sciences, Kyoto University, Kyoto 606


(Received April 10, 1985)

Synopsis. The reaction of diphenylcyclopropenethione with carbonyl-stabilized triphenylphosphonium methylides (**2**) to yield 2*H*-pyran-2-thiones was studied. It was clarified that electron-releasing substituents increased the reactivity of **2**.

The reactions of carbonyl-stabilized pyridinium and sulfonium methylides with cyclopropenone or cyclopropenethione to yield 2-pyrone derivatives^{1,2} are well known. On the other hand the reaction of diphenylcyclopropenone with acyl- or alkoxycarbonyl-stabilized phosphonium methylides yields either 2-pyrone or methylenecyclopropene *via* (3+3)-cycloaddition or Wittig reaction.^{1,2} However to our knowledge none of the reaction of diphenylcyclopropenethione (**1**) with carbonyl-stabilized phosphonium methylides (**2**) has been investigated.

In the continuation of our studies on the reaction of cyclopropenone with S-, N-, and P-ylides and imides,³ we extended our attention to the use of carbonyl-stabilized phosphonium methylides (**2**) which are stable to air and moisture.

Methylides **2** were prepared according to the known methods.⁴ Since **1** dimerizes on irradiation,⁵ it was treated in the dark. Although a benzene solution of **1** and 1-(triphenylphosphoranylidene)-2-propanone **2a** at reflux gave only a tarry mixture, the reaction at room temperature for 3 d yielded a brown crystalline mass (51%) together with triphenylphosphine in 84% yield after TLC purification. The structure of the product was confirmed to be 3,4-diphenyl-6-methyl-2*H*-pyran-2-thione **3a** on the basis of IR ($\nu_{C=S}$ 1130 cm⁻¹) and NMR spectroscopic studies. The ¹³C-NMR spectrum of **3a** gave a doublet at $\delta=111.1$ (C⁵, $J_{C^5H^5}=170$ Hz) and a quartet at $\delta=20.2$ (C⁷, $J_{C^7H^7}=140$ Hz) with long range couplings $J_{C^5H^5}=3$, $J_{C^5H^7}=4$, and $J_{C^6H^5}$ and $H^7=6$ Hz), indicating the presence of the framework:




A similar treatment of the carbonyl-stabilized methylides **2b—d** with **1** afforded 2*H*-pyran-2-thiones **3b—d**. The methylides substituted with aroyl groups **2e—i** were less reactive toward **1** than **2a—d**. Thus, the ylides **2e—i** reacted with **1** at toluene reflux to give **3e—i** in moderate yields (Table 1 and 2). The known 2*H*-pyran-2-thione **3f** was obtained in a higher yield than the previously reported (71%).² Physical properties are collected in Table 2.

As is evident from the Table 1, increasing the electron

releasing power of the substituent R increases the reactivity of **2**. The result correlates well with the reported mechanism for the reaction of cyclopropenone with S- and P-ylides.¹⁾ Thus, the formation of 2*H*-pyran-2-thiones can be explained either by an initial attack on the thiocarbonyl group of **1** by the betaine oxygen atom of **2** or a Michael type addition of the methine of **2** to the C=C bond of **1** followed by isomerization to yield the product.

In conclusion, phosphonium methylides **2** have proven to be convenient reagents for the preparation of 2*H*-pyran-2-thiones in moderate yields.



Scheme 1.

Experimental

General. Melting points are uncorrected. ¹H-NMR spectra were recorded on a Hitachi R-24B (60 MHz) spectrometer using TMS as an internal standard and ¹³C-NMR spectra on a JEOL JNM FX-60 spectrometer (15.04 MHz).

Materials. Diphenylcyclopropenethione **1** was prepared according to the literature.⁶ Phosphonium methylides

TABLE 1. THE REACTION OF **1** WITH **2**.

2	Reaction Temp/°C	Conditions Time	Yield 3 /%
2a	25	3d	51
2b	25	4d	39
2c	25	1d	72
2d	25	1d	46
2e	110	6h	74
2f	110	8h	87
2g	110	4d	70
2h	110	6d	39
2i	110	6d	37

TABLE 2. PHYSICAL PROPERTIES OF 3.

3	Mp	¹ H-NMR(CDCl ₃)	MS(M ⁺)	Found/% (Calcd/%)		
	$\theta_m/^{\circ}\text{C}$			C	H	N
3a	135—137	2.43(3H, s, Me), 6.43(1H, s, C ₆ H), 6.8—7.6(10H, m, Ar)	278	77.92 (77.66)	4.87 5.06	)
3b	148	6.57(1H, d, <i>J</i> =5Hz, C ₅ H), 6.8—7.4(10H, m, Ar), 7.78(1H, d, C ₆ H)	264	77.05 (77.24)	4.47 4.57	
3c	151—152	3.05(6H, s, 2Me), 5.44(1H, s, C ₆ H), 6.6—7.2(10H, m, Ar)	307	74.42 (74.24)	5.51 5.57	4.37 (4.56)
3d	96—103	2.70(3H, s, Me), 6.70(1H, s, C ₆ H), 7.0—7.9(10H, m, Ar)	310	69.71 (69.64)	4.65 4.54	4.94 (5.15)
3e	194—196	2.28(3H, s, Me), 6.7—7.9(14H, m, Ar)	354	81.16 (81.32)	5.32 5.11	)
3f	169—170 ^{a)}	6.9—7.8(m, Ar)	340	65.71 (65.87)	3.92 3.60	
3g	210—211	6.6—7.9(m, Ar)	419		)	
3h	213—214	6.9—8.1(m, Ar)	385	71.73 (71.67)	3.84 3.92	3.52 (3.63)
3i	194—196	6.5—8.4(m, Ar)	385	71.85 (71.67)	3.74 3.92	3.90 (3.63)

a) Ref. mp 165—166°C.²⁾2 were obtained by the known methods.⁴⁾

Reaction of 1 with 2. An equimolar mixture of **1** and **2** (**2a—d** in benzene at room temperature and **2e—i** in toluene at reflux temperature) was allowed to react in the dark. The disappearance of **1** was checked by TLC at suitable time intervals. Then, the solvent was removed *in vacuo* to give a crystalline mass. TLC separation over silica gel afforded **3** and triphenylphosphine. Yields and physical properties of **3a—i** are summarized in Tables 1 and 2.

¹³C-NMR spectroscopic data. **3a**(CDCl₃) $\delta=20.2$ (q, Me), 111.1 (d, C⁵), 127.5 (d), 127.9 (d), 128.0 (d), 128.4 (d), 128.7 (d), 130.6 (d), 136.1 (s), 136.8 (s), 137.0 (s), 147.9 (s), 165.8 (s, C⁶), and 198.0 (s, C²). **3f**(CDCl₃) $\delta=108.5$ (d, C⁵), 125.7 (d), 127.5 (d), 127.9 (d), 128.1 (s), 128.4 (d), 128.7 (d), 129.0 (d), 130.4 (s), 130.6 (d), 131.3 (d), 136.8 (s), 137.0 (s), 137.4 (s), 147.7 (s), 164.4 (s, C⁶), and 196.9 (s, C²).

References

- 1) K. T. Potts and J. S. Baum, *Chem. Rev.*, **74**, 189 (1974);

K. Matsumoto and Y. Ikemi, *Heterocycles*, **14**, 1445 (1980) and references cited therein.

2) T. Eicher, E. Angerer, and A. Hansen, *Justus Liebigs Ann. Chem.*, **746**, 102 (1971); T. Eicher, U. Sebold, and U. Stegmann, *Tetrahedron Lett.*, **24**, 2977 (1983) and references cited therein.

3) H. Yoshida, S. Sogame, Y. Takishita, and T. Ogata, *Bull. Chem. Soc. Jpn.*, **56**, 2438 (1983).

4) H. J. Bestmann and R. Zimmermann, "Organic Phosphorus Compounds," ed by G. M. Kosolapoff and L. Maier, Wiley-Interscience, New York (1972) Vol. 3.

5) A. Schoenberg and M. Mamluk, *Tetrahedron Lett.*, **1971**, 4993.

6) H. Yoshida, M. Nakajima, and T. Ogata, *Synthesis*, **1981**, 36.