

Favorskii-Type Rearrangement of α,α' -Dihalo Ketones Induced by Enolates of (Diethoxyphosphinyl)acetic Esters and Its Application to the Synthesis of α,α' -Divinyl Ketones


Takashi SAKAI,* Eiichiro AMANO, Kazuyoshi MIYATA,
Masanori UTAKA, and Akira TAKEDA

Department of Synthetic Chemistry, School of Engineering, Okayama University,
Tsushima, Okayama 700
(Received October 16, 1986)

Synopsis. A Favorskii-type rearrangement of 1,3-dihalo-3-methyl-2-butanone with enolate of ethyl (diethoxyphosphinyl)acetate or *t*-butyl (diethoxyphosphinyl)acetate gave ethyl 2-(diethoxyphosphinyl)-5-methyl-3-oxo-4-hexenoate (**3a**) or its *t*-butyl ester (**3b**). Compound **3b** was applied to the preparation of (*E*)-5-methyl-1-(*p*-tolyl)-1,4-hexadien-3-one and diethyl (4-methyl-2-oxo-3-pentenyl)phosphonate, which are the important intermediates for natural product syntheses.

Recently, we have made considerable efforts^{1a-d} to extend the base of the Favorskii rearrangement² to carbanions such as enolates of malonic ester and (phenylsulfonyl)acetic ester. In a preceding paper,^{1d} we reported that sodium enolates of β -keto esters reacted with 1,3-dihalo-3-methyl-2-butanone to give dihydro-4-pyrone derivatives via the Favorskii-type rearrangement. In this paper, we elucidate that the reaction is adaptable to enolates of (diethoxyphosphinyl)acetic esters, giving α -(diethoxyphosphinyl) β -keto esters (**3a,b**) in good yields. Products having a phosphinyl-functionality were utilized for the preparation of α,α' -divinyl ketones.

A reaction of 1,3-dichloro-3-methyl-2-butanone (**1a**) with sodium enolate of ethyl (diethoxyphosphinyl)acetate (**2a**) in THF at room temperature gave ethyl 2-(diethoxyphosphinyl)-5-methyl-3-oxo-4-hexenoate (**3a**) in 42% yield (Scheme 1). The yield was improved up to 79% by using 3-bromo-1-chloro-3-methyl-2-butanone (**1b**). The *t*-butyl ester **3b** was similarly obtained in 36% yield from **1a** or in 81% yield from **1b**. The reaction can be explained in parallel with that described previously.^{1d}




Scheme 1.


Attempts to apply the above reaction to 3-bromo-3-methyl-2-butanone (**4**) and 3,4-dibromo-4-methyl-2-pentanone (**6**) resulted in giving **5** (43% yield) and **8** (41% yield), respectively (Scheme 2). The formation of **8** suggests to us that the enolate **2a** may attack C-3 methylene carbon, instead of the carbonyl carbon, of

the cyclopropanone intermediate **7**, which is produced under the basic conditions in a manner similar to the conventional Favorskii rearrangement.² It is interesting that the mode of the ring cleavage³ of **7** is in marked contrast to that of the Favorskii rearrangement although a further investigation is required to confirm the mechanism.

The Horner–Wadsworth–Emmons reaction of sodium



Scheme 2.



Scheme 3.

enolate derived from **3a** with benzaldehyde (**9a**) was carried out in refluxing 1,2-dimethoxyethane (DME), giving α -benzylidenation product **10a** in 31% yield⁴ (Scheme 3). A similar reaction with *p*-tolualdehyde (**9b**) gave **10b** in only 17% yield.⁴ On the other hand, sodium enolate derived from **3b** was allowed to react with aldehyde **9b** at an elevated temperature (130 °C) in diglyme to give (*E*)-5-methyl-1-(*p*-tolyl)-1,4-hexadien-3-one (**12**) in 47% yield. In this case, deesterification of **11** may have occurred thermally⁵ in the course of the reaction. Furthermore, the treatment of **3b** with a catalytic amount of *p*-toluenesulfonic acid in refluxing toluene gave (4-methyl-2-oxo-3-pentenyl)-phosphonate (**13**) in 78% yield. Compounds **12** and **13** are the important intermediates for the syntheses of *ar*-turmerone^{1d,6} and related sesquiterpenes.⁷

Experimental

IR spectra were taken on a JASCO Model A-102 spectrometer. ¹H NMR spectra (60 MHz) were measured with a JEOL Model JNM-60 SI spectrometer by using Me₄Si as an internal standard. Preparative TLC was carried out on silica gel (Kieselgel 60 PF₂₅₄, Merck A. G. Darmstadt).

Materials: 1,3-Dichloro-3-methyl-2-butanone (**1a**)⁸ was prepared by chlorination of 3-methyl-2-butanone with 2.4 equiv of sulfur chloride. 3-Bromo-3-methyl-2-butanone (**4**)⁹ was prepared by bromination of the starting ketone with *N*-bromosuccinimide. 3-Bromo-1-chloro-3-methyl-2-butanone (**1b**)¹⁰ was obtained by chlorination of **4** with sulfur chloride. (Diethoxyphosphinyl)acetates **2a**¹¹ and **2b**¹² were prepared by the reported procedures.

Ethyl 2-(Diethoxyphosphinyl)-5-methyl-3-oxo-4-hexenoate (3a). To a suspension of NaH (48 mg, 2 mmol) in THF (6 ml) was added a solution of ethyl (diethoxyphosphinyl)acetate (448 mg, 2 mmol) in THF (2 ml); stirring was continued at room temperature for 2 h. To this was added **1b** (199 mg, 1 mmol) with stirring during an additional 48 h at room temperature. The mixture was poured into water and acidified with 10% HCl. The organic layer was extracted with ether, washed with aqueous NaHCO₃ and brine, dried (MgSO₄), and concentrated. The residue was purified by preparative TLC on silica gel (hexane–acetone, 3:1) to give 242 mg (79%) of **3a**: IR (neat) 1740, 1693, 1620, and 1555 cm⁻¹; ¹H NMR (CCl₄) δ =1.32 (9H, t, *J*=7 Hz), 1.96 (3H, d, *J*=1.5 Hz), 2.14 (3H, d, *J*=1.2 Hz), 3.74–4.36 (6.77H, m), 6.32 (0.77H, m), 6.79 (0.23H, m), 14.65 (0.23H, br s, enol OH). Found: C, 50.87; H, 7.64%. Calcd for C₁₃H₂₃O₆P: C, 50.98; H, 7.57%.

A similar reaction of **1a** with **2a** gave **3a** in 42% yield.

***t*-Butyl 2-(Diethoxyphosphinyl)-5-methyl-3-oxo-4-hexenoate (3b).** A similar reaction of **1a** and **1b** with sodium enolate of *t*-butyl (diethoxyphosphinyl)acetate (**2b**) gave **3b** in 36% and 81% yields, respectively. **3b**: IR (neat) 1730, 1690, 1620, and 1555 cm⁻¹; ¹H NMR (CCl₄) δ =1.31 (6H, t, *J*=7 Hz), 1.45 (9H, s), 1.93 (3H, br s), 2.09 (3H, br s), 3.67–4.36 (4.8H, m), 6.30 (0.2H, m), 6.73 (0.8H, m), 13.90 (0.2H, br s, enol OH). Found: C, 53.97; H, 8.06%. Calcd for C₁₅H₂₇O₆P: C, 53.89; H, 8.18%.

Ethyl (*E*)-4-Bromo-3,4-dimethyl-2-pentenoate (5):¹³ 43% yield after purification by column chromatography (hexane–ether, 30:1); IR (neat) 1725 and 1640 cm⁻¹, ¹H NMR (CCl₄) δ =1.28 (3H, t, *J*=7 Hz), 1.93 (6H, s), 2.36 (3H, s), 4.11 (2H, q, *J*=7 Hz), 5.85 (1H, m).

Ethyl 2-(Diethoxyphosphinyl)-6-methyl-4-oxo-5-heptenoate (8). To a mixture of mesityl oxide (98.1 mg,

1.0 mmol), NaHCO₃ (35 mg, 0.42 mmol), and ether (1 ml), was added bromine (159 mg, 1.0 mmol) dropwise; the mixture was stirred for 2 h. The resulting ether solution of 3,4-dibromo-4-methyl-2-pentanone (**6**)¹⁴ was slowly introduced to a suspension of **2a** which was prepared from ethyl (diethoxyphosphinyl)acetate (448 mg, 2.0 mmol) and NaH (48 mg, 2 mmol) in THF (2 ml) in a manner similar to that described above. After being stirred for 21 h at room temperature, the usual work up of the mixture gave 128 mg (41%) of **8** after purification by preparative TLC (hexane–acetone, 4:1; twice development, *R*_f=0.16–0.26). **8**: IR (neat) 1740, 1695, and 1625 cm⁻¹; ¹H NMR (CCl₄) δ =1.28, 1.30, 1.32 (each with 3H, t, *J*=7 Hz), 1.88 (3H, br s), 2.10 (3H, br s), 2.7–3.7 (3H, m), 4.13 (6H, q, *J*=7 Hz), 6.02 (1H, m). Found: C, 52.43; H, 7.93%. Calcd for C₁₄H₂₅O₆P: C, 52.50; H, 7.87%.

Ethyl 2-Benzylidene-5-methyl-3-oxo-4-hexenoate (10a).

To a suspension of NaH (72 mg, 3 mmol) in DME (3 ml) was added a solution of ester **3a** (918 mg, 3 mmol) and benzaldehyde (**9a**) (501 mg, 3.6 mmol) in DME (4 ml). The mixture was stirred for 1 h at room temperature and then for an additional 3 h under reflux. It was poured into water and acidified with 10% HCl. The organic layer was extracted with ether, washed with brine, dried (MgSO₄), and concentrated. The residual oil was purified by preparative TLC (hexane–acetone, 3:1) to give 240 mg (31%) of **10a**: IR (neat) 1720, 1705, 1672, and 1618 cm⁻¹; ¹H NMR (CCl₄) δ =1.30 (3H, t, *J*=7 Hz), 1.84 (2.2H, br s), 1.95 (0.8H, br s), 2.17 (3H, br s), 4.20 (2H, q, *J*=7 Hz), 5.92 (0.7H, m), 6.20 (0.3H, m), 7.26 (5H, m), 7.40 (0.3H, s), 7.47 (0.7H, s). Found: C, 74.46; H, 6.94%. Calcd for C₁₆H₁₈O₃: C, 74.40; H, 7.02%.

Ethyl 5-Methyl-3-oxo-2-(*p*-tolylmethylene)-4-hexenoate (10b). A similar reaction of **3a** with *p*-tolualdehyde (**9b**) gave **10b** (17%): IR (neat) 1719, 1670, and 1610 cm⁻¹; ¹H NMR (CDCl₃) δ =1.27 (3H, t, *J*=7 Hz), 1.83 (1.7H, s), 1.94 (1.3H, s), 2.18 (3H, br s), 2.33 (3H, s), 4.21 (1.1H, q, *J*=7 Hz), 4.28 (0.9H, q, *J*=7 Hz), 6.02 (0.56H, m), 6.27 (0.44H, m), 6.9–7.4 (4H, m), 7.43 (0.44H, s), 7.51 (0.56H, s). Found: C, 75.16; H, 7.34%. Calcd for C₁₇H₂₀O₃: C, 74.97; H, 7.40%.

(*E*)-5-Methyl-1-(*p*-tolyl)-1,4-hexadien-3-one (12). A similar reaction of enolate of **3b** with **9b** was carried out in diglyme at 130 °C for 16 h. The crude product was purified by preparative TLC (hexane–acetone, 3:1) to afford **12** (47%), whose ¹H NMR spectral data were identical with those reported.⁶

Diethyl (4-Methyl-2-oxo-3-pentenyl)phosphonate (13).

A mixture of **3b** (131 mg, 0.39 mmol), *p*-toluenesulfonic acid (40 mg, 0.23 mmol), and toluene (3 ml) was stirred for 3.5 h under reflux. The cooled mixture was washed with aqueous NaHCO₃ and water. The organic layer was separated and the aqueous layer was extracted with ether. The combined organic layer was dried (MgSO₄) and then concentrated. Distillation of the crude product [bp 100–105 °C (0.3 mm Hg) (1 mmHg=133.322 Pa)] gave **13** (89 mg, 78%): IR (neat) 1680, 1620, and 1255 cm⁻¹; ¹H NMR (CCl₄) δ =1.33 (6H, t, *J*=7 Hz), 1.93 (3H, br s), 2.13 (3H, br s), 2.88 (2H, d, *J*=22 Hz), 3.97 and 4.11 (each with 2H, q, *J*=7 Hz), 6.20 (1H, m). Found: C, 59.25; H, 9.53%. Calcd for C₁₀H₁₉O₂P: C, 59.39; H, 9.47%. These spectral data are consistent with those of the corresponding dimethyl ester.^{6a}

References

- 1) a) T. Sakai, E. Amano, A. Kawabata, and A. Takeda, *J. Org. Chem.*, **45**, 43 (1980); b) T. Sakai, T. Katayama, and A. Takeda, *J. Org. Chem.*, **46**, 2924 (1981); c) T. Sakai, H. Tabata, and A. Takeda, *J. Org. Chem.*, **48**, 4618 (1983); d) T. Sakai, K. Miyata, M. Ishikawa, and A. Takeda, *Tetrahedron*

Lett., **26**, 4727 (1985).

2) A. S. Kende, "Organic Reactions," John Wiley & Sons, New York (1960), Vol. 11, Chap. 4.

3) Synthesis of α,β -unsaturated aldehyde by the hydrolysis of 1-methoxy-(1-mesyloxyalkyl)cyclopropane through a similar type of the ring-cleavage: E. J. Corey and P. Ulrich, *Tetrahedron Lett.*, **1975**, 3685.

4) Products other than **10a,b** were obtained as a complex mixture. Their purification and structural identification were unsuccessful. The starting esters **3a,b** were not recovered.

5) Formation of radical species in the pyrolysis of β -keto esters was reported: M. B. S. Kirkiacharian, *C. R. Acad. Sci. Ser. C.*, **269**, 721 (1969).

6) a) Preparation of **13** by the reaction of 3-methyl-2-pentenoyl chloride with lithium (dimethoxyphosphinyl)-methanide and its application to the synthesis of **12**: J. Motoyoshiya, M. Miyajima, K. Hirakawa, and T. Kakurai, *J. Org. Chem.*, **50**, 1326 (1985); b) Preparation of **12** by the reaction of **9b** with lithium enolate of mesityl oxide and

subsequent dehydration: O. S. Park, Y. Grillasca, G. A. Garcia, and L. A. Maldonado, *Synth. Commun.*, **7**, 345 (1977), and references cited in these literatures.

7) L. A. Paquette, W. E. Fristad, D. S. Dime, and T. R. Bailey, *J. Org. Chem.*, **45**, 3017 (1980).

8) N. Schamp, N. De. Kimpe, and W. Coppens, *Tetrahedron*, **31**, 2081 (1975).

9) J. W. Thorpe and J. Warkentin, *Can. J. Chem.*, **51**, 927 (1973).

10) A. A. Petrov, *J. Gen. Chem.*, **15**, 931 (1945).

11) H. Takahashi, K. Fujiwara, and M. Ohta, *Bull. Chem. Soc. Jpn.*, **35**, 1498 (1962).

12) G. F. Griffiths, G. W. Kenner, S. W. McCombie, K. M. Smith, and N. J. Sutton, *Tetrahedron*, **32**, 275 (1976).

13) C. E. Moppett and J. K. Sutherland, *J. Chem. Soc.*, **1968**, 3040.

14) a) H. Rupe and S. Kessler, *Chem. Ber.*, **42**, 4715 (1909); b) K. S. Ayyar, R. C. Cookson, and D. A. Kagi, *J. Chem. Soc., Perkin Trans. 1*, **1975**, 1727.