

Silaethene in Pyrolysis of Spiro[silacyclobutane-1,9'-[9H-9]silafluorene]

Daiyo TERUNUMA,* Makoto NAKAMURA, Eiji MIYAZAWA, and Hiroyuki NOHIRA

Department of Applied Chemistry, Faculty of Engineering, Saitama University, Urawa, Saitama 338

(Received January 26, 1987)


Synopsis. A new silaspiro compound, i.e., spiro[silacyclobutane-1,9'-[9H-9]silafluorene](1) was synthesized in 75% yield. It was found that the pyrolysis of 1 at 650 °C under a reduced pressure gave a head-to-tail silaethene dimer in 15% yield. Copyrolysis of 1 in the presence of benzophenone gave 1,1-diphenylethene and a siloxane oligomer in 50 and 30% yield, respectively.

Current attention has been concentrated to the preparation and characterization of a silicon-carbon double bond.¹⁾ It is suggested that 1-silafulvene systems have strong π stability and high reactivity, on the contrary 2-silafluorene systems have weak π stability and low reactivity.²⁾ In this connection, the generation of 1-silafluorene was already investigated.³⁾

In the course of our studies on the synthesis of silicon-containing cyclic compounds,⁴⁾ we have interested in spiro[silacyclobutane-1,9'-[9H-9]silafluorene] (1). Because the generation of 2-silafluorene type intermediate would be expected with extrusion of ethene in the pyrolysis of 1.


Results and Discussion

Compound 1 was obtained by treating 1,1-dichlorosilacyclobutane with biphenyl-2,2'-diyl dilithium in 75% yield (Scheme 1).



Scheme 1.


The mass spectrum of 1 shows relatively simple fragmentation including the base peak (m/z 194) corresponded to the silaethene fragment (Scheme 2).



Scheme 2.

Then the pyrolysis of 1 was carried out at 650 °C under a reduced pressure (ca. 10^{-3} Torr[†]). A white crystalline product was obtained. The parent peak of the mass spectrum of the product was consistent with the molecular weight of a dimer of 2-silafluorene derivative (2) (m/z 388). Thus it is clear that 2 is generated under these reaction conditions and successive reaction of 2 occurs to give the dimer (Scheme 3).


† 1 Torr = 133.322 Pa.



Scheme 3.

On the other hand, the silaethene can be dimerized in the two manners,⁵⁾ head-to-head (A) and head-to-tail (B).

NMR spectrum (²⁹Si) of the dimer (δ +2.1 based on TMS) suggests 1,3-disilacyclobutane structure (B).⁶⁾ The disilacyclobutane derivative has been known to undergo ring-opening reaction in the presence of lithium aluminum hydride (LAH).⁷⁾ The dimer was treated with LAH. The expected product from A and B are shown in Scheme 4.




Scheme 4.

NMR spectra data of the product [¹³C NMR (CDCl₃) δ -5.20 (t, CH₂), -2.27 (q, CH₃)] shows (c) in Scheme 4 which indicates the dimer structure of (B).

Furthermore, the reaction of the dimer with dimethylphenylsilane in the presence of platinum/carbon⁸⁾ afforded corresponding ring-opened product (d) (Scheme 5), but no reaction occurred with *m*-chloroperbenzoic acid (MCPBA).


These results show that the dimer has the structure of type B.



Scheme 5.

Then copyrolysis of 1 with benzophenone was investigated. The reaction was carried out at 540 °C under a

reduced pressure (ca. 10^{-3} Torr). 1,1-Diphenylethene and a siloxane oligomer were obtained in a 50 and 30% yield, respectively. Although we could not find out the direct evidence for the existence of the intermediates, it seems to be reasonable to consider that the reaction proceeds via a manner similar to that described in the copolyrlysis of the silacyclobutane derivatives with benzophenone⁹ (Scheme 6).



Scheme 6.

Experimental

The melting points and boiling points are uncorrected. The IR spectra were determined with a JASCO IR-A 302 spectrometer. The NMR spectra were determined at 90 MHz with a JEOL FX90Q and at 200 MHz with a JEOL FX200 spectrometers in CDCl₃ using TMS as an internal standard. The mass spectra were recorded on a JEOL-01SG instrument. Biphenyl-2,2'-diyl dilithium was prepared in a manner similar to that described in the literature.¹⁰

Spiro[silacyclobutane-1,9'-[9H-9]silafuorene] (1). In a flask lithium (2.0 g, 0.29 gram-atoms) and dry ether (30 ml) were placed. A solution of 2,2'-diiodobiphenyl (12.0 g, 29.6 mmol) in dry ether (50 ml) was added to the mixture at reflux temperature with stirring under a dry nitrogen atmosphere. The mixture was refluxed for 1 h after the addition was complete. In another flask, a solution of 1,1-dichlorosilacyclobutane (4.0 g, 28.4 mmol) in dry ether (40 ml) was placed. The ethereal solution of biphenyl-2,2'-diyl dilithium was added to the solution of the chlorosilane by means of a syringe at 0°C. The reaction mixture was stirred at 0°C for 1 h, allowed to stand at room temperature for 1 h. The reaction mixture was hydrolyzed and the organic layer was separated and dried (Na₂SO₄). Evaporation and distillation gave **1** in 4.4 g (75%) yield. Bp 131°C/0.25 Torr. n_D^{25} 1.6511. IR (neat) 2900, 1600, 1430, 1350, 1260, 1130, 1060, 850, 740, 720, 680, 540 cm⁻¹. ¹H NMR (CDCl₃) δ =1.56 (t, 4, Si-CH₂), 2.38 (quintet, CH₂), 7.1–7.8 (m, 8, Ar). ¹³C NMR (CDCl₃) δ =15.0 (Si-CH₂), 19.29 (CH₂), 120.55, 127.48, 130.68, 133.11, 135.61, 147.58 (Ar). ²⁹Si NMR (CDCl₃) δ =12.9. UV (cyclohexane) $\epsilon_{\max}$ (nm) 237 (4.51×10⁴), 244 (3.84×10⁴), 281 (1.40×10⁴), 293 (1.21×10⁴). Found: C, 81.00; H, 6.51%. Calcd for C₁₅H₁₄Si: C, 81.02; H, 6.34%.

Pyrolysis of 1. Pyrolysis was carried out using a vacuum flow system. Compound **1** (527 mg, 2.4 mmol) was added into a quartz tube (l=40 cm, d=0.8 cm) maintained at 640°C through preheated tube (ca. 300°C) under a reduced pressure (4×10⁻³ Torr). The products were collected in a liq. N₂ trap, and washed out with benzene. Evaporation and filtration gave crystalline product. It was purified by recrystallization from benzene. Yield 70 mg (15%). Mp 323–326°C. MS M⁺ 388. IR (KBr) 3050, 1600, 1435, 1340, 1260, 1130, 1080, 940, 770, 730, 710, 630 cm⁻¹. ¹H NMR (CDCl₃) δ =1.17 (s, 4H, Si-CH₂), 7.2–8.0 (m, 16H, Ar). ¹³C NMR (CDCl₃) δ =1.10 (Si-CH₂), 121.1, 128.0, 131.0, 132.8, 137.5, 147.5 (Ar). ²⁹Si NMR (CDCl₃) δ =+2.1. UV (cyclohexane) $\epsilon_{\max}$ (nm) 238 (9.65×10⁴), 245 (1.02×10⁵), 283 (3.44×10⁴).

Copolyrlysis of **1** (500 mg, 2.25 mmol) in the presence of benzophenone (410 mg, 2.25 mmol) was carried out at 540°C in a manner similar to that described above. 1,1-

Diphenylethene and the siloxane oligomer were separated from the reaction mixture by TLC (benzene-hexane 4:1) in 50% (200 mg) and 30% (130 mg) yield, respectively. 1,1-Diphenylethene; ¹H NMR (CCl₄) δ =5.35 (s, 2H, =CH₂), 7.20 (s, 10H, Ph). Oligomer IR (KBr) 3000, 1590, 1420, 1260, 1120, 1060, 740, 500 cm⁻¹. ¹H NMR (CCl₄) δ =7.0–8.2 (Ar).

Reaction of the Dimer with LAH. LAH (50 mg, 1.3 mmol) was added to a solution of the dimer (200 mg, 0.5 mmol) in dry ether (30 ml). The mixture was refluxed for 1 h with stirring. Usual work-up and purification by TLC (benzene) gave ring-opened product in 120 mg (60%) yield. ¹H NMR (CDCl₃) δ =0.40 (s, H, Si-CH₃), 0.61 (d, 2H, Si-CH₂-Si), 4.87 (t, 1H, Si-H), 7.1–7.8 (m, 16H, Ar). ¹³C NMR (CDCl₃) δ =-5.20 (t, CH₂), -2.27 (q, Si-CH₃), 120.77, 127.33, 130.20, 130.33, 132.99, 133.64, 135.46, 138.19, 147.69, 148.34 (Ar).

Reaction of the Dimer with Dimethylphenylsilane in the Presence of Pt/C Catalyst. A mixture of the dimer (39.8 mg, 0.10 mmol), dimethylphenylsilane (1.00 g, 7.4 mmol), and 5% Pt/C (10 mg) was placed in a test tube. The tube was sealed under a dry nitrogen atmosphere, and it was kept at 100°C for 55 h. After evaporation of excess dimethylphenylsilane, the product was separated by TLC (hexane-benzene 3:1) and recrystallized from pentane. Yield 16.1 mg (30%). ¹H NMR (CDCl₃) δ =-0.06 (s, 6H, Si-CH₃), 0.44 (s, 2H, Si-CH₂-Si), 0.58 (d, 2H, Si-CH₂), 4.82 (t, 1H, Si-H), 7.1–7.8 (m, 21H, Ar).

The authors are grateful to Professor Yasuhiro Nakadaira for help and advice in the determination of the structure of the dimer, and Associate Professor Sumio Tokita for letting us know the improved preparation method of 2,2'-diiodobiphenyl.

The authors also wish to thank the Shin-etsu Chemical Industry Co. for the gift of silanes.

References

- 1) H. Sakurai, *Kagaku No Ryoiki*, **29**, 742 (1975); L. E. Gusel'nikov and N. S. Nametkin, *Chem. Rev.*, **79**, 529 (1979); A. G. Brook, *J. Organomet. Chem.*, **300**, 21 (1986).
- 2) G. Bertrand, G. Trinquier, and P. Mazerolles, "Organometallic Chemistry Reviews" ed by D. Seyferth, Elsevier, Amsterdam (1981) Vol. 12, 1.
- 3) Y. Nakadaira, H. Sakaba, and H. Sakurai, *Chem. Lett.*, **1980**, 1071; T. J. Barton, G. T. Burns, E. V. Arnold, and J. Clardy, *Tetrahedron Lett.*, **7** (1981); T. J. Barton and C. R. Tully, *J. Organomet. Chem.*, **172**, 11 (1979).
- 4) D. Terunuma, N. Shibuya, and H. Nohira, *Bull. Chem. Soc. Jpn.*, **55**, 2287 (1982).
- 5) A. G. Brook, J. W. Harris, I. Lennon, M. EISheikh, *J. Am. Chem. Soc.*, **101**, 83 (1979); M. Ishikawa, T. Fuchikami, M. Kumada, T. Higuchi, and S. Miyamoto, *J. Am. Chem. Soc.*, **101**, 1348 (1979); N. S. Nametkin, L. E. Gusel'nikov, V. M. Vdovin, P. L. Grinberg, V. I. Zav'yakov, and V. D. Oppengein, *Dokl. Akad. Nauk SSSR*, **171**, 630 (1966).
- 6) A. M. Krapivin, M. Magi, V. I. Svergun, R. Z. Zaharjan, E. D. Babich, and N. V. Ushakov, *J. Organomet. Chem.*, **190**, 9 (1980).
- 7) A. M. Devine, P. A. Griffin, R. N. Haszeldine, M. J. Newlands, and A. E. Tipping, *J. Chem. Soc., Dalton Trans.*, **1975**, 1434.
- 8) D. R. Weyenberg and L. E. Nelson, *J. Org. Chem.*, **30**, 2618 (1965).
- 9) D. N. Roark and L. H. Sommer, *Chem. Commun.*, **1973**, 167; C. M. Golino, R. D. Bush, D. N. Roark, and L. H. Sommer, *J. Organomet. Chem.*, **66**, 29 (1974).
- 10) G. Wittig and W. Herwig, *Chem. Ber.*, **87**, 1511 (1954).