

The Magnetic Field Effects on Electrolysis. III.¹⁾ The Anodic Oxidation of Phenylacetate Ion

Takeshi WATANABE,[†] Yoshifumi TANIMOTO,^{*,††} Ryoichi NAKAGAKI, Mitsuo HIRAMATSU,^{†††}
and Saburo NAGAKURA

Institute for Molecular Science, Myodaiji, Okazaki 444

(Received April 2, 1987)

Synopsis. In anodic oxidation of the phenylacetate ion, the formation yield of benzaldehyde increases by about 40% in the presence of a magnetic field (0.67 T), while 1,2-diphenylethane is insensitive to the field. The results are interpreted in terms of the magnetohydrodynamic mechanism.

In previous studies,^{1,2)} we have investigated external magnetic field effects on inorganic electrochemical processes and demonstrated that the magnetohydrodynamic (MHD) effect plays an important role in redox reactions. In the MHD mechanism, a flow of electrons and ions is affected by an external magnetic field, resulting in the perturbation on the mass-transport process near the electrode. As an extension of the above-mentioned investigation, we have studied the magnetic field effects on organic electrochemical processes.

In the present paper, we report on the magnetic field effects on the electrolysis of the phenylacetate ion, as an example of the Kolbe reaction, which may be preparatively useful.

Experimental

Phenylacetic acid was prepared by the hydrolysis of ethyl phenylacetate,³⁾ and was purified by recrystallization. Solvents for high-performance liquid chromatography (HPLC) grade, methanol, dioxane and water, were used as received.

The electrolysis was carried out using a potentio/galvano stat (Hokuto Denko, HA-102). An aerated methanol–dioxane solution (50 ml), containing phenylacetic acid (0.05 mol dm⁻³) and its potassium salt (0.05 mol dm⁻³), was electrolyzed between two platinum plate electrodes (5 mm × 10 mm) at a constant density of 9 mA cm⁻² for about 90 min (0.1 F mol⁻¹). Argon or oxygen gas was bubbled through the solution during the electrolysis, when necessary. A magnetic field was applied with an electromagnet (Tokin SEE-10).

The reaction products were analyzed on an HPLC apparatus (Waters model 590 liquid chromatograph) and a UV detector (Waters 481 LC-spectrophotometer). The products were identified from the agreement of their UV absorption spectra and retention times with those of authentic samples, by monitoring the absorbance at 260 nm.

[†] Present address: Toray Research Center, INC, Sonoyama, Otsu 520.

^{††} Adjunct associate professor of Institute for Molecular Science (1983–85). Faculty of Pharmaceutical Sciences, Kanazawa University, Takara-machi, Kanazawa 920.

^{†††} Visiting research fellow from Hamamatsu Photonics K. K. (1983–85). Hamamatsu Photonics K. K., Ichino-cho, Hamamatsu 435.

Results and Discussion

Figure 1a shows a typical chromatogram of the reaction mixture obtained by electrolysis in the absence of a magnetic field. The five peaks in the chromatogram have been identified as benzaldehyde (PhCHO), benzyl methyl ether (PhCH₂OCH₃), toluene (PhCH₃), benzyl phenylacetate (PhCH₂COOCH₂Ph), and 1,2-diphenylethane (PhCH₂CH₂Ph), though the other two peaks with retention times of 10.2 and 16.8 min remain unidentified. Figure 1b shows the effects of a magnetic field (0.67 T) on the product yields. In the presence of a magnetic field, the yield of benzaldehyde increases by about 40%, while those of 1,2-diphenylethane and others are insensitive to the magnetic field (summarized in Table 1).

The electrolysis of substituted phenylacetate ions in methanol was studied in detail by Coleman et al.⁴⁾ According to their proposed mechanism, the plausible




Fig. 1. Typical chromatograms of the reaction mixture after the electrolysis of an aerated solution of phenylacetic acid and its potassium salt in the absence (a) and in the presence (b) of a magnetic field (0.67 T).

To our knowledge, this is the first report of the

magnetic field effect in organic electrochemistry, though a large number of papers have been concerned with inorganic electrochemistry.^{1,2,9-11} We believe that the present finding demonstrates the applicability of an external magnetic field effect to the mechanistic study of organic electrochemistry.

References

- 1) Part II of this series: T. Watanabe, Y. Tanimoto, R. Nakagaki, M. Hiramatsu, T. Sakata, and S. Nagakura, *Bull. Chem. Soc. Jpn.*, **60**, 4163 (1987).
 - 2) T. Watanabe, Y. Tanimoto, T. Sakata, R. Nakagaki, M. Hiramatsu, and S. Nagakura, *Bull. Chem. Soc. Jpn.*, **58**, 1251 (1985).
 - 3) R. Adams and A. F. Thal, *Org. Synth.*, Coll. Vol. 2, 63 (1922).
 - 4) J. P. Coleman, J. H. P. Utley, and B. C. L. Weedon, *J. Chem. Soc., Chem. Commun.*, **1971**, 438; J. P. Coleman, R. Lines, J. H. P. Utley, and B. C. L. Weedon, *J. Chem. Soc., Perkin Trans. 2*, **1974**, 1064.
 - 5) Y. Takegami, Y. Fujimura, H. Ishii, and T. Iwamoto, *Kogyo Kagaku Zasshi*, **68**, 196 (1965).
 - 6) For example, see: Y. Tanimoto, H. Hayashi, S. Nagakura, H. Sakuragi, and K. Tokumaru, *Chem. Phys. Lett.*, **41**, 267 (1976); Y. Tanimoto, T. Watanabe, R. Nakagaki, M. Hiramatsu, and S. Nagakura, *ibid.*, **116**, 341 (1985); R. Nakagaki, M. Hiramatsu, T. Watanabe, Y. Tanimoto, and S. Nagakura, *J. Phys. Chem.*, **89**, 3222 (1985); Y. Tanimoto, M. Takayama, M. Itoh, R. Nakagaki, and S. Nagakura, *Chem. Phys. Lett.*, **129**, 414 (1986); Y. Tanimoto, N. Okada, M. Itoh, K. Iwai, K. Sugioka, F. Takemura, R. Nakagaki, and S. Nagakura, *ibid.*, **136**, 42 (1987).
 - 7) F. J. J. de Kanter, J. A. den Hollander, A. H. Huizer, and R. Kaptein, *Mol. Phys.*, **34**, 857 (1985).
 - 8) G. L. Closs and O. D. Redwine, *J. Am. Chem. Soc.*, **107**, 6131 (1985).
 - 9) S. Mohanta and T. Z. Fahidy, *Can. J. Chem. Eng.*, **50**, 248 (1972).
 - 10) F. Takahashi, Y. Sakai, and T. Tamura, *Electrochim. Acta*, **28**, 1147 (1983).
 - 11) R. Aogaki and K. Fueki, *J. Electrochem. Soc.*, **131**, 1295 (1984).
-