




Since the formation of allyl alcohols on permanganate oxidation of alkenes is unfamiliar to our knowledge, further study of the structure of **2** and **3** was required. Grignard methylation of mayurone **7** gave **2** and **3** in a ratio of 1:9. Relative values of the lanthanoid-induced shift (LIS) using $\text{Eu}(\text{fod})_3$ and $\text{Pr}(\text{fod})_3$ are summarized in Table 1. The data show that the OH group of **3** is closer to the cyclopropane ring than that of **2**. Thus, the OH group in **3** must be in an α -configuration, and that in **2** in a β -configuration. Interestingly, all of the three tertiary methyl groups remote from the OH group in the epimers **2** and **3** showed the same chemical shifts. The difference in the LIS values of the bridge head proton, H-C(7), in **2** and **3** is small. These results show that the epimers have the same ring conformation in which the cyclohexene ring has a boat form with a flag pole of a C(1) atom. The LIS data observed for two vinyl protons and two methylene protons on the




Table 1. Lanthanoid-Induced Shift (LIS) of **2** and **3**

Proton	Chemical shift δ /ppm		Relative LIS	
	2	3	2	3
(Cyclopropane)				
H α on C(6)	0.02	0.22	0.449	0.754
H β on C(6)	0.44	0.39	0.321	0.465
H on C(7)	1.55	1.48	0.990	0.916
(Vinyl)				
H on C(9)	5.24	5.10	0.786	0.983
H on C(10)	5.15	4.92	0.368	0.340
(Methyl)				
Me on C(8)	1.28	1.40	1.00	1.00
Me	0.66	0.65	0.152	0.207
Me	1.07	1.06	0.143	0.156
Me	1.17	1.15	0.216	0.216

cyclopropane ring are consistent with this conformation.

Oxidation in Aqueous Acetone: Gas chromatographic analysis of the neutral product obtained under Kawamura's conditions showed two main peaks besides more than 12 minor peaks. Chromatographic separation with silica gel gave a mixture of *cis*-8,9-diols **10** and **11** as one of the main products. The mixture of **10** and **11** showed a single peak on gas chromatography. The ratio of **10** to **11** was about 1:4 by NMR analysis. Further chromatography of the mixture on neutral alumina gave 8 β ,9 β -diol **10**, mp 68–71 °C and 8 α ,9 α -diol **11**, mp 71–3 °C. Repeated recrystallization of a mixture of **10** and **11** from hexane gave binary eutectic crystals of mp 92 °C corresponding to Kawamura's glycol.

Another main product **8** was separated as an oil. IR showed the presence of OH and C=O groups. NMR showed an isolated $\text{CH}_2\text{-(C=O)}$ group, four tertiary methyl groups and a cyclopropane ring. NaBH_4 reduction of **8** gave **10** and a *trans*-diol **12**, mp 62–3 °C in a ratio of 2:1. CrO_3 oxidation of **10** in pyridine gave **8**. Thus **8** was 8 β -hydroxythujopsan-9-one. Semicarbazone, mp 230 °C.

The epimeric 8 α -hydroxythujopsan-9-one **9** was obtained as a crystalline minor product, mp 97–8 °C. NaBH_4 reduction of **9** gave **11** and a *trans*-diol **13**, mp 76–8 °C in a ratio of about 2:3. CrO_3 oxidation of **11** gave **9**. α -Ketol rearrangement of **8** was easy as described above. However, the rearrangement of **9** was found to be slow.

4 9 -Thujopsen-8 β -ol (**2**) and -8 α -ol (**3**), the main products under nonaqueous conditions, became minor products.

Experimental

The ^1H NMR spectra were measured with JEOL JNM-PMX 60si in CDCl_3 solution. GLC analyses were performed on a Shimadzu GC-5A gas chromatograph with PEG-HT glass column 1.0 or 1.5 m. The column temperature was elevated from 100 to 220 °C at a rate of 5 °C min^{-1} .

Thujopsene **1** was separated from Hiba wood neutral oil purchased from Soda Aromatic Co., by a spinning band distillation with Tokyo Kagaku Seiki Co., Model SB 818 (1.8 m height) and the fraction of bp 124–6 °C/10 mmHg † (purity 95% up) was used immediately after passing through a silica-gel column to eliminate some autoxidation products.

Oxidation in Dry Acetone: Thujopsene **1** 30.4 g (0.149 mol) was dissolved in 250 ml of dry acetone and 47 g (0.30 mol) of KMnO_4 was added portionwise below 6 °C under nitrogen atmosphere. The mixture was stirred for 12 h and MnO_2 was filtered off. Evaporation of the acetone left 11.8 g of residue. The MnO_2 was washed twice with acetone and additional 7.7 g of extract was obtained. The combined residue was dissolved in hexane and shaken with aqueous Na_2CO_3 solution and NaOH solution successively, giving 14 g of neutral products. Gas chromatography showed unreacted hydrocarbon **1** 33.5%, thujopsadiene **6** 2%, **2** 28.5%, **3** 14.5%, **8** 6.5% and **5** 5%. Extraction of MnO_2 with hot water gave neutral product 1.1 g. Gas chromatography showed 67% of thujopsane-8,9-diol.

† 1 mmHg=133.322 Pa.

Separation by Column Chromatography: The separation was accomplished by passage through an alumina column (320 g). Hexane eluted hydrocarbons and **5** successively. Hexane-CH₂Cl₂ (4:1) eluted **2**, hexane-CH₂Cl₂ (1:1) eluted **3** and CH₂Cl₂ eluted **4**. Δ^9 -Thujopsen-8 β -ol (**2**): Oil. IR (neat) 3400 (OH), 3060, 1665, 1100, 910, and 760 cm⁻¹. ¹H NMR (CDCl₃) δ =0.02 (1H, dd, J =5 and 4 Hz), 0.44 (1H, dd, J =9 and 4 Hz), 0.66 (3H, s), 1.07 (3H, s), 1.17 (3H, s), 1.28 (3H, s), 1.0–1.7 (6H, m), 1.55 (1H, ddd, J =9, 5 and 2 Hz), 5.15 (1H, d, J =10 Hz), and 5.24 (1H, dd, J =10 and 2 Hz). Δ^9 -Thujopsen-8 α -ol (**3**): mp 110–2°C. IR (KBr) 3300 (OH), 3060, 1660, 1125, 1090, 940, 750, and 738 cm⁻¹. ¹H NMR (CDCl₃) δ =0.29 (1H, dd, J =5 and 4 Hz), 0.39 (1H, dd, J =9 and 4 Hz), 0.65 (3H, s), 1.0–1.7 (6H, m), 1.06 (3H, s), 1.15 (3H, s), 1.40 (3H, s), 1.48 (1H, ddd, J =9, 5 and 2 Hz), 4.92 (1H, d, J =10 Hz), and 5.10 (1H, dd, J =10 and 2 Hz). Ring contracted hydroxy ketone **4**: mp 69–70°C. IR (KBr) 3450, 3370, 3200, 1698, 1350, 1235, and 1048 cm⁻¹. ¹H NMR (CDCl₃) δ =0.35–0.82 (2H, m), 0.60 (3H, s), 1.03 (3H, s), 1.13 (3H, s), 1.3–2.3 (7H, m, broad), 1.33 (1H, d, J =14 Hz), 1.72 (1H, d, J =14 Hz), 2.31 (3H, s), and 3.75 (1H, broad). 8 β ,9-Epoxy-10-thujopsanone (**5**): mp 75–7°C. IR (KBr) 3050, 3025, 1690 (C=O), 1372, 1250, 1090, 850, and 818 cm⁻¹. ¹H NMR (CDCl₃) δ =0.00 (1H, dd, J =6 and 5.5 Hz), 0.78 (1H, dd, J =8.5 and 6 Hz), 0.67 (3H, s), 1.05 (3H, s), 1.33 (3H, s), 1.52 (3H, s), 1.3–2.3 (6H, m), 1.77 (1H, dd, J =8.5 and 5.5 Hz) and 3.14 (1H, s).

Isomerization of 8-Hydroxy-9-thujopsanone **8 to **4** by Alumina:** Eighty three mg of **8** was dissolved in hexane and loaded on a column containing 10.5 g alumina deactivated with 1.05 ml of water. After one day, 82 mg of **4** was eluted with ether as the sole product.

Synthesis of **2 and **3** from Mayurone (**7**):** To a Grignard reagent from 0.080 ml of methyl iodide in 3 ml ether was added dropwise 21 mg of mayurone in 1 ml ether. After 15 min at 0°C, the mixture was cooled on a Dry Ice methanol bath, hydrolyzed with aqueous ammonium acetate, and extracted with hexane-ether (1:1). The extract was washed twice with saturated sodium chloride, dried over sodium sulfate and evaporated to give 21 mg of residue. GLC of the residue showed 7% of **2**, and 63% of **3** besides 11% of **6** and 12% of an unidentifiable product.

Lanthanoide-Induced Shift (LIS) of **2 and **3**:** ¹H NMR spectra were measured for ca. 20 mg of **2** and **3** in 0.35 ml CDCl₃ by adding stepwise Eu(fod)₃ and Pr(fod)₃ (3–5 mg for each measurement). The LIS values of CH₃-C(8) were from +2.4 to –8.0 ppm for **2** and from +3.4 to –5.5 ppm for **3**. A good straight line was obtained when the chemical shift values of each proton at various concentrations of the shift reagents were plotted against the corresponding values of CH₃-C(8). The relative value of LIS (1.00 for CH₃-C(8)) was obtained from the slope. The chemical shift values of cyclopropane ring protons were also obtained by extrapolating the concentration of the shift reagents to zero. The results are summarized in Table I.

Oxidation in Aqueous Acetone: Potassium permanganate 14.0 g (88.6 mmol) was added portionwise to a mixture of thujopsene 14.0 g (68.5 mmol) in 140 ml of acetone and 14 ml of water under ice water cooling. The mixture was stirred overnight and filtered. The filtrate was evaporated under reduced pressure, and the residue was dissolved in hexane. The MnO₂ was extracted with hot water several times. The oil which separated from the water

layer was extracted with hexane. The hexane solutions were combined and extracted with aqueous sodium carbonate solution. They afforded 13.4 g of neutral product and 1.7 g of acidic product. Gas chromatography of the neutral product yielded unreacted hydrocarbon **1** 26%, thujopsadiene **6** 0.4%, **2** 8.0%, **3** 4.0%, **8** 22.1%, **9** 4.5% and cis-diols (**10** and **11**) 25.1%.

Separation: The neutral product 27 g was chromatographed on a column (45 mm i.d. $\times$ 49 cm) packed with 350 g of Fuji gel CQ-3 silica gel. The eluent was fed with a Chemco Low Prep Pump Model 81-M-2. Hexane eluted hydrocarbons. Most of **2** and **3** were dehydrated to **6** on a SiO₂ column. Hydroxy ketone **8**, 5.86 g was eluted with hexane-CH₂Cl₂ (4:1). Hydroxy ketone **9**, 0.95 g was eluted with hexane-CH₂Cl₂ (1:1). Then, a mixture of cis-diols, **10** and **11**, 6.95 g was eluted with CH₂Cl₂. A similar oxidation product was chromatographed on alumina containing 10% H₂O gave **4** instead of **8**. **8**: oil. IR (neat) 3480, 3050, 1715, 1140, 1090, and 958 cm⁻¹. ¹H NMR (CDCl₃) δ =0.43–0.86 (2H, m), 0.62 (3H, s), 1.06 (3H, s), 1.1–1.66 (7H, m), 1.20 (3H, s), 1.28 (3H, s), 1.66 (1H, d, J =14 Hz), 2.35 (1H, d, J =14 Hz), and 3.37 (1H, broad). **9**: mp 97–8°C. IR (KBr) 3480, 3070, 3050, 1704, 1168, and 1106 cm⁻¹. ¹H NMR (CDCl₃) δ =0.44–0.80 (2H, m), 0.70 (3H, s), 1.06 (3H, s), 1.18 (3H, s), 1.4–1.75 (7H, m), 1.52 (3H, s), 2.10 (1H, d, J =16.5 Hz), 2.31 (1H, d, J =16.5 Hz) and 2.92 (1H, broad).

Reduction of **8 and **9**:** **8** (385 mg) was dissolved in methanol (20 ml) and reduced with excess NaBH₄. The product (319 mg) was chromatographed on a silica-gel column (10 g). Methylene chloride eluted 175 mg of **10**, mp 68–71°C. Ether eluted 85 mg of 8 β ,9 α -diol **12**. **12**: mp 61–3°C (recrystallized from hexane). IR (Nujol) 3440, 3150, 1170, 1140, 1103, 1080, and 1035 cm⁻¹. ¹H NMR (CDCl₃) δ =–0.01–0.57 (2H, m), 0.55 (3H, s), 0.9–1.9 (9H, m), 0.96 (3H, s), 1.07 (3H, s), 1.18 (3H, s), 2.47 (2H, broad), and 3.75 (1H, X part of ABX, $J_{AX}+J_{BX}$ =16.5 Hz).

Hydroxy ketone **9**, 387 mg was reduced similarly, to give 53 mg of **11**, mp 72–3°C and 82 mg of 8 α ,9 β -diol **13**. **13**: mp 76–8°C (from hexane). IR (Nujol) 3300, 1145, 1106, and 1082 cm⁻¹. ¹H NMR (CDCl₃) δ =0.29–0.66 (2H, m), 0.79 (6H, s), 1.09 (3H, s), 1.1–1.7 (9H, m), 1.33 (3H, s), 2.25–2.57 (3H, m, broad), and 3.51 (1H, X part of ABX, $J_{AX}+J_{BX}$ =16.5 Hz).

References

- 1) Part VIII of this series: S. Nagahama, *Bull. Chem. Soc. Jpn.*, **37**, 1029 (1964).
- 2) Part I: S. Akiyoshi and S. Nagahama, *Bull. Chem. Soc. Jpn.*, **30**, 886 (1957).
- 3) S. Nagahama, H. Kobayashi, and S. Akiyoshi, *Bull. Chem. Soc. Jpn.*, **35**, 1140 (1962).
- 4) Unpublished. The authors are indebted to professor S. Ito, Tohoku University for the IR and NMR spectra.
- 5) J. Kawamura, *Bull. Govt. Forest Expt. Station*, **30**, 59 (1930).
- 6) P. C. Traas, L. M. van der Linde, and H. J. Takken, *Recueil Trav. Chim. Pays-Bas*, **93**, 264 (1974).
- 7) This is corresponding to Ohloff's⁸ Δ^3 -thujopsen-2 α -ol.
- 8) G. Ohloff, H. Strickler, B. Willhalm, C. Borer, and M. Hinder, *Helv. Chim. Acta*, **53**, 623 (1970).