




Scheme 4.

minimal, was treated similarly with thionyl chloride as described in 2. The chlorosulfonium chloride of 5 could not be isolated as a salt in the reaction of 6 with thionyl chloride in the absence of antimony pentachloride. Hydrolysis of the reaction mixture in situ gave sulfoxide 6 in 9%, sulfide 5 in 20%, and aldehyde 7 in 41% yields (Scheme 3). This reaction may have proceeded through the formation of the Pummerer product such as α -chloro sulfide 9 which subsequently is converted into 7 by hydrolysis of 9 as illustrated in Scheme 4. However, an alternative process involving the formation of the sulfonium salt 10 by the reaction of 9 with 5 cannot be excluded (Scheme 4). The structure of 7 was confirmed by the formation of 2,4-dinitrophenyl hydrazone 11. Thus, this distinct difference of reactivity between 2 and 6 should depend on the degree of the transannular S-S interaction in the six- and eight-membered rings.

Experimental

^1H NMR spectra were measured on a Hitachi R-600 FT-NMR spectrometer. The IR spectra were obtained on JASCO A-3 spectrometer. Secondary ion mass spectra were taken by Hitachi M-80B mass spectrometer. Elemental analyses were carried out by the Chemical Analytical Center at this University.

1,5-Dithiacyclooctane was synthesized by a modification of the method of Meadow and Reid.⁶⁾ 1,4-Dithiacyclohexane was obtained from Wako Pure Chemical Industries, Ltd.

Cyclic Bis-Sulfide Monosulfoxides. The monosulfoxides were obtained by general oxidation using *m*-chloroperbenzoic acid.

1,5-Dithiacyclooctane 1-Oxide (2): Yield 76%; mp 27–29 °C; IR (neat) 1010 cm^{-1} (S=O); ^1H NMR (CDCl_3) δ =2.18–2.43 (m, 4H, CH_2), 2.54–2.76 (m, 4H, SCH_2), and 3.04–3.23 [m, 4H, $\text{S}(\text{O})\text{CH}_2$].

1,4-Dithiacyclohexane 1-Oxide (6): Yield 36%; mp 125.5–127.0 °C; IR (KBr) 1020 cm^{-1} (S=O); ^1H NMR (CDCl_3) δ =2.40–2.70 (m, 2H, SCH_2 -e), 2.91–3.22 [m, 4H, $\text{S}(\text{O})\text{CH}_2$], and 3.43–3.77 (m, 2H, SCH_2 -a).

Preparation of 2,2,8,8-Tetradeuterated 1,5-Dithiacyclooctane 1-Oxide (2a). Deuteration of sulfoxide 2 was carried out in $\text{NaOD-D}_2\text{O}$ under nitrogen atmosphere at 100 °C for 24 h. After usual work-up the tetradeuterated

sulfoxide 2a was obtained in 91% yield; the deuterium content of 2a was more than 95 atom% by ^1H NMR spectroscopy.

Isolation of Chlorosulfonium Chloride 3. Addition of pure thionyl chloride (1 mmol) in anhydrous dichloromethane (1 ml) to a solution of 1,5-dithiacyclooctane 1-oxide (2) (1 mmol) in anhydrous dichloromethane (1 ml) under N_2 atmosphere at 0 °C resulted in a white crystalline precipitate. Upon filtration in a dry box under rigorously anhydrous conditions, 1-chloro-5-thia-1-thioniacyclooctane chloride (3) was obtained in 74% yield as stable hygroscopic crystalline salt: Mp 80 °C; SIMS m/z (rel intensity) 185(42), 183(100) 151(17), 107(44), and 73(23).

Reactions of Sulfoxide 2 with Thionyl Chloride in the Presence of Antimony Pentachloride. To a stirred solution of sulfoxide 2 (1 mmol) and antimony pentachloride (1 mmol) in anhydrous dichloromethane (1 ml) was added thionyl chloride (1 mmol) in anhydrous dichloromethane under nitrogen at 0 °C. 1-Chloro-5-thia-1-thioniacyclooctane hexachloroantimonate (4) was obtained in 75% yield as stable hygroscopic crystalline salt: Mp 106–107 °C (decomp); ^1H NMR (CD_3CN) δ =2.32–2.60 (m, 4H, CH_2), 2.71–3.52 (m, 4H, SCH_2), and 3.58–4.16 [m, 4H, $\text{S}(\text{Cl})\text{CH}_2$]; MS m/z 183 (M^+-SbCl_6).

Hydrolysis of Chlorosulfonium Salt 3. The salt 3 was treated with aqueous NaHCO_3 on ice bath. The mixture was extracted with dichloromethane. The combined organic phase was dried over MgSO_4 , filtered, and concentrated under vacuum to give sulfoxide 2 in 70% and sulfide 1 in 15% yields. The salt 4 was hydrolyzed by the same procedure as 3.

Reaction of Sulfoxide 6 with Thionyl Chloride. To a stirred solution of 1,4-dithiacyclohexane 1-oxide (6) (1 mmol) in anhydrous dichloromethane (1 ml) was added thionyl chloride (1 mmol) in anhydrous dichloromethane (1 ml) under nitrogen at 0 °C. Then the reaction mixture was treated with aqueous NaHCO_3 to afford sulfide 5 in 20%, sulfoxide 6 in 9%, and aldehyde 7 in 41% yields. The hydrazone of 7 was prepared as follows. To a solution of aldehyde 7 (19 mg) in ethanol (3 ml) was added 2,4-dinitrophenylhydrazine (50 mg, 0.1% HCl solution). The reaction mixture was stirred at room temperature for 2 h. The solution was extracted with carbon tetrachloride, and the extract was dried over Na_2SO_4 . After removal of the solvent, hydrazone 11 was purified by silica-gel column chromatography. 11: Mp 125–126 °C; IR (KBr) 1610 cm^{-1} (C=N), 1504 and 1321 cm^{-1} (NO_2); ^1H NMR (CDCl_3) δ =2.70–3.19 (m, 10H), 3.47 (d, $J=6\text{ Hz}$, 2H, $\text{SCH}_2\text{CH}=\text{N}$), 4.10 (m, 1H, SCHS), 7.47 (t, $J=6\text{ Hz}$, 1H, $\text{CH}=\text{N}$), 7.92 (d, $J=9\text{ Hz}$, 1H, ArH), 8.36 (dd, $J=9, 3\text{ Hz}$, 1H, ArH), 9.13 (d, $J=3\text{ Hz}$, 1H, ArH), and 11.11 (br s, 1H, NH). Found: C, 38.69; H, 4.17; N, 12.89%. Calcd for $\text{C}_{14}\text{H}_{18}\text{N}_4\text{O}_4\text{S}_4$: C, 38.38; H, 4.14; N, 12.85%.

References

- 1) a) F. G. Bordwell and B. M. Pitt, *J. Am. Chem. Soc.*, **77**, 572 (1955); b) W. E. Truce, G. Birum, and E. T. McBee, *J. Am. Chem. Soc.*, **74**, 3594 (1952); c) H. Böhme and H. J. Gran, *Justus Liebigs Ann. Chem.*, **577**, 68 (1952).
- 2) J. T. Doi and W. K. Musker, *J. Am. Chem. Soc.*, **100**, 3533 (1978); W. K. Musker, *Acc. Chem. Res.*, **13**, 200 (1980).
- 3) N. Furukawa, A. Kawada, and T. Kawai, *J. Chem. Soc., Chem. Commun.*, **1984**, 1151.
- 4) N. Furukawa, A. Kawada, T. Kawai, and H. Fujihara, *J. Chem. Soc., Chem. Commun.*, **1985**, 1266.
- 5) In the ^1H NMR spectrum of $[\text{2H}_4]\text{-4}$ in CD_3CN , the splitting pattern is complicated and could not be resolved.
- 6) J. R. Meadow and E. E. Reid, *J. Am. Chem. Soc.*, **56**, 2177 (1934).