

An Unusual Cyclization of Trifluoroacetohydroximoyl and -hydrazonoyl Bromides with Malononitrile¹⁾

Kiyoshi TANAKA,* Takashi SUZUKI, Seiji MAENO, and Keiryō MITSUHASHI


Department of Industrial Chemistry, College of Technology, Seikei University, Musashino, Tokyo 180
(Received June 23, 1987)

Synopsis. Trifluoroacetohydroximoyl bromide etherate (**1**) reacted with an excess of malononitrile in the presence of sodium methoxide, giving a fused isoxazolopyridine **3**, participated by two molecules of malononitrile, along with the conventional isoxazole **2**. Similar concomitant formation of **7** and the fused pyrazolopyridines **8** was also recognized from the hydrazonoyl bromides **6** and their product ratio was found to depend on the reaction conditions, particularly on the concentration of malononitrile used.

It has been well explored that nitrile oxides and nitrilimines offer a wide variety of isoxazoles and pyrazoles, respectively, through the reaction with the doubly activated methylene compounds.^{2–4)} So far, we have demonstrated the successful synthetic routes to 3-trifluoromethylisoxazoles and -pyrazoles by the reac-

tions of trifluoroacetohydroximoyl bromide etherate (**1**) and *N*-aryltrifluoroacetohydrazonoyl bromides (**6**), respectively, with various active methylene compounds such as β -keto esters,⁵⁾ β -diketones, and cyanoacetates.⁶⁾ As an extension of this study, we now wish to describe an unusual cyclization of **1** and **6** with two molecules of malononitrile other than the conventional cyclization giving 5-amino-4-cyanoazole derivatives.

The bromide **1** reacted with an equimolar malononitrile in the presence of sodium methoxide to give 5-amino-4-cyano-3-trifluoromethylisoxazole (**2**) in 45% yield (Table 1). The orientation of the cycloaddition was deduced by the chemical shifts of its 4- and 5-carbons compared with those of 4-methoxycarbonyl analogue **2'**⁶⁾ (Table 2). However, the treatment with an excess of malononitrile resulted in the formation of brown tarry matter, from which 4,6-diamino-5-cyano-3-trifluoromethylisoxazolo[5,4-*b*]pyridine (**3**) could be isolated in 18% yield. Thus obtained bicyclic product **3** was converted into the 6-chloro analogue, 4-amino-6-chloro-5-cyano-3-trifluoromethylisoxazolo[5,4-*b*]pyridine (**4**), by diazotization followed by chlorination with hydrochloric acid. The structure of **3** and **4** was supported by their elemental analysis and spectral data, particularly by ¹³C NMR analysis (Table 2). The similar behavior producing the fused pyridine was also recognized in case of the hydrazonoyl bromides **6**. Thus, the reaction of **6a** with three times of malononitrile afforded a mixture of 5-amino-4-cyano-1-phenyl-3-trifluoromethylpyrazole (**7a**) (38%) and 4,6-diamino-



Scheme 1.

Table 1. Analytical and Spectral Data of Products

Compound	Formula	Calcd (Found)/%			IR ν/cm^{-1}	¹ H NMR δ
		C	H	N		
2	C ₅ H ₂ F ₃ N ₃ O	33.91 (33.70)	1.14 (1.12)	23.73 (23.83)	3340, 3250, 3200, 3160 (NH ₂) 2240 (C=N), 1660 (C=N)	7.5 (br. s)
3	C ₈ H ₄ F ₃ N ₅ O	39.52 (39.70)	1.66 (1.57)	28.80 (28.71)	3442, 3330, 3220 (NH ₂) 2199 (C=N), 1638 (C=N)	6.7 (br. s, 2H) 7.1 (br. s, 2H)
4	C ₈ H ₂ F ₃ N ₄ OCl	36.59 (36.77)	0.77 (0.68)	21.34 (21.38)	3475, 3330, 3225 (NH ₂) 2230 (C=N), 1655 (C=N)	3.2 (br. s)
7a	C ₁₁ H ₇ F ₃ N ₄	52.39 (52.05)	2.80 (2.74)	22.22 (22.56)	3458, 3368, 3312, 3212, 3190 (NH ₂), 2224 (C=N), 1642 (C=N)	5.65 (br. s, 2H) 7.5 (s, 5H) ^{a)}
7b	C ₁₁ H ₆ F ₃ N ₅ O ₂	44.46 (44.69)	2.03 (1.99)	23.56 (23.76)	3405, 3309, 3219, 3195 (NH ₂) 2205 (C=N), 1640 (C=N)	7.5 (br. s, 2H) 8.15 (A ₂ X ₂ , 4H)
8a	C ₁₄ H ₉ F ₃ N ₆	52.84 (52.90)	2.85 (2.76)	26.41 (26.70)	3505, 3461, 3323, 3231, 3209 (NH ₂), 2200 (C=N), 1661, 1640 (C=N)	6.6 (br. s, 2H), 6.85 (br. s, 2H) 7.1–8.3 (m, 5H)
8b	C ₁₄ H ₈ F ₃ N ₇ O ₂	46.29 (46.08)	2.22 (2.18)	26.99 (26.84)	3464, 3372, 3332, 3239 (NH ₂) 2199 (C=N), 1661 (C=N)	6.8 (br. s, 2H), 7.1 (br. s, 2H) 8.5 (A ₂ X ₂ , 4H)
10	C ₁₄ H ₇ F ₃ N ₅ OCl	49.80 (50.29)	2.09 (2.07)	20.74 (20.27)	3480, 3325, 3220 (NH ₂) 2230 (C=N), 1660 (C=N)	3.3 (br. s, 2H) 7.4–8.0 (m, 5H)

a) Measured in deuteriochloroform.

The IR spectra were recorded on a JASCO A-100 spectrometer. Samples were run as potassium bromide pellets. The ^1H and ^{13}C NMR spectra were measured with a JEOL

JNM-GX270 spectrometer using tetramethylsilane as an internal standard, the chemical shifts being given in δ ppm downfield. Samples were prepared by dissolving in deuteriochloroform-bis(deuteriomethyl) sulfoxide unless otherwise noted. The UV spectra were observed with a Hitachi 340 spectrometer. The MS spectra were obtained on a Finnigan 4023 GC-MS DS spectrometer. The bromides **1** and **6** were prepared by the methods reported in our previous paper.⁶

Reaction of 1 with Malononitrile. (A) A solution of **1** (3.65 g, 13.7 mmol) in 20 cm³ of methanol was rapidly added dropwise to a solution of malononitrile (0.91 g, 13.8 mmol) and sodium methoxide (0.75 g, 13.9 mmol) in 25 cm³ of methanol. After stirring at room temperature for 2 h, the solvent was removed to leave a residue which was extracted with diethyl ether. The extracts were washed with water and brine, dried over magnesium sulfate, and evaporated. The resulting residue was chromatographed on silica gel with chloroform to give 1.10 g (45%) of the isoxazole **2**, mp 149–150°C (recrystallized from hexane-chloroform).

(B) Three times of malononitrile was reacted on a similar procedure to the above and the solvent was removed. The resulting mixture was washed with chloroform and extracted with acetone. The extracts were evaporated and the residual solid was sublimed, giving 0.60 g (18%) of **3** which was further purified by recrystallization from chloroform-acetone (sublimed at around 200°C).

Reaction of 6a with Malononitrile. (A) To a solution of malononitrile (0.89 g, 13.5 mmol) and sodium methoxide (0.73 g, 13.5 mmol) in 20 cm³ of methanol was added a solution of **6a** (1.20 g, 4.5 mmol) in 10 cm³ of methanol. After the mixture was refluxed for 2 h, the solvent was removed to leave a residue which was extracted with diethyl ether. The extracts were dried over magnesium sulfate and evaporated under reduced pressure. To the residue was added excess hexane and the deposited solid was collected on a filter to give 0.05 g (3%) of **8a** which was recrystallized from acetone-methanol, mp 249.5–250.5°C. On the other hand, hexane-extracts were evaporated and the resulting mixture was chromatographed (silica gel, hexane-chloroform, 4:1) to give 0.43 g (38%) of **7a**, which was further purified by recrystallization from hexane-chloroform, mp 115–116°C, along with 0.20 g (20%) of oily **9a**. The spectral data of **9a** are consistent with those of our authentic sample.⁷

(B) A solution of **6a** (4.80 g, 18.0 mmol) in 20 cm³ of methanol was added to a solution of malononitrile (3.56 g, 53.9 mmol) and sodium methoxide (2.92 g, 54.1 mmol) in 40 cm³ of methanol and the mixture was heated under reflux for 2 h. Usual work-up yielded 1.01 g (22%) of **7a** and 2.27 g (40%) of **8a**.

Reaction of 6b with Malononitrile. A mixture of malononitrile (0.76 g, 11.5 mmol), sodium methoxide (0.62 g, 11.5 mmol), and **6b** (1.20 g, 3.8 mmol) in 30 cm³ of methanol was stirred at room temperature for 15 h. The formed solid was collected on a filter, washed with water, and dried at the pump to give 0.10 g (7%) of **8b** which was recrystallized from chloroform-ethyl acetate, mp 309°C (decomp). On the other hand, the filtrate was concentrated and chromatographed (silica gel, hexane-ethyl acetate, 1:1) to give 0.72 g (64%) of **7b** which was further purified by recrystallization from chloroform-ethyl acetate, mp 213–214°C.

Chlorination of 3 into 4. Isoxazolopyridine **3** (0.40 g, 1.6 mmol) was dissolved in 50 cm³ of 6 mol dm⁻³ hydrochloric

acid under reflux. To this solution was added dropwise a solution of 0.40 g of sodium nitrite in 5 cm³ of water at the same temperature. The mixture was refluxed for 2 h and cooled overnight in a refrigerator. The deposited solid was collected on a filter and passed through a short column (silica gel, hexane-ethyl acetate, 3:1) to give 0.09 g (21%) of **4**, which was recrystallized from hexane-diethyl ether, mp 165–166°C.

Chlorination of 8a into 10. In a similar manner to the above, diazotization of 0.25 g of **8a** was carried out at the same temperature for 10 min, giving 0.13 g (49%) of **10**, mp 221–222°C (recrystallized from hexane-ethanol).

Reaction of 1 with Dimeric Malononitrile. A mixture of 2.40 g (18.2 mmol) of dimeric malononitrile and 1.00 g (18.5 mmol) of sodium methoxide in 30 cm³ of methanol was stirred for 40 min. And to this mixture was added dropwise a solution of 4.00 g (15.0 mmol) of **1** in 10 cm³ of methanol over a period of 30 min. After stirring at room temperature for 25 h, the solvent was removed to leave a residue which was extracted with acetone and evaporated under reduced pressure. The resulting mixture was chromatographed (silica gel, hexane-ethyl acetate, 4:1) to give 0.32 g (9%) of **5** which was further purified by recrystallization from hexane-ethanol, mp 290–291.5°C (decomp), IR 3460, 3330, 3230, 3160 (NH₂), 2210 (C≡N), 1630 cm⁻¹ (C=N, C=C), ¹H NMR δ =7.1 (br.s, 2H), 7.4 (br.s, 2H), ¹³C NMR δ =159.6, 157.3, 150.8 (J_{CCF} =33.3 Hz), 118.5 (J_{CF} =277.8 Hz), 112.4, 111.4, 78.5, 72.2, MS (CI, m/z) 228 (M+H)⁺, 208 (M+H-HF)⁺, UV $\lambda_{\text{max}}^{\text{EtOH}}$ (e) 238 (4.143×10⁴), 324 (5.516×10³), 328 (5.600×10³).

Found: C, 42.19; H, 1.49; N, 30.64%. Calcd for C₈H₄F₃N₅: C, 42.30; H, 1.77; N, 30.83%.

References

- 1) Part XIII in "Applications of the fluorinated 1,3-dipolar compounds as the building blocks of the heterocycles with fluorine groups." Part XII: K. Tanaka, M. Ohsuga, Y. Sugimoto, Y. Okafuji, and K. Mitsunashi, *J. Fluorine Chem.*, in press.
- 2) S. A. Lang, Jr. and Y. -i Lin, "Isoxazoles and their Benzo Derivatives," in "Comprehensive Heterocyclic Chemistry," ed by A. Katritzky and C. W. Rees, Pergamon Press, Oxford (1984); A. S. Shawali and C. Párkányi, *J. Heterocyclic Chem.*, **17**, 833 (1980).
- 3) C. Grundmann and S. K. Datta, *J. Org. Chem.*, **34**, 2016 (1969).
- 4) A. O. Abdehamid, C. Párkányi, A. S. Shawali, and M. A. Abdalla, *J. Heterocyclic Chem.*, **21**, 1049 (1984).
- 5) K. Tanaka, M. Kishida, S. Maeno, and K. Mitsunashi, *Bull. Chem. Soc. Jpn.*, **59**, 2631 (1986).
- 6) K. Tanaka, T. Suzuki, S. Maeno, and K. Mitsunashi, *J. Heterocyclic Chem.*, **23**, 1535 (1986).
- 7) For the formation of **9a** in the reaction of **6a** with sodium methoxide in methanol, see K. Tanaka, S. Maeno, and K. Mitsunashi, *Bull. Chem. Soc. Jpn.*, **58**, 1841 (1985).
- 8) H. Junek, B. Thierrichter, and G. Lukas, *Chem. Ber.*, **113**, 1195 (1980).
- 9) H. Junek, M. Mittelbach, and B. Thierrichter, *Monatsh. Chem.*, **110**, 1279 (1979); A. Sakurai and H. Midorikawa, *Bull. Chem. Soc. Jpn.*, **41**, 430 (1968).