

Facile Conversion of Aldehydes into 1-Substituted 2-Chloro-3,3-difluoro-2-propen-1-ols with 1,1-Dibromo-1-chloro-2,2,2-trifluoroethane/Magnesium Reagent


Makoto FUJITA and Tamejiro HIYAMA*

Sagami Chemical Research Center, 4-4-1 Nishiohnuma, Sagamihara, Kanagawa 229

(Received May 13, 1988)

Synopsis. Aldehydes RCHO are converted in excellent yields into chlorodifluoroallyl alcohols RCH(OH)CCl=CF₂ upon treatment with CClBr₂CF₃ and magnesium in tetrahydrofuran at 0 °C.


Aldehyde adducts of halogenated ethenyl anions -CX=CF₂ (X=Cl or F) have been shown to be useful intermediates for the synthesis of α-halo-α,β-unsaturated ketones and acid derivatives,¹⁾ as well as artificial pyrethroids.²⁾ The aldehyde addition of CX=CF₂ anions was earlier realized at -130 °C with LiCX=CF₂,^{1a-c)} and recently achieved under milder conditions using R₃SiCX=CF₂/F⁻,³⁾ CF₃CCl₃/Zn/AlCl₃(cat),⁴⁾ or CF₃CCl₃/Pb^{1h)} reagent. Even these reagents possess problems from practical viewpoints. For example, preparation of the fluorinated vinylsilanes is tedious;⁵⁾ the zinc and lead reagents seem to lack wide applicability.⁶⁾ Reported herein is an alternative facile way to 1-substituted 2-chloro-3,3-difluoro-2-propen-1-ols **2** through the reaction of aldehydes **1** with CClBr₂CF₃ and magnesium, both of which are commercially readily available and economical (Eq 1).



a: R = Ph-

b: R = 4-Cl-C₆H₄-

c: R = PhCH=CH-

d: R = PhCH(CH₃)-e: R = *c*-C₆H₁₁-

Results and Discussion

The adducts **2** were produced in excellent yields, when aldehydes were treated with 3 molar equivalents of 1,1-dibromo-1-chloro-2,2,2-trifluoroethane and 6 mol of magnesium chips in tetrahydrofuran (THF). Results are summarized in Table 1. In addition to common aliphatic and aromatic aldehydes, readily enolizable aldehydes (e.g. **1d** and **1f**) give the desired adducts. Cinnamaldehyde (**1c**) afforded a 1,2-adduct only. The adduct **2f** is a versatile intermediate for synthetic pyrethroids having a CH=C(Cl)CF₃ side chain.²⁾

The reaction course of the transformation deserves

particular comments. When 1:1 mole ratio of CClBr₂CF₃ and Mg was applied, any kinds of aldehyde adducts were not detected, but formation of CClBr=CF₂ was observed by ¹⁹F NMR of the crude reaction mixture. Thus, the organomagnesium reagent responsible for the carbonyl addition is concluded to be a Grignard reagent BrMgCCl=CF₂ generated in situ by the reaction of magnesium metal and the CClBr=CF₂.⁷⁾ In this sense, the mechanism of the total transformation contrasts sharply to that of the CF₃CCl₃/Zn/AlCl₃ (cat) reagent, which first gives aldehyde adducts RCH(OH)CCl₂CF₃ and then reduces the adducts to give products of type **2**.

Although the Grignard reagent BrMgCCl=CF₂ is recorded by Stelin et al.,⁷⁾ its carbonyl addition has never been achieved due possibly to the instability of the reagent. Noteworthy is that the in situ prepared reagent can do the transformation at 0 °C as reported herein.

Experimental

Materials. Magnesium chips and 1,1-dibromo-1-chloro-2,2,2-trifluoroethane were purchased from Nakarai Chemicals Co. Ltd. and PCR Co. Ltd., respectively, and used directly. Aldehydes were purchased from Tokyo Kasei Kogyo Co. Ltd. Ethyl 2-formyl-3,3-dimethylcyclopropanecarboxylate was prepared by a procedure previously reported.^{2b)} Tetrahydrofuran was dried over benzophenone ketyl and freshly distilled before use.

Reaction of Aldehydes with CClBr₂CF₃/Mg Reagent. A Typical Procedure. Magnesium chips 144 mg (6.0 mmol) were suspended in THF 2 mL solution of benzaldehyde (**1a**) 106 mg (1.0 mmol). The whole was cooled at 0 °C, and THF 2 mL solution of 1,1-dibromo-1-chloro-2,2,2-trifluoroethane 825 mg (3.0 mmol) was added dropwise. After 3 h at 0 °C, sat ammonium chloride aq solution 15 mL was added, and the mixture was extracted with diethyl ether (10 mL×3). The ethereal extract was dried over anhydrous magnesium sulfate, concentrated under reduced pressure, and subjected to preparative TLC (silica gel, dichloromethane-hexane 1:1) to give 2-chloro-3,3-difluoro-1-phenyl-2-propen-1-ol 144 mg (**2a**) (70%) as a colorless oil. Spectral data are completely in accord with those of the authentic sample.^{4b)}

Table 1. Transformation of RCHO to RCH(OH)CCl=CF₂^{a)}

Run	Aldehyde	Conditions	Product	Yield/%
1	1a	0 °C, 3 h	2a	70
2	1b	0 °C, 3 h	2b	51
3	1c	0 °C, 4 h	2c	59
4	1d	0 °C, 4 h	2d	77
5	1e	0 °C, 4 h	2e	73
6	1f	0 °C, 4 h	2f	42

a) The molar ratio **1**:CClBr₂CF₃:Mg=1:3:6.

In a similar manner, **2b–f** were prepared. Physical data of **2d**, **2e**, and **2f** have been already reported.^{4b)}

2-Chloro-3,3-difluoro-1-(4-chlorophenyl)-2-propen-1-ol (2b): A colorless oil; ¹H NMR (CDCl₃) δ=2.20 (broad s, 1H), 5.72 (broad s, 1H), and 7.37 (s, 4H); ¹⁹F NMR (CDCl₃-CFCl₃) δ=-87.0 (d, J=38 Hz, 1F) and -91.3 (d, J=38 Hz, 1F); IR (neat) 3400, 1746, 1500, 1296, 1096, 1012, 846, and 734 cm⁻¹; MS (70 eV) m/z (rel intensity) 240 (M⁺+2, 9), 238 (M⁺, 10), 220 (64), 218 (100), 183 (45), 182 (29), 154 (20), 143 (41), 120 (43), 99 (21), 77 (20), 75 (30), 74 (21), and 50 (25).

Found: C, 45.09; H, 2.66%. Calcd for C₉H₆Cl₂F₂O: C, 45.22; H, 2.53%.

2-Chloro-1,1-difluoro-5-phenyl-1,4-pentadien-3-ol (2c): ¹H NMR (CDCl₃) δ=2.17 (broad s, 1H), 5.17 (broad s, 1H), 6.20 (dd, J=5.0, 16.0 Hz, 1H), 6.70 (d, J=16.0 Hz, 1H), and 7.20–7.45 (m, 5H). ¹⁹F NMR (CDCl₃-CFCl₃) δ=-86.3 (d, J=40 Hz, 1F) and -90.0 (d, J=40 Hz, 1F); IR (neat) 3380, 1740, 1288, 1012, 993, 965, 753, and 693 cm⁻¹.

Found: m/z 230.0332. Calcd for C₁₁H₉F₂ClO: M, 230.0310.

References

- 1) a) J. F. Normant, J. P. Foulon, D. Masure, R. Sauvetre, and J. Villieras, *Synthesis*, **1975**, 122. b) D. Masure, R. Sauvetre, J. F. Normant, and J. Villieras, *Synthesis*, **1976**, 761. c) C. Chuit, R. Sauvetre, D. Masure, M. Baudry, J. F. Normant, and J. Villieras, *J. Chem. Res. (S)*, **1977**, 104. d) R. Sauvetre, D. Masure, C. Chet, and J. F. Normant, *Synthesis*, **1978**, 128. e) D. Masure, C. Chuit, R. Sauvetre, and J. F. Normant, *Synthesis*, **1978**, 458. f) J. P. Gillet, R. Sauvetre, and J. F. Normant, *Synthesis*, **1986**, 355. g) F. Tellier, R. Sauvetre, J. F. Normant, and C. Chutt, *Tetrahedron Lett.*, **29**, 3335 (1987). h) H. Tanaka, S. Yamashita, Y. Katayama, and S. Torii, *Chem. Express*, **2**, 751 (1987).
- 2) a) M. Fujita and T. Hiyama, *Tetrahedron Lett.*, **27**, 3659 (1986). b) M. Fujita, K. Kondo, and T. Hiyama, *Bull. Chem. Soc. Jpn.*, **60**, 4385 (1987).
- 3) a) M. Fujita and T. Hiyama, *J. Am. Chem. Soc.*, **107**, 4085 (1985). b) M. Fujita, M. Obayashi, and T. Hiyama, *Tetrahedron*, in press.
- 4) a) M. Fujita and T. Hiyama, *Tetrahedron Lett.*, **27**, 3655 (1986). b) idem, *Bull. Chem. Soc. Jpn.*, **60**, 4377 (1987).
- 5) a) S. Martin, R. Sauvetre, and J. F. Normant, *Tetrahedron Lett.*, **24**, 5615 (1983); *Synthesis*, **1986**, 538. b) T. Hiyama, K. Nishide, and M. Obayashi, *Chem. Lett.*, **1984**, 1765.
- 6) We failed to obtain the CCl=CF₂ adducts of cyclopropanecarbaldehydes and α,β-unsaturated aldehydes with the CF₃CCl₃/Zn/AlCl₃ (cat) reagent.
- 7) CF₂=CFMgBr from CF₂=CFBr: R. N. Stelin, R. D. Yatsenke, L. N. Pinkina, and I. L. Khunyants, *Khim. Nauka Prom.*, **4**, 810 (1959); *Chem. Abstr.*, **54**, 10838.