

Convenient Synthesis of Cyclic Trithiocarbonates from 1,2- or 1,3-Dihaloalkanes and Sodium Trithiocarbonate in the Presence of Phase-Transfer Catalyst

Akira SUGAWARA,* Tsukasa SATO, and Ryu SATO*,†

Department of Industrial Chemistry, Tsuruoka Technical College,
Tsuruoka, Yamagata 997

†Department of Resource Chemistry, Faculty of Engineering Iwate University,
Morioka 020

(Received June 22, 1988)

Synopsis. Cyclic trithiocarbonates, such as 1,3-dithiolane-2-thiones and 1,3-dithiane-2-thiones, were conveniently synthesized by treating 1,2- or 1,3-dihaloalkanes with sodium trithiocarbonate in the presence of a phase-transfer catalyst.

Much attention has been paid to the synthesis of cyclic trithiocarbonates, which are very important intermediates for the preparation of insecticides,¹⁾ biologically active compounds²⁾ and the donor of a superconductor.³⁾ Many procedures for the synthesis of cyclic trithiocarbonates have hitherto been developed: for example, a reaction of potassium *O*-methyl dithiocarbonate or carbon disulfide with epoxide^{4,5)} a reaction of potassium *O*-methyl dithiocarbonate with 2-chloroethanols,⁶⁾ a reaction of sodium trithiocarbonate with 1,3-trimethylene dimesylates,²⁾ and a reaction of ammonium trithiocarbonates with α,ω -dithioalkanes.⁷⁾ These procedures have, however, some problems regarding the availability of the starting materials. In the course of our investigations on the synthesis of alkanediyl and alkyl aryl trithiocarbonates,⁸⁾ we found a convenient synthesis of cyclic trithiocarbonates, 1,3-dithiolane-2-thione (**2a–d**) and 1,3-dithiane-2-thione (**4a–c**), from 1,2- or 1,3-dihaloalkanes with sodium trithiocarbonate in phase-transfer catalyst (PTC) systems (Eqs. 1 and 2).

Triocylmethylammonium chloride (TOMAC) was employed as a phase-transfer catalyst. As shown in




Table 1, 1,3-dithiolane-2-thione (**2a**) was obtained in high yield from 1,2-dibromoethane (Run 4). The reaction in the absence of PTC resulted in a poor yield of 4-methyl-1,3-dithiolane-2-thione (**2b**) (Runs 8–10). Surprisingly, 1,2-ethanedithiol also afforded 1,3-

Table 1. Reaction of Substituted 1,2-Dihaloethane and Sodium Trithiocarbonate in the Presence of PTC

Run ^{a)}	Substrate 1			React		Catalyst	Yield of 2	
	R ¹	R ²	X	Temp/°C	Time/h	mmol	% ^{b)}	
1	H	H	Br	60	8	0	84	2a
2	H	H	Br	60	8	0.02	92	2a
3	H	H	Br	60	3	0.02	76	2a
4	H	H	Br	60	8	0.08	97	2a
5 ^{c)}	H	H	SH	Reflux	2	0	90	2a
6	H	H	Cl	60	8	0.08	58	2a
7	H	H	I	60	8	0.08	— ^{d)}	
8	CH ₃	H	Br	60	8	0	25	2b
9	CH ₃	H	Br	60	8	0.08	67	2b
10	CH ₃	H	Br	60	20	0.08	81	2b
11	C ₂ H ₅	H	Br	60	20	0.08	72	2c
12	CH ₃	CH ₃	Br	60	25	0.08	10	2d
13	C ₆ H ₅	C ₆ H ₅	Br	60	8	0.08	— ^{d)}	

a) 1,2-Dihaloethane, 2 mmol; Na₂CS₃, 6 mmol. b) Isolated yield based on substrate. c) 1,2-Ethanedithiol, 2 mmol; CS₂, 6 mmol; NaOH, 2.5 mmol. d) Products were ethene derivatives.

dithiolane-2-thione (**2a**) in high yield upon treating with carbon disulfide and sodium hydroxide, as shown in Table 1 (Run 5). It should be noted that the treatment of 1,2-diiodoethane and 1,2-dibromo-1,2-diphenylethane with sodium trithiocarbonate gave *trans*-1,2-diphenylethane, since the ethene was a dehalogenates product from the diiodide or dibromide (Runs 7 and 12).⁹ Interestingly, 4-mercaptomethyl-1,3-dithiane-2-thione (**5**) was obtained in good yield upon treating 2,3-dibromo-1-propanol (59%) and 1,3-dibromo-2-propanol (37%) with sodium trithiocarbonate. The formation of **5** may be interpreted in terms of a substitution of the hydroxyl group with trithiocarbonate followed by cyclization as depicted in Scheme 1. The novel thiolane **5** is conceived to be a useful intermediate for the synthesis of biologically active compounds, based on previous results.²⁾


Next, we studied the synthesis of 4-methyl-1,3-dithiane-2-thione (**4b**) by a reaction of 1,3-dibromobutane (**3**) with sodium trithiocarbonate in the presence of PTC. As shown in Table 2, three 1,3-

dithiane-2-thiones (**4a—c**) were obtained in moderate yields. The treatment of 1,4-dibromobutane with sodium trithiocarbonate by the above-mentioned method was not prepared seven-membered cyclic trithiocarbonate.

As mentioned above, 1,3-dithiolane-2-thione **2** and 1,3-dithiane-2-thione **4** were synthesized in satisfactory yields by a reaction of the corresponding 1,2- or 1,3-dihaloalkanes with sodium trithiocarbonate in the presence of PTC. A plausible pathway for this reaction may be considered to be an internal nucleophilic attack of the thiolate anion of intermediate (**6**) as shown in Scheme 2. This novel method for the preparation of cyclic trithiocarbonates has a great advantage from the point of synthetic convenient and utility.

Experimental

All melting points were uncorrected. IR spectra were obtained on a Jasco IR-G spectrophotometer and NMR spectra




Scheme 1.

Table 2. Synthesis of 1,3-Dithiane-2-thiones **4** from Substituted 1,3-Dibromopropane with Na₂CS₃ in the Presence of PTC

Run ^{a)}	Substrate 3			React		Yield of 4 / % ^{b)}	
	R ¹	R ²	X	Temp/°C	Time/h		
1 ^{c)}	H	H	Br	60	1	7	4a
2	H	H	Br	60	1	38	4a
3 ^{d)}	H	H	SH	100	3	45	4a
4	CH ₃	H	Br	r t	1	23	4b
5 ^{d)}	CH ₃	H	Br	60	1	17	4b
6	CH ₃	H	Br	60	1	60	4b
7	CH ₃	H	Br	60	5	66	4b
8	CH ₃	CH ₃	Br	60	20	31	4c

a) 1,3-Dibromopropane, 2 mmol; Na₂CS₃, 6 mmol; TOMAC, 0.08 mmol. b) Isolated yield based on substrate **3**. c) Absence of TOMAC. d) 1,3-Propanedithiol, 2 mmol; CS₂, 6 mmol; NaOH, 2.5 mmol.



Scheme 2.

were measured with a Varian Gemine-200 spectrophotometer (200 MHz) in CDCl_3 using TMS as in internal standard. Elemental analyses were determined with a Yanagimoto MT-3.

General Procedure. To a solution of 1,2-dibromoethane (2 mmol) in benzene (5 ml) were added Na_2CS_3 30% aqueous solution (6 mmol) and trioctylmethylammonium chloride (0.08 mmol); then, the mixture was heated with stirring at 60°C for 8 h. After completion of the reaction, the benzene layer was washed with water and then dried on sodium sulfate. A yellow oil obtained by the evaporation of benzene was chromatographed on silica gel using dichloromethane-hexane (3:1) as an eluent to give the desired 1,3-dithiolane-2-thione (**2a**) in a yield of 97%. The structures of the products were characterized by ^1H NMR, IR, MS, and elemental analyses. The physical and spectral data of thione **2a**, **4a**, and **4b** were identified with those of authentic samples.^{2,9,10}

4-Methyl-1,3-dithiolane-2-thione (2b): Oil; IR (neat) 1032, 1048, and 1077 cm^{-1} ; ^1H NMR (CDCl_3) $\delta=1.65$ (d, 3H, $J=6.6\text{ Hz}$, CH_3), 3.70 (q, 1H, $J=11.8, 7.4\text{ Hz}$, CH), 4.04 (q, 1H, 7.4, 5.7 Hz, CH), and 4.55 (m, 1H, $J=11.8, 6.6$, and 5.7 Hz, CH). Found: C, 32.19; H, 4.09%. Calcd for $\text{C}_4\text{H}_6\text{S}_3$: C, 31.96; H, 4.02%.

4-Ethyl-1,3-dithiolane-2-thione (2c): Oil; IR (neat) 1048 and 1088 cm^{-1} ; ^1H NMR (CDCl_3) $\delta=1.09$ (t, 3H, $J=7.4\text{ Hz}$, CH_3), 1.99 (m, 2H, CH_2), 3.72 (q, 1H, $J=12.0, 7.6\text{ Hz}$, CH), 4.00 (q, 1H, $J=12.0, 5.6\text{ Hz}$, CH), and 4.34 (m, 1H, CH). Found: C, 36.70; H, 4.97%. Calcd for $\text{C}_5\text{H}_8\text{S}_3$: C, 36.54; H, 4.91%.

4,5-Dimethyl-1,3-dithiolane-2-thione (2d): Oil; IR (neat) 1050, 1065, and 1090 cm^{-1} ; ^1H NMR (CDCl_3) (mixture of cis and trans) $\delta=1.52, 1.60$ (d, 6H, CH_3) and 4.10, 4.50 (m, 2H, CH). Found: C, 36.69; H, 4.66%. Calcd for $\text{C}_6\text{H}_8\text{S}_3$: C, 36.54; H, 4.91%.

4,6-Dimethyl-1,3-dithiane-2-thione (4c): Oil; IR (neat) 930, 990, and 998 cm^{-1} ; ^1H NMR (CDCl_3) $\delta=1.46$ (d, 6H,

$J=7.0\text{ Hz}$, CH_3), 2.19 (t, 2H, $J=6.0\text{ Hz}$, CH_2), and 3.64 (m, 2H, $J=7.0, 6.0\text{ Hz}$, CH_2). Found: C, 40.65; H, 5.76%. Calcd for $\text{C}_6\text{H}_{10}\text{S}_3$: C, 40.40; H, 5.63%.

4-Mercaptomethyl-1,3-dithiolane-2-thione (5): Oil; IR (neat) 2500 and 1060 cm^{-1} ; ^1H NMR (CDCl_3) $\delta=1.77$ (t, 1H, $J=9.0\text{ Hz}$, SH), 3.05 (m, 2H, CH_2), 3.99 (q, 1H, $J=12.2, 4.4\text{ Hz}$, CH), 4.19 (q, 1H, $J=12.2, 5.4\text{ Hz}$, CH), and 4.36 (m, 1H, CH). Found: C, 26.45; H, 3.32%. Calcd for $\text{C}_4\text{H}_6\text{S}_4$: C, 26.34; H, 3.32%.

References

- 1) O. Serpold, H. J. Renner, J. Weissflog, and H. Schneider, Ger. (East) Patent 11045 (1956); *Chem. Abstr.*, **53**, 6519c (1959).
- 2) S. Kubota, E. Taniguchi, M. Eto, and K. Maekawa, *Agric. Biol. Chem.*, **41**, 1621 (1977).
- 3) D. B. J. Easton and D. Leaver, *J. Chem. Soc., Chem. Commun.*, **1965**, 585.
- 4) S. M. Iqbal and L. N. Owen, *J. Chem. Soc.*, **1960**, 1030; C. G. Overberger and A. Drucker, *J. Org. Chem.*, **29**, 360 (1964); J. A. Durden, Jr., H. A. Stansbury, Jr., and W. H. Catlette, *J. Am. Chem. Soc.*, **82**, 3082 (1960).
- 5) P. Alain and T. Clande, Fr. Patent 1510016 (1966); *Chem. Abstr.*, **70**, 46876p (1969).
- 6) N. Sawa, M. Hoda, and Y. Hosokawa, Jpn. Patent 7343741 (1973); *Chem. Abstr.*, **81** 13781e (1974).
- 7) Y. Hori, Y. Nagano, T. Fukuhara, S. Teramoto, and H. Taniguchi, *Nippon Kagaku Kaishi*, **1987**, 1408.
- 8) A. Sugawara, M. Shirahata, S. Sato, and R. Sato, *Bull. Chem. Soc. Jpn.*, **57**, 3353 (1984); A. Sugawara, K. Hasegawa, K. Suzuki, Y. Takahashi, and R. Sato, *ibid.*, **60**, 435 (1987).
- 9) J. Nakayama, H. Machida, and M. Hoshino, *Tetrahedron Lett.*, **24**, 3001 (1983).
- 10) S. Tanimoto, T. Oida, K. Hatanaka, and T. Sugimoto, *Tetrahedron Lett.*, **22**, 655 (1981).