

Halogenation Using Quaternary Ammonium Polyhalides. XIX.¹⁾ Aromatic Chlorination of Arenes with Benzyl- trimethylammonium Tetrachloroiodate

Shoji KAJIGAESHI,* Yasuhiro UEDA, Shizuo FUJISAKI, and Takaaki KAKINAMI†

Department of Industrial Chemistry, Faculty of Engineering, Yamaguchi University, Tokiwadai, Ube 755

†Department of Industrial Chemistry, Ube Technical College, Tokiwadai, Ube 755

(Received December 15, 1988)

Synopsis. The reaction of arenes with a calculated amount of benzyltrimethylammonium tetrachloroiodate in acetic acid at room temperature or at 70 °C gave nuclear chloro-substituted arenes in fairly good yields.


Direct aromatic chlorination of arenes (**1**) with chlorine has usually been carried out in the presence of a catalyst such as iodine, iron(III) chloride, or an aluminium-mercury couple. As the chlorinating agents which are easier to handle than toxic gaseous chlorine, sulfuryl chloride,²⁾ iodine trichloride,³⁾ titanium(IV) chloride,⁴⁾ copper(II) chloride,⁵⁾ alumina-supported copper(II) chloride,⁶⁾ antimony(V) chloride,⁷⁾ tellurium(IV) chloride,⁸⁾ trichloroisocyanuric acid,⁹⁾ and poly(*N*-chloromaleimide),¹⁰⁾ etc. have been used to chlorinate **1**.

As one part of our investigation concerning the halogenation of aromatic compounds with quaternary ammonium polyhalides, we have recently found that a new reagent, benzyltrimethylammonium tetrachloro-

iodate (BTMA ICl₄), is an effective chlorinating agent.^{1,11)} In this paper we wish to report on the aromatic chlorination of **1** by the use of BTMA ICl₄.

Results and Discussion

A reaction of **1** with BTMA ICl₄ in acetic acid at room temperature or at 70 °C gave aromatic chloro-substituted arenes (**2**) together with benzyltrimethylammonium dichloroiodate (BTMA ICl₂).¹²⁾ Thus, the produced BTMA ICl₂ is easily separable from the reaction mixture, since BTMA ICl₂ is hardly

Table 1. Aromatic Chlorination of Arenes(**1**) with BTMA ICl₄ in AcOH










Substrate 1	Molar ratio BTMA ICl ₄ / 1	Reaction conditions		Product 2	Yield ^{a)} %	Mp(°C) or Bp(°C/mmHg)	
		Time/h	Temp/°C			Found	Reported
	1.0	48	70		16	132—133/760	131.65—131.66/ ¹³⁾ 760
(1a)				(2a)			
	1.0	24	70		88 ^{c)}	158—161/760	—
(1b)				(2b)			
	1.0	20	70		30 ^{d)}	187—189/760	—
(1c)				(2c)			
	1.0	20	70		66 ^{e)}	213—215/760	—
(1d)				(2d)			
	1.0	20	70		70	194—196/760	193—197/760 ¹⁴⁾
(1e)				(2e)			
	1.0	20	rt		90 ^{f)}	185—187/760	—
(1f)				(2f)			
	1.0	20	70		66	185—187/760	186.8/760 ¹⁵⁾
(1g)				(2g)			

Table 1. (Continued)

Substrate 1	Molar ratio BTMA ICl ₄ /1	Reaction conditions		Product 2	Yield ^{a)} %	Mp(°C) or Bp(°C/mmHg)	
		Time/h	Temp/ °C			Found	Reported
	1.0	20	rt		81	218—220/760	86—87/16 ¹⁶⁾
(1h)				(2h)			
	1.0	20	rt		52	67—68	70.5—71 ¹⁶⁾
(1i)				(2i)			
	1.0	22	rt		90	205—206/760	86—86.8/14 ¹⁷⁾
(1j)				(2j-1)			
1j	2.0	22	70		95	59—60	58—59 ¹⁶⁾
				(2j-2)			
1j	3.0	24	70		98	205.5—206	204—205 ¹⁸⁾
				(2j-3)			
	1.0	20	rt		76	242—244/760	131—132/24 ¹⁶⁾
(1k)				(2k-1)			
1k	2.0	20	rt		96	189—191	193 ¹⁹⁾
				(2k-2)			
	1.0	20	rt		73	43.5—44.5	47.5—48 ¹⁶⁾
(1l)				(2l-1)			
1l	2.0	20	rt		94	187—190	189—189.5 ¹⁶⁾
				(2l-2)			
	1.0	20	rt		96	152—153	155 ²⁰⁾
(1m)				(2m)			
	1.0	24	70		65	256—258/760	106.5/5 ²¹⁾
(1n)				(2n)			
	1.0	18	70		73	271—273/760	132.5—136/12 ²²⁾
(1o)				(2o)			
	1.0	15	70		95	273—275/760	120—122/4.5 ²⁾
(1p)				(2p)			
	1.0	18	rt		40	212—213	214—215 ²³⁾
(1q)				(2q-2)			
1q	2.0	24	rt	2q-2	99	212—213	214—215 ²³⁾

a) Yield of isolated product. b) Product was obtained as a mixture of 4- and 2-chloro derivatives, and its isomer ratio was determined by GC analysis and / or ¹H NMR spectrum. c) 4-Chloro deriv. / 2-chloro deriv. =3/7. d) 4-Chloro deriv. / 2-chloro deriv. =1/1. e) 4-Chloro deriv. / 2-chloro deriv. =7/3. f) 4-Chloro deriv. / 2-chloro deriv. =2/1.

soluble in acetic acid at room temperature. The results are summarized in Table 1.

The reaction of such polyalkylbenzenes as 1,3,5-trimethylbenzene (**1j**), 1,2,3,4-tetramethylbenzene (**1k**), and 1,2,4,5-tetramethylbenzene (**1l**), with equivalent weights of BTMA ICl_4 gave the desired mono-, di-, or trichloro-substituted arenes selectively. For example, the reactions of **1j** with 1.0, 2.0, and 3.0 equiv of BTMA ICl_4 gave 1-chloro- (**2j-1**), 1,3-dichloro- (**2j-2**), and 1,3,5-trichloro-2,4,6-trimethylbenzene (**2j-3**), in good yields, respectively.

We emphasize that the stable reagent BTMA ICl_4 is an excellent chlorinating agent for **1** since it can be treated safely and quantitatively in comparison with toxic gaseous chlorine. Readily separable by-product BTMA ICl_2 can be easily converted to BTMA ICl_4 by treatment with chlorine in dichloromethane.²⁴

However, benzene and isopropylbenzene gave chloro-substituted products in poor yields, respectively, even with a Lewis acid-catalyst such as iron(III) chloride, titanium(IV) chloride, or zinc chloride.

Experimental

1-Chloro-2,4,6-trimethylbenzene (2j-1); General Procedure: To a solution of 1,3,5-trimethylbenzene (**1j**) (0.50 g, 4.16 mmol) in acetic acid (30 ml) was added BTMA ICl_4 (1.74 g, 4.16 mmol). The mixture was stirred for 22 h at room temperature. During the period of stirring BTMA ICl_4 , which was only slightly soluble in acetic acid, gradually reacted with **1j** to give a product and a yellow precipitate. The precipitate (BTMA ICl_2 , 1.02 g, 2.93 mmol) was filtered off. The filtrate was treated with 5% NaHSO_3 (20 ml) and then extracted with hexane (25 ml $\times$ 4). The hexane solution was washed with 5% NaHCO_3 (30 ml) and was purified by column chromatography on alumina. The eluent was concentrated in vacuo to give an oily residue which was distilled, affording **2j-1** as a colorless liquid; yield 0.58 g (90%); bp 205–206°C/760 mmHg (lit.¹⁷) bp 86–86.8°C/14 mmHg).

References

- 1) Part XVIII; S. Kajigaeshi, Y. Shinmasu, S. Fujisaki,

and T. Kakinami, *Chem. Lett.*, **1989**, 415.

- 2) P. B. D. de la Mare and H. Suzuki, *J. Chem. Soc. C*, **1967**, 1586.

- 3) E. Campaigne and W. Thompson, *J. Am. Chem. Soc.*, **72**, 629 (1950).

- 4) G. K. Chip and J. S. Grossert, *Can. J. Chem.*, **50**, 1233 (1972).

- 5) D. C. Nonhebel, *Org. Synth.*, **43**, 15 (1963).

- 6) M. Kodomari, H. Satoh, and S. Yoshitomi, *J. Org. Chem.*, **53**, 2093 (1988).

- 7) P. Kovacic and A. K. Sparks, *J. Am. Chem. Soc.*, **82**, 5740 (1960).

- 8) M. Albeck and S. Shaik, *J. Chem. Soc., Perkin Trans. 1*, **1975**, 1223.

- 9) E. C. Juenge, D. A. Beal, and W. P. Duncan, *J. Org. Chem.*, **35**, 719 (1970).

- 10) C. Yaroslavsky and E. Katchalski, *Tetrahedron Lett.*, **1972**, 5173.

- 11) S. Kajigaeshi, T. Kakinami, H. Ikeda, and T. Okamoto, *Chem. Express*, **3**, 659 (1988).

- 12) S. Kajigaeshi, T. Kakinami, H. Yamasaki, S. Fujisaki, M. Kondo, and T. Okamoto, *Chem. Lett.*, **1987**, 2109.

- 13) R. H. Myers, M. E. Hobbs, and P. M. Gross, *J. Am. Chem. Soc.*, **76**, 4737 (1954).

- 14) D. R. Lyon, F. G. Mann, and G. H. Cookson, *J. Chem. Soc.*, **1947**, 662.

- 15) R. L. Datta and F. V. Fernandes, *J. Am. Chem. Soc.*, **38**, 1809 (1916).

- 16) L. I. Smith and C. L. Moyle, *J. Am. Chem. Soc.*, **58**, 1 (1936).

- 17) G. Illuminati and G. Marino, *J. Am. Chem. Soc.*, **78**, 4975 (1956).

- 18) C. P. Smith and G. L. Lewis, *J. Am. Chem. Soc.*, **62**, 949 (1940).

- 19) A. H. White, B. S. Biggs, and S. O. Morgan, *J. Am. Chem. Soc.*, **62**, 16 (1940).

- 20) L. J. Andrews and R. M. Keefer, *J. Am. Chem. Soc.*, **79**, 5169 (1957).

- 21) A. I. Vogel, *J. Chem. Soc.*, **1948**, 654.

- 22) J. Sauer, R. Huisgen, and A. Hauser, *Chem. Ber.*, **91**, 1461 (1958).

- 23) Y. Nagai and M. Tanabe, *Kogyo Kagaku Zasshi*, **60**, 294 (1957).

- 24) S. Kajigaeshi, T. Kakinami, M. Moriwaki, T. Tanaka, and S. Fujisaki, *Tetrahedron Lett.*, **29**, 5783 (1988).