


Photolysis of Tetraarylmethanes and 3-(Triarylmethyl)pyridines

Min SHI, Yoshiki OKAMOTO,* and Setsuo TAKAMUKU
The Institute of Scientific and Industrial Research, Osaka University,
Ibaraki, Osaka 567
(Received March 12, 1990)

Synopsis. Upon UV irradiation in benzene–methanol (1:2) tetraarylmethanes or 3-(triarylmethyl)pyridines underwent an α,α -elimination of two aryl groups to give biaryls or 3-arylpyridine, and two corresponding carbene intermediates. The latter afforded methyl ethers by the O–H insertion to methanol.

It has been reported that upon UV irradiation in cyclohexane tetraphenylmethane (**1a**) undergoes an α,α -elimination of two phenyl groups to give biphenyl (**2a**) and diphenylcarbene (**3a**), which reacts with cyclohexane to afford diphenylmethane and diphenyl (cyclohexyl)methane.¹⁾ However, its quantum yield is very small ($\phi=7.1\times 10^{-4}$).²⁾ The mechanism of this reaction can be explained as follows^{3,4)}: in an excited state, an intramolecular interaction among the four benzene rings occurs, and two phenyl groups are eliminated via an initial bonding between their C-1 positions (ipso coupling), yielding the carbene **3a** (Scheme 1). An analogous photochemical generation of carbene has also been proposed in triptycene derivatives.⁵⁾



Scheme 1.

Since the interaction among the aryl rings might be more enhanced by the increase of their polarizability by introducing substituents, we prepared and examined some substituted tetraarylmethanes (**1b–i**).

In order to clarify the scope and limitation of this carbene generation, we also prepared 3-(triarylmethyl)pyridines (**4a–e**) and 2-(triphenylmethyl)thiophene (**5**) and studied their photochemical behaviors.⁶⁾


Results and Discussion

Upon UV irradiation of a benzene–methanol (1:2) solution of (4-methoxyphenyl)triphenylmethane (**1d**) (5 mmol dm⁻³) for 1 h in a quartz tube using a high-pressure mercury lamp (300 W) under argon atmosphere, **2a** and 4-methoxybiphenyl (**2d**) were obtained in the yields of 6 and 18% (total quantum yield=0.018), respectively. A mixture of benzene and methanol was used to solubilize **1** and to trap the carbenes **3a** and **3d**.


The elimination of **2a** and **2d** from **1d** resulted in the generation of two kinds of carbenes **3a** and **3d**, affording corresponding methyl ethers **6a** and **6d**⁷⁾ in the yields comparable to those of **2a** and **2d**, respectively (Scheme 2). After 4 h irradiation, the compound (**1d**) was consumed completely and the combined yield of **2a** and **2d** was 50%.

The photolyses of other tetraarylmethanes (**1b, c, f–i**) were carried out under the same conditions. The results are summarized in Table 1 and Scheme 3. These results indicate that the introduction of substituents into the benzene ring causes the preferential formation of the unsymmetric biaryl and the significant increase in total quantum yields of **2**, regardless of the nature of substituent, electron releasing or withdrawing. This suggests that the interaction between the benzene rings in electronically excited state is much enhanced by the introduction of substituents.

However, the introduction of a hydroxyl, an amino, or a phenyl group into the benzene ring depressed the yield of **2a**. For example, even upon a prolonged irradiation of triphenyl(4-hydroxyphenyl)methane (**1g**)



Scheme 2.



Scheme 3.

Table 1. Photolysis of Tetraarylmethane (**1b**—**i**) in Benzene–Methanol (1:2)


Compd			Conv./% ^{a)}	Φ^b	Product (yield/%)			
1	R	R'			2a	6b — i	2b — i	6a
1b	H	CH ₃	40	0.014	(6)	6b (6)	2b (17)	(18)
1c	CH ₃	H	44	0.015	(7)	6c (6)	2c (18)	(18)
1d	OCH ₃	H	46	0.018	(6)	6d (6)	2d (18)	(18)
1e	H	Cl	42	0.014	(6)	6e (3)	2e (16)	(17)
1f	Cl	H	43	0.014	(6)	6f (3)	2f (17)	(17)
1g	OH	H	20	—	(3)	6g (0)	2g (0)	(0)
1h	NH ₂	Cl	25	—	(2)	6h (2)	2h (0)	(0)
1i	C ₆ H ₅	H	10	—	(2)	6i (0)	2i (0)	(0)

a) Irradiated in a quartz tube for 1 h. b) The total quantum yields of **2**.


Table 2. Photolysis of 3-(Triarylmethyl)pyridine (**4a**—**d**) in Benzene–Methanol (1:2)

Compd			Conv./% ^{a)}	Φ^b	Product (yield/%)	
4	R	R'			2	6
4a	H	H	30	0.01	2a (6), 2j (11)	6a (12), 6j (5)
4b	H	OH	5	0	2a (0), 2k (0)	6a (0), 6k (0)
4c	H	Cl	40	0.014	2a (6), 2l (17)	6a (17), 6l (5)
4d	CH ₃	H	40	0.012	2m (8), 2n (10)	6m (10), 6n (7)

a) Irradiated in a quartz tube for 1 h. b) The total quantum yields of **2**.



Scheme 4.



Scheme 5.

or triphenyl(3-methyl-4-aminophenyl)methane (**1h**) in benzene–methanol (1:2), **2a** was obtained only in 2–3% yields, respectively (Table 1). These facts can be interpreted as follows: in an electronically excited singlet state, the hydroxyl group of the phenol derivatives **1g** is apt to dissociate due to the increase of positive charge on oxygen atom.⁸⁾ Therefore, the α,α -elimination of aryl groups of **1g** hardly occurs. In the case of **1h**, the similar interpretation can be also applied. *p*-Biphenyl(triphenyl)methane (**1i**) was also stable upon irradiation. The biphenyl group possesses a lowest T_1 (π,π^*) state, so that an intramolecular energy transfer takes place from the highest (π,π^*) state localized on phenyl group to the lowest state localized on the biphenyl group. As the results, the α,α -elimination of aryl groups of **1i** may be also depressed.

In order to examine the competitive elimination of a heteroaromatic ring, the photolysis of 3-(triphenyl-

methyl)pyridine (**4a**) was carried out under the same conditions. The two products, **2a** and 3-phenylpyridine (**2j**), were obtained. The yield of **2j** (11.0%) was higher than that of **2a** (6.0%) (total quantum yield=0.01). The methyl ethers **6a** and **6j** derived from the corresponding carbenes **3a** and **3j** were likewise obtained (Scheme 4). The substituent effect on the photolysis of **4** was also studied. Upon UV-irradiation of 2-chloro-5-(triphenyl)pyridine (**4c**) or 3-(tri-*p*-tolylmethyl)pyridine (**4d**), the yield of 2-chloro-3-phenylpyridine (**2l**) or 3-*p*-tolylpyridine (**2n**) was larger than that of **2a** or 4,4'-dimethylbiphenyl (**2m**), respectively (Scheme 5, Table 2). These results indicate that the interaction between a phenyl and pyridyl groups is larger than that between phenyl ones.

2-Hydroxy-5-(triphenylmethyl)pyridine (**4b**) and 3-[(4-biphenyl)diphenylmethyl]pyridine (**4e**) were found to be stable upon irradiation. These photo-stability were explained by the same reason described above.

2-Triphenylmethylthiophene (**5**) was also stable upon UV-irradiation in methanol. This may be due to its lower photo-excited energy level of **5** ($UV_{\max}=300$ nm) compared to that of **1** or **4** ($UV_{\max}=260-270$ nm).

Experimental

The compounds **1b-i** and **4a-e** were prepared according to the literature.⁹⁻¹³

Triphenyl(*m*-tolyl)methane (1b).⁹ Mp 162–164 °C; UV (CH_3OH) $_{\max}=265$ nm (ϵ 980 dm³mol⁻¹cm⁻¹).

Triphenyl(*p*-tolyl)methane (1c).⁹ Mp 206–208 °C; UV (CH_3OH) $_{\max}=265$ nm (ϵ 1100 dm³mol⁻¹cm⁻¹).

Triphenyl(4-methoxyphenyl)methane (1d). Mp 200–202 °C; UV(CH_3OH) $_{\max}=272$ nm (ϵ 2000 dm³mol⁻¹cm⁻¹); Found: C, 89.08; H, 6.40%. Calcd for C₂₆H₂₂O: C, 89.11; H, 6.33%.

Triphenyl(3-chlorophenyl)methane (1e).⁹ Mp 190–192 °C; UV(CH_3OH) $_{\max}=266$ nm (ϵ 1000 dm³mol⁻¹cm⁻¹).

Triphenyl(4-chlorophenyl)methane (1f).¹⁰ Mp 216–218 °C; UV(CH_3OH) $_{\max}=270$ nm (ϵ 1300 dm³mol⁻¹cm⁻¹).

Triphenyl(4-hydroxyphenyl)methane (1g).¹⁰ Mp 281–282 °C; UV(CH_3OH) $_{\max}=274$ nm (ϵ 2000 dm³mol⁻¹cm⁻¹).

Triphenyl(3-chloro-4-aminophenyl)methane (1h).⁹ Mp 246–248 °C; UV(CH_3OH) $_{\max}=300$ nm (ϵ 1500 dm³mol⁻¹cm⁻¹).

***p*-Biphenyl(triphenyl)methane (1i).**¹² Mp 197–199 °C; UV(CH_3OH) $_{\max}=263$ nm (ϵ 20000 dm³mol⁻¹cm⁻¹).

3-(Triphenylmethyl)pyridine (4a).¹³ Mp 269–270 °C; UV (CH_3OH) $_{\max}=263$ nm (ϵ 1000 dm³mol⁻¹cm⁻¹).

2-Hydroxy-5-(triphenylmethyl)pyridine (4b).¹³ Mp >300 °C; UV $_{\max}=262$ nm (ϵ 980 dm³mol⁻¹cm⁻¹).

2-Chloro-5-(triphenylmethyl)pyridine (4c).¹³ Mp 256–260 °C; UV(CH_3OH) $_{\max}=263$ nm (ϵ 1100 dm³mol⁻¹cm⁻¹).

3-(Tri-*p*-tolylmethyl)pyridine (4d).¹³ Mp 256–260 °C; UV (CH_3OH) $_{\max}=270$ nm (ϵ 1600 dm³mol⁻¹cm⁻¹); Found: C, 89.10; H, 7.01; N, 3.76%. Calcd for C₂₇H₂₅N: C, 89.22; H, 6.93; N, 3.85%.

3-[(4-Biphenyl)diphenylmethyl]pyridine (4e).¹³ Mp 190–192 °C; UV(CH_3OH) $_{\max}=260$ nm (ϵ 20000 dm³mol⁻¹cm⁻¹).

2-(Triphenylmethyl)thiophene (5). The compound **5** was prepared by the reaction of triphenylmethyl chloride with 2-thienylmagnesium bromide. Mp 194–196 °C (decomp); UV(CH_3OH) $_{\max}=300$ nm (ϵ 400 dm³mol⁻¹cm⁻¹); ¹H NMR (CDCl₃) $\delta=6.76$ (1H, dd, $J=3.6, 1.2$ Hz), 6.95 (1H, dd, $J=5.2, 3.6$ Hz), 7.0–7.50 (16H, m, Ar and $-\text{CH}=\text{C}-$); MS (70 eV) m/z (rel intensity) 326 (M^+ , 80), 249 (100), and 78 (70); Found: C, 84.58; H, 5.60%. Calcd for C₂₃H₁₈S: C, 84.62; H, 5.56%.

Preparation of Some Authentic Samples. Diphenylmethyl Methyl Ether (6a): Diphenylmethyl methyl ether was prepared by the methylation of diphenylmethanol with dimethyl sulfate.¹⁴ ¹H NMR (CDCl₃) $\delta=3.24$ (3H, s, OCH₃), 5.10 (1H, s, $-\text{CH}-$), 7.0–7.40 (10H, m, Ar); Found: m/z 198.1040. Calcd for C₁₄H₁₄O: M, 198.1044.

(3-Methylphenyl)phenylmethyl Methyl Ether (6b): Colorless oil. ¹H NMR (CDCl₃) $\delta=2.20$ (3H, s, CH₃), 3.20 (3H, s, OCH₃), 5.12 (1H, s, $-\text{CH}-$), 7.0–7.40 (9H, m, Ar); Found: m/z 212.1202. Calcd for C₁₅H₁₆O: M, 212.1200.

(4-Methylphenyl)phenylmethyl Methyl Ether (6c): Colorless oil. ¹H NMR (CDCl₃) $\delta=2.21$ (3H, s, CH₃), 3.22 (3H, s, OCH₃), 5.12 (1H, s, $-\text{CH}-$), 7.0–7.40 (9H, m, Ar); Found: m/z 212.1200. Calcd for C₁₅H₁₆O: M, 212.1200.

(4-Methoxyphenyl)phenylmethyl Methyl Ether (6d): Mp 64–66 °C. ¹H NMR (CDCl₃) $\delta=3.24$ (3H, s, OCH₃), 3.64

(6H, s, OCH₃), 5.10 (1H, s, $-\text{CH}-$), 7.0–7.40 (9H, m, Ar); Found: m/z 228.1148. Calcd for C₁₅H₁₆O₂: M, 228.1150.

(3-Chlorophenyl)phenylmethyl Methyl Ether (6e): Colorless oil. ¹H NMR (CDCl₃) $\delta=3.22$ (3H, s, OCH₃), 5.12 (1H, s, $-\text{CH}-$), 7.0–7.50 (9H, m, Ar); Found: m/z 232.0650, 234.0622. Calcd for C₁₄H₁₃OCl: M, 232.0655, 234.0625.

(4-Chlorophenyl)phenylmethyl Methyl Ether (6f): Colorless oil. ¹H NMR (CDCl₃) $\delta=3.24$ (3H, s, OCH₃), 5.10 (1H, s, $-\text{CH}-$), 7.0–7.50 (9H, m, Ar); Found: m/z 232.0648, 234.0620. Calcd for C₁₄H₁₃OCl: M, 232.0655, 234.0652.

The compounds (**6j**, **6l**, and **6n**) were identified by use of high GC-MS measurement, because these compounds were too difficult to prepare by usual methods.

3-(α -Methoxybenzyl)pyridine (6j) Found: m/z 199.0994. Calcd for C₁₃H₁₃NO: M, 199.0996.

2-Chloro-5-(α -methoxybenzyl)pyridine (6l) Found: m/z 233.0606, 235.0577. Calcd for C₁₃H₁₂NOCl: M, 233.0607, 235.0577.

Di-*p*-tolylmethyl Methyl Ether (6m): Colorless oil. ¹H NMR (CDCl₃) $\delta=2.20$ (6H, s, CH₃), 5.10 (1H, s, $-\text{CH}-$), 7.0–7.50 (8H, m, Ar); Found: m/z 210.1406. Calcd for C₁₆H₁₈O: M, 210.1408.

3-(α -Methoxy-4-methylbenzyl)pyridine (6n): Found: m/z 213.1150. Calcd for C₁₄H₁₅NO: M, 213.1153.

A General Procedure of Photolysis. Irradiation and analysis of **1** or **4** were carried out in the same manner as reported previously.⁴

Measurement of the Quantum Yields. The quantum yields were measured on the base of **2** in the same manner as reported previously.⁴

References

- 1) T. D. Walsh and D. R. Powers, *Tetrahedron Lett.*, **1970**, 3855.
- 2) We have determined the quantum yield in the photolysis of tetraphenylmethane under the same conditions as described in experimental.
- 3) M. Shi, Y. Okamoto, and S. Takamuku, *Chem. Lett.*, **1989**, 1297.
- 4) M. Shi, Y. Okamoto, and S. Takamuku, *J. Chem. Soc., Chem. Commun.*, **1989**, 151. M. Shi, Y. Okamoto, and S. Takamuku, *Bull. Chem. Soc. Jpn.*, **63**, 453 (1990).
- 5) H. Iwamura and K. Yoshimura, *J. Am. Chem. Soc.*, **96**, 2652 (1974).
- 6) M. Shi, Y. Okamoto, and S. Takamuku, *Chem. Express*, **5**, 25 (1990).
- 7) G. L. Closs and B. E. Rabinow, *J. Am. Chem. Soc.*, **98**, 8191 (1976).
- 8) T. Matsuura, "Yuuki-hikari-kagaku," Kagaku-dojo, Kyoto (1983), p. 179.
- 9) R. A. Benkeser and R. B. Gosnell, *J. Am. Chem. Soc.*, **78**, 4914 (1956).
- 10) L. H. Welsh and N. L. Drake, *J. Am. Chem. Soc.*, **60**, 58 (1938).
- 11) R. A. Benkeser and W. Schroeder, *J. Am. Chem. Soc.*, **80**, 3314 (1958).
- 12) C. S. Schoepfle and S. G. Trepp, *J. Am. Chem. Soc.*, **54**, 4059 (1932).
- 13) R. Adams, J. Hine, and J. Campbell, *J. Am. Chem. Soc.*, **71**, 387 (1949).
- 14) C. M. Welch and H. A. Smith, *J. Am. Chem. Soc.*, **72**, 4748 (1950).