

Base-Induced Addition-Elimination Reactions of Electron-Deficient Vinylarenes Containing a Carbonyl or Ester Group Utilizing 4-Phenyl-3*H*-1,2,4-triazole-3,5(4*H*)-dione (PTAD)

Kazuyoshi SEGUCHI* and Satoko TANAKA

Faculty of Home Science, Mukogawa Women's University, 6-46 Ikebiraki-cho, Nishinomiya 663


(Received March 22, 1991)

Synopsis. The reaction of electron-deficient vinylarenes such as styrylketones and cinnamate with PTAD afforded double Diels–Alder products and Diels–Alder ene products; the latter underwent novel base-induced elimination of 4-phenyl-1,2,4-triazolidine-3,5-dione via a stable carbanion.

The reactions of 4-phenyl-3*H*-1,2,4-triazole-3,5(4*H*)-dione (PTAD) have been used extensively in organic syntheses, and a diversity of reactions with alkenes have been documented, such as [4+2],¹⁾ [2+2],²⁾ ene,³⁾ and dipolar reactions.^{4,5)} In particular, the reactions of PTAD with electron-rich vinylarenes exemplified by styrene,^{1b)} vinylpyridines,⁶⁾ and benzyldienebicyclic compounds,^{2,5)} resulted in 1:2 adducts. However, no studies have described that the 3,5-dioxo-4-phenyl-1,2,4-triazolidin-1-yl group (PTAD·H) is a latent leaving group. We have found that PTAD·H served as a good leaving group in the reaction products of PTAD with styrylketones or cinnamate.

Results and Discussion

The reaction of 1-phenyl-1-penten-3-one **1a** with PTAD in dichloromethane at 25 °C afforded the double Diels–Alder product **2a** and the Diels–Alder ene product **3a** through a 1:1 adduct as depicted in Scheme 1 in




Scheme 1.

isolated yields of 27% and 24%, respectively. Pursuit of the reaction by ¹H NMR spectroscopy in CDCl₃ at appropriate intervals failed to detect the 1:1 adduct because of the higher reactivity of the diene. The Diels–Alder ene product decomposed partially during purification procedures (silica-gel chromatography and recrystallization). Careful recrystallization from ace-

tonitrile gave pure **3a** as a white powder. The spectral data and the elemental analyses supported the proposed structure. The ¹H NMR showed two doublet signals at δ=5.33 and 6.02 corresponding to 4-H and 5-H, respectively. The ¹³C NMR also showed two doublet signals at δ=52.7 and 63.9. A molecular ion peak (*m/z* 510) was observed on the mass spectrum. When a catalytic amount of triethanolamine was added to an acetonitrile solution of **3a** at 25 °C, the solution turned yellow after a few minutes to afford yellow crystals of **4a** in 94% yield after conventional purifications. This structure was again assigned on the basis of spectral and elemental analyses. The ¹H NMR showed a singlet proton signal at a lower field (δ=6.21) in addition to an ethyl signal (δ=1.25 and 2.32) and a phenyl signal (δ=7.03–8.35). Compound **4a** resulted from the novel base-induced elimination of 4-phenyl-1,2,4-triazoline-3,5-dione (PTAD·H₂). This elimination reaction could be carried out in situ without isolation of **3a**.

Thus, the in situ reaction was extended to a series of electron-deficient vinylarenes such as styrylketones, cinnamate, and cinnamionitrile.⁷⁾ As shown in Table 1, similar successive addition–elimination reactions took place in all cases except for **1h** and **1j**. Concerning **1h**, unidentified products were obtained. **1j** was inert to



Scheme 2.

Table 1. Product Yields of **4** by Addition–Elimination Reaction in situ

	Starting compound		Yield (%) of 4 ^{a)}
	X	R	
1a	H	Et	34 (94)
1b	H	Me	27 (90)
1c	H	Ph	16 (87)
1d	<i>p</i> -OMe	Me	52
1e	<i>p</i> -OMe	Ph	61 (92)
1f	<i>p</i> -Cl	Me	26 (70)
1g	<i>o</i> -Cl	Me	33 (68)
1h	<i>p</i> -NO ₂	Me	0 (30) ^{b)}
1i	<i>p</i> -OMe	OEt	27
1j	Cinnamionitrile		0 (78) ^{b)}

a) Yield based on **1**. The values in parentheses show yields based on isolated crude **3**. b) The values in parentheses show recovered yields of starting materials.

PTAD possibly because of the lower electron-density of the diene.

In order to elucidate the elimination mechanism in particular, the deuterium exchange reaction of **3a** was investigated by ^1H NMR (CDCl_3) spectroscopy in the presence of deuterium oxide and a catalytic amount of sodium carbonate. Addition of sodium carbonate caused the immediate exchange at C-4 more rapidly than the formation of **4a**. Furthermore, the recovered starting material was completely deuterated at C-4. These findings suggest that the elimination reaction may proceed via a carbanion as shown in Scheme 2. Presumably, the electron-withdrawing carbonyl or ester group and amide groups each stabilize the conjugate base of **3**. In summary, the PTAD-H group is a good leaving group in the elimination reaction via a carbanion.

Experimental

Materials and Instruments. Substituted styrylketones, cinnamate, and cinnamionitrile **1a–j** were synthesized according to the reported procedures or obtained from a commercial source. PTAD was freshly prepared by Cookson's method.⁸⁾ The ^1H NMR and ^{13}C NMR spectra were measured in CDCl_3 solution using Me_4Si as an internal standard on a Hitachi R-600 or a JEOL FX-200 spectrometer. The mass spectra were recorded on a JEOL JMS DX-303 spectrometer, operating at an ionizing potential of 70 eV. The IR spectra were taken as KBr disks using a Shimadzu R-460 spectrometer. The elemental analyses were performed using a Yanagimoto Model MT-3 CHN analyzer.

Reaction of 1a with PTAD. PTAD (13.8 mmol) was added portionwise over 2 h in the dark at 25 °C to a solution of **1a** (6.59 mmol) in dichloromethane (10 ml) and the mixture was stirred overnight. The solvent was removed and the residue was flash-chromatographed on silica gel (Wakogel C-200) with dichloromethane as an eluent. **3a** from the first eluate and **2a** from the second eluate were obtained in yields of 24% and 27%, respectively.

2a; Mp 184 °C (acetone); ^1H NMR δ =0.99 (t, 3H), 2.52 (q, 2H), 4.10 (bs, 1H), 5.08–5.30 (m, 1H), 5.30–5.51 (m, 1H), 6.28–6.50 (m, 2H), 6.65–6.90 (m, 2H), and 7.45–7.68 (m, 10H); ^{13}C NMR δ =7.0 (q), 32.1 (t), 52.9 (d), 54.1 (d), 60.0 (d), 62.1 (d), 118.6 (d), 125.4 (d), 125.6 (d), 128.3 (d), 128.4 (d), 129.0 (d), 129.1 (d), 129.3 (d), 130.1 (s), 130.9 (s), 131.0 (d), 131.2 (d), 151.7 (s), 154.8 (s), 154.9 (s), and 200.8 (s); IR 1782, 1776, and 1709 cm^{-1} ; MS m/z (%) 510 (6, M^+) and 333 (100, M^+ –PTADH₂). Found: C, 63.70; H, 4.22; N, 16.44%. Calcd for $\text{C}_{27}\text{H}_{22}\text{N}_6\text{O}_5$: C, 63.50; H, 4.35; N, 16.47%.

3a; Mp 139–140 °C (acetonitrile); ^1H NMR δ =1.06 (t, 3H), 2.62 (q, 2H), 5.33 (d, 1H, J =2 Hz), 6.02 (d, 1H, J =2 Hz), 7.16–7.38 (m, 14H), and 8.40 (bs, 1H); ^{13}C NMR δ =7.1 (q), 32.2 (t), 52.7 (d), 63.9 (d), 114.6 (s), 116.4 (d), 124.7 (d), 125.3 (d), 126.2 (d), 128.4 (d), 128.6 (d), 129.2 (d), 129.7 (d), 130.8 (d), 131.4 (d), 133.2 (s), 146.8 (s), 151.8 (s), 153.8 (s), and 201.5 (s); IR 3160, 1764, and 1708 cm^{-1} ; MS m/z (%) 510 (0.4, M^+) and 333 (100, M^+ –PTADH₂). Found: C, 63.33; H, 4.27; N, 16.42%. Calcd for $\text{C}_{27}\text{H}_{22}\text{N}_6\text{O}_5$: C, 63.50; H, 4.35; N, 16.47%.

General Procedure of in situ Addition-Elimination Reaction. PTAD (13.8 mmol) was added portionwise to the stirred acetonitrile solution (10 ml) of **1** (6.59 mmol) as described above. The mixture was stirred at 25 °C overnight and then a drop of triethanolamine was added to the reaction mixture. The solution turned yellow in a few minutes. After 1 h, conventional purification procedures (concentration by a rotary evaporator, silica-gel column chromatography, and recrystallization) gave yellow crystals.

4a; Mp 202–203 °C (benzene); ^1H NMR δ =1.25 (t, 3H), 2.32 (q, 2H), 6.21 (s, 1H), and 7.03–8.35 (m, 9H); ^{13}C NMR δ =8.1 (q), 34.4 (t), 113.3 (d), 115.4 (d), 119.7 (s), 125.7 (d), 125.9 (d), 128.1 (d), 128.7 (d), 129.2 (d), 130.6 (s), 131.1 (d), 133.2 (s), 134.6 (s), 143.9 (s), 146.2 (s), and 194.6 (s); IR 1746 and 1638 cm^{-1} ; MS m/z (%) 333 (100, M^+). Found: C, 68.56; H, 4.41; N, 12.56%. Calcd for $\text{C}_{19}\text{H}_{15}\text{N}_3\text{O}_3$: C, 68.44; H, 4.54; N, 12.61%.

4b; Mp 201–202 °C (benzene–hexane); ^1H NMR δ =2.48 (s, 3H), 6.31 (s, 1H), and 7.09–8.25 (m, 9H); ^{13}C NMR δ =28.1 (q), 114.7 (d), 115.4 (d), 119.6 (s), 125.7 (d), 125.9 (d), 128.4 (d), 128.7 (d), 129.2 (d), 130.6 (s), 131.4 (d), 133.4 (s), 134.8 (s), 144.1 (s), 146.4 (s), and 190.7 (s); IR 1746 and 1690 cm^{-1} ; MS m/z (%) 319 (100, M^+). Found: C, 67.74; H, 3.98; N, 13.01%. Calcd for $\text{C}_{18}\text{H}_{13}\text{N}_3\text{O}_2$: C, 67.69; H, 4.11; N, 13.17%.

4c; Mp 174–175 °C (benzene–hexane); ^1H NMR δ =6.12 (s, 1H), and 6.99–8.38 (m, 14H); ^{13}C NMR δ =114.2 (d), 115.4 (d), 120.0 (s), 125.8 (d), 127.9 (d), 127.5 (d), 128.8 (d), 129.1 (d), 129.4 (d), 127.9 (s), 130.6 (s), 131.1 (d), 131.6 (s), 133.9 (d), 134.4 (s), 143.1 (s), 145.7 (s), and 185.5 (s); IR 1752 and 1692 cm^{-1} ; MS m/z (%) 381 (100, M^+). Found: C, 72.52; H, 3.77; N, 10.92%. Calcd for $\text{C}_{23}\text{H}_{15}\text{N}_3\text{O}_3$: C, 72.42; H, 3.97; N, 11.06%.

4d; Mp 202–203 °C (benzene); ^1H NMR δ =2.48 (s, 3H), 3.85 (s, 3H), 6.44 (s, 1H), and 6.61–7.90 (m, 8H); ^{13}C NMR δ =27.7 (q), 55.7 (q), 101.5 (d), 111.7 (s), 112.1 (d), 117.0 (d), 125.1 (d), 128.8 (d), 129.3 (d), 129.9 (d), 130.7 (s), 130.8 (s), 136.6 (s), 144.0 (s), 146.8 (s), 162.5 (s), and 190.2 (s); IR 1743 and 1706 cm^{-1} ; MS m/z (%) 349 (100, M^+). Found: C, 65.50; H, 4.24; N, 11.92%. Calcd for $\text{C}_{19}\text{H}_{15}\text{N}_3\text{O}_3$: C, 65.30; H, 4.33; N, 12.04%.

4e; Mp 209 °C (benzene–hexane); ^1H NMR δ =3.86 (s, 3H), 6.25 (s, 1H), and 7.33–8.10 (m, 13H); ^{13}C NMR δ =55.7 (q), 101.6 (d), 111.5 (d), 112.5 (d), 116.6 (s), 125.8 (d), 128.5 (d), 128.7 (d), 129.1 (d), 129.4 (d), 130.7 (s), 133.7 (d), 135.8 (s), 136.1 (s), 143.0 (s), 146.1 (s), 162.2 (s), and 185.3 (s); IR 1758, 1709, and 1653 cm^{-1} ; MS m/z (%) 411 (100, M^+). Found: C, 69.81; H, 4.03; N, 10.45%. Calcd for $\text{C}_{24}\text{H}_{17}\text{N}_3\text{O}_4$: C, 70.05; H, 4.17; N, 10.22%.

4f; Mp 241–243 °C (ethanol); ^1H NMR δ =2.51 (s, 3H), 6.78 (s, 1H), and 7.02–7.58 (m, 8H); ^{13}C NMR δ =28.2 (q), 113.8 (d), 115.8 (d), 118.2 (s), 125.8 (d), 125.9 (d), 128.9 (d), 129.1 (d), 129.4 (d), 130.4 (s), 133.5 (s), 135.5 (s), 137.3 (s), 143.9 (s), 146.3 (s), and 190.4 (s); IR 1762, 1712, and 1682 cm^{-1} ; MS m/z (%) 355 (39, M^+ +2) and 353 (100, M^+). Found: C, 61.17; H, 3.47; N, 11.67%. Calcd for $\text{C}_{18}\text{H}_{12}\text{N}_3\text{O}_3\text{Cl}$: C, 61.09; H, 3.42; N, 11.89%.

4g; Mp 216–217 °C (ethanol); ^1H NMR δ =2.55 (s, 3H), 6.79 (s, 1H), and 7.38–7.58 (m, 8H); ^{13}C NMR δ =28.2 (q), 110.5 (d), 113.7 (d), 118.5 (s), 125.9 (d), 126.5 (d), 128.9 (d), 129.3 (d), 130.5 (s), 131.6 (d), 133.0 (s), 134.4 (s), 143.7 (s), 146.2 (s), and 190.4 (s); IR 1757, 1717, and 1690 cm^{-1} ; MS m/z (%) 355 (41, M^+ +2) and 353 (100, M^+). Found: C, 61.89; H, 3.40; N, 11.60%. Calcd for $\text{C}_{18}\text{H}_{12}\text{N}_3\text{O}_3\text{Cl}$: C, 61.09; H, 3.42; N, 11.89%.

4i; Mp 182 °C (ethanol); ^1H NMR δ =1.37 (t, 3H), 3.84 (s, 3H), 4.37 (q, 2H), 7.07 (s, 1H), and 7.09–7.59 (m, 8H); ^{13}C NMR δ =14.1 (q), 55.7 (q), 62.3 (t), 101.3 (d), 111.4 (d), 112.3 (d), 118.2 (d), 123.0 (s), 125.9 (d), 128.6 (d), 129.2 (d), 129.5 (d), 130.8 (s), 136.9 (s), 143.3 (s), 146.7 (s), 160.2 (s), and 162.3 (s); IR 1780, 1773, 1711, and 1707 cm^{-1} ; MS m/z (%) 379 (100, M^+). Found: C, 63.07; H, 4.42; N, 11.05%. Calcd for $\text{C}_{20}\text{H}_{17}\text{N}_3\text{O}_5$: C, 63.30; H, 4.52; N, 11.08%.

References

- 1) a) M. E. Burrage, R. C. Cookson, S. S. Gupte, and I. D. R. Stevens, *J. Chem. Soc., Perkin Trans. 2*, **1975**, 1325; b) R. C. Cookson, S. S. H. Giliani, and I. D. R. Stevens, *J. Chem.*

Soc. C, **1967**, 1905.

2) W. Adam, V. Lucchini, E.-M. Peters, L. Pasquato, H. G. Schnering, K. Seguchi, H. Walter, and B. Will, *Chem. Ber.*, **122**, 133 (1989).

3) C.-C. Cheng, C. A. Seymour, M. A. Petti, and F. D. Greene, *J. Org. Chem.*, **49**, 2910 (1984).

4) W. Adam, O. De Lucchi, and K. Hill, *Chem. Ber.*, **102**, 6384 (1980).

5) W. Adam, V. Lucchini, L. Pasquato, E.-M. Peters, K.

Peters, H. G. Schnering, and K. Seguchi, *Chem. Ber.*, **119**, 2932 (1986).

6) P. B. Terentev, A. N. Kost, and V. G. Kartsev, *Khim. Geterotsikl. Soedin.*, **1976**, 702; *Chem. Abstr.*, **85**, 94301 (1976).

7) The details of the cycloadditions will be published elsewhere.

8) R. C. Cookson, S. S. Gupte, I. D. R. Stevens, and C. T. Watts, *Org. Synth.*, **51**, 121 (1971).
