

Reactions of 3-Isopropenyl- and 3-Acetyltropolone with Quarternary Ammonium Tribromides

Yoichiro MATSUNAGA and Kimiaki IMAFUKU*

Department of Chemistry, Faculty of Science, Kumamoto University,
Kurokami, Kumamoto 860
(Received July 20, 1991)

Synopsis. Treatments of 3-isopropenyltropolone with quarternary ammonium tribromides in tetrahydrofuran afforded 3-methyl-8*H*-cyclohepta[*b*]furan-8-one. The reactions in methanol–dichloromethane gave 7-bromo-3-methyl-8*H*-cyclohepta[*b*]furan-8-one. Bromination of 3-acetyltropolone with the tribromides in tetrahydrofuran produced 3-(bromoacetyl)-tropolone, while the reaction in the methanolic solvent gave 7-bromo- and 5,7-dibromo-substituted 3-acetyltropolones. The brominations of 4'-hydroxyacetophenone were also carried out.

It is well-known that, on bromination with bromine, tropolones generally give substitution products.^{1–3)} 3-Acetyltropolone was brominated to yield 3-acetyl-5,7-dibromotropolone,⁴⁾ while 3-isopropenyltropolone gave 3-methyl-8*H*-cyclohepta[*b*]furan-8-one and its 5,7-dibromo-substituted product.⁵⁾

Recently, solid quarternary ammonium tribromides have been used as brominating agents in order to avoid the disadvantage of volatile, toxic, and corrosive properties of liquid bromine.^{6,7)} In the treatment of 3-acetyltropolone with phenyltrimethylammonium tribromide, an acetyl group was brominated to yield 3-(bromoacetyl)tropolone, which is applicable as a useful synthon for syntheses of tropolones bearing heterocyclic side chain.⁸⁾

The present paper deals with the reactions of 3-isopropenyl- (**1**) and 3-acetyltropolone (**4**) with tetrabutylammonium tribromide (TBABr₃), benzyltrimethylammonium tribromide (BTMABr₃), and phenyltrimethylammonium tribromide (PTMABr₃). We investigated the differences in the reactivity of the reagents and the effects of solvents on the reactions.

Results and Discussion


Previously, it was found that alkenes are treated with BTMABr₃ to give dibromo adducts on the carbon–carbon double bond.⁹⁾ A solution of 3-isopropenyltropolone (**1**) in an aprotic solvent, tetrahydrofuran, was stirred for 2 h at room temperature in the presence of an equimolar amount of TBABr₃. From the reaction mixture, a cyclized product, 3-methyl-8*H*-cyclohepta[*b*]furan-8-one (**2**),¹⁰⁾ was isolated in a 41% yield. The reactions with BTMABr₃ and PTMABr₃ under the same conditions also gave the compound **2**. These results are summarized in Table 1.

On the other hand, the reactions of **1** with TBABr₃, BTMABr₃, and PTMABr₃ in a protic solvent, methanol–dichloromethane (2:5), yielded the cyclized product **2** and its 7-bromo-substituted product **3** (Table 1). The treatment of the compound **2** with quarternary ammonium tribromide did not give the brominated compound **3**. This means that **1** was initially brominated to produce 7-bromo-3-isopropenyltropolone (not isolated) as an intermediate, which cyclized to the product **3**.

Previously, we reported that bromination of 3-acetyltropolone (**4**) with PTMABr₃ in tetrahydrofuran yielded 3-(bromoacetyl)tropolone (**5**).⁸⁾ In a similar manner, **4** was treated with an equimolar amount of TBABr₃, BTMABr₃, and PTMABr₃ by stirring for 2 h at room temperature to afford **5** (Table 1). From the reaction with TBABr₃, the starting material **4** was also recovered.

The reactions of 3-acetyltropolone (**4**) with quarternary ammonium tribromide in methanol–dichloromethane (2:5) for 1 h gave 3-acetyl-7-bromotropolone (**6**) and 3-acetyl-5,7-dibromotropolone (**7**)⁴⁾ as substitution products (Table 1).

In the chemistry of benzenoid compounds, it was



Scheme 1. a: Quarternary ammonium tribromide/Tetrahydrofuran. b: Quarternary ammonium tribromide/CH₃OH–CH₂Cl₂.

Table 1. Reactions with Quarternary Ammonium Tribromide

Substrate	Reagent	Solvent	Reaction time/h	Product (Yield/%)
1	TBABr ₃	THF	2	2 (41)
1	BTMABr ₃	THF	2	2 (46)
1	PTMABr ₃	THF	2	2 (56)
1	TBABr ₃	CH ₃ OH-CH ₂ Cl ₂	2	2 (54) 3 (7)
1	BTMABr ₃	CH ₃ OH-CH ₂ Cl ₂	2	2 (52) 3 (6)
1	PTMABr ₃	CH ₃ OH-CH ₂ Cl ₂	2	2 (56) 3 (3)
4	TBABr ₃	THF	2	5 (22) 4 ^a) (13)
4	BTMABr ₃	THF	2	5 (38)
4	PTMABr ₃	THF	2	5 (46)
4	TBABr ₃	CH ₃ OH-CH ₂ Cl ₂	1	6 (15) 7 (34)
4	BTMABr ₃	CH ₃ OH-CH ₂ Cl ₂	1	6 (16) 7 (31)
4	PTMABr ₃	CH ₃ OH-CH ₂ Cl ₂	1	6 (19) 7 (54)
8	TBABr ₃	THF	3	9 (49) 8 ^a) (16)
8	BTMABr ₃	THF	2	9 (51) 8 ^a) (27)
8	PTMABr ₃	THF	2	9 (55) 8 ^a) (19)
8	TBABr ₃	CH ₃ OH-CH ₂ Cl ₂	0.5	9 (21) 10 (59)
8	BTMABr ₃	CH ₃ OH-CH ₂ Cl ₂	0.5	9 (16) 10 (69)
8	PTMABr ₃	CH ₃ OH-CH ₂ Cl ₂	0.5	9 (15) 10 (76)


a) Recovered substrate.

reported that the reactions of phenols with BTMABr₃ gave 2,4,6-tribromophenols,¹¹⁾ while the reactions of acetophenones gave 2-bromoacetophenones by using an equimolar amount of BTMABr₃¹²⁾ and 2,2-dibromoacetophenones by using three molar amount of the reagent.¹³⁾ However, the bromination reactions of hydroxyacetophenones bearing both the hydroxyl and acetyl group with quarternary ammonium tribromides were not reported. Thus, in comparison with the reactions of 3-acetyltropolone (**4**), the bromination of 4'-hydroxyacetophenone (**8**) with quaternary ammonium tribromides was carried out.

When a solution of **8** in tetrahydrofuran was treated with an equimolar amount of TBABr₃ at room temperature, the color of the reagent disappeared after stirring for 2 h. The reaction mixture was worked up to give 2-bromo-4'-hydroxyacetophenone (**9**) in a 49% yield, besides the unchanged **8**. The reactions with BTMABr₃ and PTMABr₃ also gave **9** and the unchanged **8** (Table 1). On the other hand, in the reactions in methanol-dichloromethane (2:5), the color of the reagents disappeared after stirring for 30 min and then **9** and 3'-bromo-4'-hydroxyacetophenone (**10**) were obtained in 15–21 and 59–76% yields, respectively (Table 1).

The yields of the reactions by using three quarternary ammonium tribromides increased in order of TBABr₃ < BTMABr₃ < PTMABr₃ in both tetrahydrofuran and methanolic solvent. It can be said that four inductively electron-donating alkyl groups of TBABr₃ repressed the bromination reaction, while the electron-attracting phenyl group of PTMABr₃ accelerated the reactions.

We also found a striking difference in the effects of the solvents on the bromination reactions. As shown in Table 1, the bromination of 3-acetyltropolone (**4**) in the aprotic tetrahydrofuran took place at the side-chain acetyl group, while the reaction in the protic methanol-dichloromethane took place at the 5- and 7-positions in



Scheme 2.

the tropolone nucleus. The reactions of 4'-hydroxyacetophenone (**8**) in tetrahydrofuran afforded the 2-brominated product **9**. The reaction in the methanolic solvent gave two types of products **9** and **10**. The latter was preferential to the former. When the compounds **5** and **9** were stirred at room temperature in the methanolic solvent in absence of the brominating agent, they were recovered quantitatively, respectively. On the other hand, the compounds **5** and **9** were treated with an equimolar amount of PTMABr₃ in methanol-dichloromethane to afford 5,7-dibromo-3-(bromoacetyl)tropolone (**11**) and 2,3'-dibromo-4'-hydroxyacetophenone (**12**) in 58 and 85% yields, respectively. In the reaction of 3-isopropenyltropolone (**1**), the 7-brominated product **3** was found in the methanolic solvent, besides the product **2**.

Thus, in the methanolic solvent, the brominating agent might react with methanol to produce methyl hypobromite^{11,13)} which brominates the tropolone and benzene nucleus as a more active brominating species.

Experimental

The melting points were determined with a Yanagimoto MP-

S2 apparatus and are uncorrected. The IR spectra were taken on a JASCO A-102 spectrophotometer. The ^1H NMR spectra were recorded with a JEOL JNM-PMX60SI spectrometer. The mass spectra were measured on a JEOL DX3-3HF spectrometer.

Reactions of 3-Isopropenyltropolone (1) with Quarternary Ammonium Tribromide. General Procedure. a) Reactions in Tetrahydrofuran. To a solution of 3-isopropenyltropolone (1) (486 mg, 3 mmol) in tetrahydrofuran (10 ml) was added quarternary ammonium tribromide (3 mmol) little by little. After stirring for 2 h at room temperature, the color of the reaction mixture changed gradually from orange to yellow. The mixture was diluted with water (100 ml) and extracted with ether (40 ml $\times$ 4). The extract was evaporated, chromatographed on a Wakogel B-10 plate (30 $\times$ 30 cm) with ethyl acetate-chloroform (1:1), and recrystallized from methanol to give 3-methyl-8*H*-cyclohepta[*b*]furan-8-one (2): Mp 95–97 °C (lit.¹⁰) 95–96 °C).

b) Reactions in Methanol-Dichloromethane. A solution of 3-isopropenyltropolone (1) (486 mg, 3 mmol) in methanol (20 ml)-dichloromethane (50 ml) was treated with quarternary ammonium tribromide (3 mmol) for 2 h and worked up, as described above. The evaporation residue was chromatographed on a Wakogel B-10 plate (30 $\times$ 30 cm) with ethyl acetate-chloroform (1:1). The lower fraction was recrystallized from methanol to give 3-methyl-8*H*-cyclohepta[*b*]furan-8-one (2). The upper fraction was recrystallized from methanol to give 7-bromo-3-methyl-8*H*-cyclohepta[*b*]furan-8-one (3): Pale yellow needles; mp 118–120 °C; IR (CHCl₃) 1597 cm⁻¹ (C=O); ^1H NMR (CDCl₃) δ =2.27 (3H, s, CH₃), 6.83 (1H, dd, *J*=9, 9 Hz, H-5), 7.70 (1H, m, H-2), 7.74 (1H, d, *J*=9 Hz, H-6), 8.23 (1H, d, *J*=9 Hz, H-4). Found: C, 49.98; H, 2.97%. Calcd for C₁₀H₇BrO₂: C, 50.24; H, 2.95%.

Reactions of 3-Acetyltropolone (4) with Quarternary Ammonium Tribromide. General Procedure. a) Reactions in Tetrahydrofuran. A solution of 3-acetyltropolone (4) (492 mg, 3 mmol) in tetrahydrofuran (10 ml) was treated with quarternary ammonium tribromide (3 mmol) for 2 h and worked up, as described above. The extract was evaporated and recrystallized from petroleum ether to give 3-(bromoacetyl)tropolone (5): Mp 95–97 °C (lit.⁸) 97–98 °C).

b) Reactions in Methanol-Dichloromethane. A solution of 3-acetyltropolone (4) (492 mg, 3 mmol) in methanol (20 ml)-dichloromethane (50 ml) was treated with quarternary ammonium tribromide (3 mmol) for 1 h and worked up, as described above. The extract was evaporated and chromatographed on a Wakogel B-10 plate (30 $\times$ 30 cm) with ethyl acetate. The lower fraction was recrystallized from methanol to give 3-acetyl-7-bromotropolone (6): Yellow needles; mp 87–89 °C; IR (CHCl₃) 3220 (OH), 1693 (COCH₃), 1603 cm⁻¹ (C=O); ^1H NMR (CDCl₃) δ =2.76 (3H, s, CH₃), 7.03 (1H, dd, *J*=10, 10 Hz, H-5), 7.50 (1H, d, *J*=10 Hz, H-6), 8.70 (1H, d, *J*=10 Hz, H-4); MS *m/z* (%) 244 [(*M*+2)⁺, 45], 242 (*M*⁺, 46), 201 [(*M*+2)⁺-COCH₃, 91], 199 (*M*⁺-COCH₃, 92), 163 (*M*⁺-Br, 100). Found: C, 44.25; H, 2.84%. Calcd for C₉H₇BrO₃: C, 44.47; H, 2.90%.

The upper fraction was recrystallized from methanol to give 3-acetyl-5,7-dibromotropolone (7): Mp 171–173 °C (lit.⁴) 173–174 °C).

Reactions of 4'-Hydroxyacetophenone (8) with Quarternary Ammonium Tribromide. General Procedure. a) Reactions in Tetrahydrofuran. A solution of 4'-hydroxyacetophenone (8) (408 mg, 3 mmol) in tetrahydrofuran (10 ml) was treated with quarternary ammonium tribromide (3 mmol) for 3 h (TBABr₃) or 2 h (BTMABr₃ and PTMABr₃) and worked up, as described above. The extract was evaporated, chromatographed on a Wakogel B-10 plate (30 $\times$ 30 cm) with ethyl acetate, and recrystallized from benzene to give 2-bromo-4'-

hydroxyacetophenone (9): Mp 128–130 °C (lit.¹⁴) 130 °C).

b) Reactions in Methanol-Dichloromethane. A solution of 4'-hydroxyacetophenone (8) (408 mg, 3 mmol) in methanol (20 ml)-dichloromethane (50 ml) was treated with quarternary ammonium tribromide (3 mmol) for 30 min and worked up, as described above. The extract was evaporated and chromatographed on a Wakogel B-10 plate (30 $\times$ 30 cm) with ethyl acetate. The lower fraction was recrystallized from benzene to give 2-bromo-4'-hydroxyacetophenone (9). The upper fraction was recrystallized from petroleum ether to give 3'-bromo-4'-hydroxyacetophenone (10): Mp 93–94 °C (lit.¹⁵) 92–95 °C).

Reaction of 3-(Bromoacetyl)tropolone (5) with PTMABr₃. A solution of 5 (122 mg, 0.5 mmol) in methanol (3 ml)-dichloromethane (8 ml) was treated with PTMABr₃ (188 mg, 0.5 mmol) for 1 h and worked up, as described above. The extract was evaporated and recrystallized from methanol to give 5,7-dibromo-3-(bromoacetyl)tropolone (11): Yield 116 mg (58%); yellow crystals; mp 162 °C (decomp); IR (CHCl₃) 3200 (OH), 1695 (COCH₂), 1600 cm⁻¹ (C=O); ^1H NMR (CDCl₃) δ =4.59 (2H, s, CH₂), 8.02 (1H, d, *J*=2 Hz, H-6), 8.49 (1H, d, *J*=2 Hz, H-4). Found: C, 26.71; H, 1.50%. Calcd for C₉H₅Br₃O₃: C, 26.96; H, 1.26%.

Reaction of 2-Bromo-4'-hydroxyacetophenone (9) with PTMABr₃. A solution of 9 (108 mg, 0.5 mmol) in methanol (3 ml)-dichloromethane (8 ml) was treated with PTMABr₃ (188 mg, 0.5 mmol) for 30 min and worked up, as described above. The extract was evaporated and recrystallized from benzene to give 2,3'-dibromo-4'-hydroxyacetophenone (12): Mp 140–142 °C (lit.¹⁶) 143 °C).

References

- 1) T. Nozoe, K. Takase, and H. Matsumura, "Dai Yuki Kagaku," ed by M. Kotake, Asakura Shoten, Tokyo (1960), Vol. 13, pp. 1–437.
- 2) F. Pietra, *Chem. Rev.*, **73**, 293 (1973).
- 3) D. Lloyd, "Non-benzenoid Conjugated Carbocyclic Compounds," Elsevier, Amsterdam (1984), pp. 107–125.
- 4) Z.-H. Li, Z.-T. Jin, B.-Z. Yin, and K. Imafuku, *J. Heterocycl. Chem.*, **24**, 779 (1987).
- 5) R.-H. Jin, B.-Z. Yin, Z.-T. Jin, and K. Imafuku, *J. Heterocycl. Chem.*, **27**, 583 (1990).
- 6) L. F. Fieser and M. Fieser, "Reagents for Organic Synthesis," John Wiley & Son, New York (1967), Vol. 1, p. 855; Vol. 2, p. 328 (1969); Vol. 4, p. 386 (1974); Vol. 5, p. 531 (1975); Vol. 11, p. 426 (1984); Vol. 13, p. 287 (1988).
- 7) S. Kajigaeshi and T. Kakinami, *Yuki Gosei Kagaku Kyokai Shi*, **46**, 986 (1988).
- 8) C.-Y. Qian, Z.-T. Jin, B.-Z. Yin, and K. Imafuku, *J. Heterocycl. Chem.*, **26**, 60 (1989).
- 9) S. Kajigaeshi, T. Kakinami, M. Moriwaki, S. Fujisaki, and M. Tashiro, *Rept. Inst. Adv. Mater. Stud., Kyushu Univ.*, **2**, 87 (1988).
- 10) K. Imafuku, K. Yamaguchi, and H. Matsumura, *Bull. Chem. Soc. Jpn.*, **53**, 745 (1980).
- 11) S. Kajigaeshi, T. Kakinami, H. Tokiyama, T. Hirakawa, and T. Okamoto, *Chem. Lett.*, **1987**, 627.
- 12) S. Kajigaeshi, T. Kakinami, T. Okamoto, and S. Fujisaki, *Bull. Chem. Soc. Jpn.*, **60**, 1159 (1987).
- 13) S. Kajigaeshi, T. Kakinami, H. Tokiyama, T. Hirakawa, and T. Okamoto, *Bull. Chem. Soc. Jpn.*, **60**, 2667 (1987).
- 14) N. P. Buu-Hoi, N. D. Xuong, and D. Lavit, *J. Chem. Soc.*, **1954**, 1034.
- 15) F.-C. Chen and T.-H. Tsai, *J. Taiwan Pharm. Assoc.*, **4**, 42 (1952).
- 16) K. W. Rosenmund and K. Pfroepffer, *Chem. Ber.*, **90**, 1922 (1957).