

Generation and Cycloaddition Reactions of 3-Styryl-1,2-thiaphosphole 2-Sulfide and 1,5-Diphenyl-2,4-pentadiene-1-thione

Shinichi MOTOKI,* Takaya SAKAI, Yoshihisa MATSUO,[†] Shunichi KAMETANI,^{††} and Takao SAITO*

Department of Chemistry, Faculty of Science, Science University of Tokyo,
Kagurazaka, Shinjuku-ku, Tokyo 162

[†] Mitsui Petrochemical Industries, Ltd., Chikusakaigan, Ichihara-shi, Chiba 299-01

^{††} Nippon Kayaku Co., Ltd., Shimo, Kita-ku, Tokyo 115

(Received October 9, 1991)

Synopsis. The cycloaddition reaction of 3-styryl-1,2-thiaphosphole 2-sulfide and 1,5-diphenyl-2,4-pentadiene-1-thione derivatives generated by the thermolysis of 5,6-distyryl-3,8-diaryl-2,9-dithia-1-phosphabicyclo[4.3.0]nona-3,7-diene 1-sulfides have been described. Methacrylonitrile, methyl methacrylate, α -methylstyrene, norbornene, methyl acrylate, styrene, and methyl vinyl ketone afforded the cycloadducts or their rearranged products with the thiaphosphole 2-sulfides, whereas diethyl azodicarboxylate, dimethyl acetylenedicarboxylate, and *N*-(*p*-methoxyphenyl)maleimide gave the cycloadducts with the thioketone.


In the previous paper, we reported that treatment of $\alpha,\beta:\alpha',\beta'$ -unsaturated ketones with P_2S_5 gave phosphorus-containing compounds which generated $\alpha,\beta:\alpha',\beta'$ -unsaturated thioketones and 3-aryl-5-(2-arylvinyl)-1,2-thiaphosphole 2-sulfides upon heating. The thioketone is a sulfur-containing cross-conjugated heterotriene and underwent diene-transmissive Diels–Alder reaction with dienophiles successively giving bicyclic cycloadducts.^{1,2)}

In view of the above results, diene-transmissive Diels–

Alder reaction of 5-aryl-3-styryl-1,2-thiaphosphole 2-sulfide (**1**) was now attempted. The compound **1** is a new phosphorus-containing cross conjugated heterotriene and we expected that cisoid constrained $C^5=C^4-C^3=P^2$ moiety in the ring would preferentially react with dienophiles to produce the cycloadduct **5** capable of undergoing successive cycloaddition reaction (Scheme 1).

The reaction of 1,5-diphenyl-2,4-pentadien-1-one with large excess of P_2S_5 in the presence of triethylamine gave 5,6-distyryl-3,8-diphenyl-2,9-dithia-1-phosphabicyclo[4.3.0]nona-3,7-diene 1-sulfide **2a** in 34% yield. The mass spectrum of **2a** exhibited fragment peaks of thiaphosphole **1a**-S (280, 100%) and of thioketone **3a** (250, 33%) with a weak parent peak (562, 0.4%). The 1H NMR spectrum showed six signals of olefinic protons and one methine proton. The ^{13}C NMR spectrum showed signals of one quaternary carbon and one tertiary carbon. The ^{31}P NMR spectrum exhibited a signal at $\delta=120.44$. Similarly, the reaction of 1-*p*-tolyl and 1-(*p*-chlorophenyl)-5-phenyl-2,4-pentadien-1-one with P_2S_5 gave the products **2b** (Ar=*p*-CH₃C₆H₄, yield 34%) and **2c** (Ar=*p*-ClC₆H₄, yield 22%).

As expected from the mass spectrum, **1** and **3** were



Numbering in the parentheses are those of the adducts **6f** and **7f**.

Scheme 1.

Table 1. Reaction of **2** with Dienophiles

Compound 2	Dienophile ^{a)}		Reaction time/h	Product	Yield/% ^{b)}		
	X	Y			6	7	8
2a	CH ₃ OCOC=	=CH ₂	2.0	6a	33	0	0
2b	CH ₃ OCOC=	=CH ₂	2.5	6b	32	0	0
2c	CH ₃ OCOC=	=CH ₂	1.0	6c	26	0	0
2a	NC-C=	=CH ₂	6.0	6d	24	0	0
2a	Ph-C=	=CH ₂	2.0	6e	41	0	0
2a	Norbornene		3.0	6f + 7f	33	33	0
2a	CH ₃ OCOCH=	=CH ₂	4.0	6g + 7g	8	13	8
2a	Ph-CH=	=CH ₂	10.0	6h + 7h	12	18	0
2a	CH ₃ COCH=	=CH ₂	2.0	6i + 7i	2	18	0
2a	DMAD		2.5	8j	0	0	28
2a	DAD		2.0	8k	0	0	43
2a	MI		5.0	8l	0	0	27

a) Molar ratio of dienophile/compound **2** was 3. b) Yields were based on compound **2**.

generated by the thermolysis of **2** and trapped by dienophiles **4** (Table 1). However, the reactions of **2** with methyl methacrylate, methacrylonitrile and α -methylstyrene gave only the adducts **6a–e** which can not be utilized in successive cycloaddition reaction. The ^{13}C NMR (DEPT) spectrum of **6a** exhibited the signals of quaternary ($\delta=47.5$), tertiary ($\delta=49.8$), and secondary ($\delta=47.1$) carbon atoms one by one and the ^1H NMR spectrum showed the signals of unequivalent methylene protons ($\delta=2.80$ and 3.29 , dd, $J_{\text{HH}}=13.7$, $J_{\text{HP}}=13.6$, 15.8 Hz), one methine proton ($\delta=4.48$, dd, $J_{\text{HH}}=4.9$, $J_{\text{HP}}=4.8$ Hz) and two olefinic protons ($\delta=6.87$, 6.88). On the other hand, the reaction with norbornene afforded the adduct **6f** and its isomer **7f** formed by 1,5-prototropy of **6f**. The signal of H-5 methine proton of **7f** appeared at lower field ($\delta=5.95$) than that of H-9 methine proton of **6f** ($\delta=4.71$). In the 2D ^1H NMR (H–H COSY) of **7f**, H-5 was shown to have the coupling only with an olefinic proton (H-6).

The reaction of **2a** with methyl acrylate, styrene and methyl vinyl ketone gave an inseparable mixture of phosphorus-containing adducts in each case. Analysis of the ^1H NMR spectra of the mixture referring above mentioned differences in the spectra of **6f** and **7f** proved that these mixtures were composed of the adducts **6g–i** and their rearranged products **7g–i** respectively. The ratio of **6** to **7** was estimated by the relative intensity of H-4 and H-8 signals.³⁾ The reaction of **2a** with dimethyl acetylenedicarboxylate (DMAD), diethyl azodicarboxylate (DAD), and *N*-(methoxyphenyl)maleimide (MI) resulted to afford only the cycloadducts with the thioketone (**8j–l**).

Preferential formation of **6(7)** rather than **5** is recognized by considering that bicyclic compounds **6(7)** have less steric strain than bridged compounds **5** and that disubstituted C-5 terminal carbon of **1** is less reactive than its mono substituted C-2' terminal carbon. It was also found that there is a marked contrast between the formation of the thiaphosphole adducts **6(7)** and the thioketone adducts **8**, that is, dienophiles DMAD, DAD, and MI gave **8** while the other dienophiles gave **6(7)**. For the purpose of obtaining the both adducts **6(7)** and **8**, the reaction of **2a** with the mixture of methyl methacrylate and DMAD or methyl methacrylate and DAD were examined but only the adduct **8** was formed even under such conditions. Finally, dienophiles of the type $\text{CH}_2=\text{C}(\text{CH}_3)\text{--Z}$ ($\text{Z}=\text{CO}_2\text{CH}_3$, CN, Ph, and COCH_3) produced only **6**, while those of the type $\text{CH}_2=\text{CH}\text{--Z}$ and norbornene produced both **6** and **7**. These interesting results are under further investigation.

Experimental

All the melting points are uncorrected. IR spectra were measured on a Hitachi Model 270-30 spectrometer. ^1H and ^{13}C NMR spectra were measured in CDCl_3 solution using TMS as an internal standard (JEOL JNM-FX 100, JEOL EX-270, JEOL JNM-GSX 500). ^{31}P NMR spectra were measured at 109 MHz on a JEOL EX-270 spectrometer in CDCl_3 solution using 85% H_3PO_4 as an external standard. Mass spectra were recorded on a Hitachi double-focusing mass spectrometer Model RMU-7M operating at an ionizing potential of 70 eV.

General Procedure for the Preparation of 2. A suspension of conjugated dienone (20 mmol), P_2S_5 powder (6 g) and

triethylamine (12 ml) in dry carbon disulfide (80 ml) was gently refluxed under a nitrogen atmosphere for 2–3.5 h. The reaction mixture was filtered and the filtrate was evaporated. The residue was chromatographed on silica gel (Wakogel C-200). The solvent was evaporated and the residue was recrystallized from ethyl acetate–hexane giving the product (**2a–c**).

3,8-Diphenyl-5,6-distyryl-2,9-dithia-1-phosphabicyclo[4.3.0]nona-3,7-diene 1-Sulfide (2a):⁴⁾ Pale yellow crystals; mp $150\text{--}151^\circ\text{C}$. ^1H NMR $\delta=4.09$ (ddd, H-5, $J_{4,5}=2.81$, $J_{5,1}=7.26$, $J_{\text{HP}}=12.87$ Hz), 5.73 (dd, H-4, $J_{4,5}=2.81$, $J_{\text{HP}}=2.81$ Hz), 6.20 (d, H-7, $J_{\text{HP}}=35.30$ Hz), 6.38 (dd, 5-styryl H-1', $J_{5,1'}=7.26$, $J_{1',2'}=15.84$ Hz), 6.44 (dd, 6-styryl H-1', $J_{1',2'}=16.17$, $J_{\text{HP}}=5.94$ Hz), 6.58 (dd, 6-styryl H-2', $J_{1',2'}=16.17$, $J_{\text{HP}}=6.27$ Hz), 6.71 (d, 5-styryl H-2', $J_{1',2'}=15.84$ Hz), $7.21\text{--}7.64$ (m, 20H); ^{13}C NMR (DEPT) $\delta=42.7$ (d-CH, C-5, $J_{\text{CP}}=3.6$ Hz), 63.4 (d-C, C-6, $J_{\text{CP}}=59.9$ Hz); ^{31}P NMR $\delta=120.4$. MS m/z 562 (M^+ ; very weak), 280 ($\text{M}^+\text{--thioketone-S}$; 100). Found: C, 72.85; H, 4.98%. Calcd for $\text{C}_{34}\text{H}_{27}\text{PS}_3$: C, 72.56; H, 4.84%.

General Procedure for the Generation and Reaction of 1 and 3 with Dienophiles. To a solution of **2a–c** (2 mmol) in dry benzene (20 ml) was added the dienophile (6 mmol). The mixture was stirred and refluxed under a nitrogen atmosphere. After the solvent was evaporated, the residue was chromatographed on silica gel (Wakogel C-200) with ethyl acetate–hexane (1:10–1:45) to give the each product. Benzene–hexane (1:1 for **6b**, 1:5 for **6e**) was also used as an eluent.

3-Methoxycarbonyl-3-methyl-4,8-diphenyl-9-thia-1-phosphabicyclo[4.3.0]nona-5,7-diene 1-Sulfide (6a):⁴⁾ Colorless crystals; mp $158\text{--}159^\circ\text{C}$. IR (KBr) 1722 cm^{-1} (C=O); ^1H NMR $\delta=1.60$ (s, 3H), 2.80 (dd, H-2, $J_{2,2'}=13.74$, $J_{\text{HP}}=15.75$ Hz), 3.29 (dd, H-2', $J_{2,2'}=13.74$, $J_{\text{HP}}=13.55$ Hz), 3.45 (s, 3H), 4.48 (dd, H-4, $J_{4,5}=4.94$, $J_{\text{HP}}=4.76$ Hz), 6.87 (d, H-7, $J_{\text{HP}}=31.50$ Hz), 6.88 (dd, H-5, $J_{4,5}=4.94$, $J_{\text{HP}}=38.46$ Hz), $7.23\text{--}7.57$ (m, 10H); ^{13}C NMR (DEPT) $\delta=25.6$ (CH_3), 47.1 (d- CH_2 , C-2, $J_{\text{CP}}=41.5$ Hz), 47.5 (C, C-3), 49.8 (d-CH, C-4, $J_{\text{CP}}=20.8$ Hz), 52.1 (CH_3), 118.1 (d-CH, C-7, $J_{\text{CP}}=15.9$ Hz), 134.5 (d-C, C-8, $J_{\text{CP}}=2.4$ Hz), 140.1 (d-CH, C-5, $J_{\text{CP}}=4.9$ Hz), 143.4 (d-C, C-6, $J_{\text{CP}}=62.3$ Hz), 174.5 (d-C, C=O, $J_{\text{CP}}=13.5$ Hz). MS m/z 412 (M^+ ; 19), 312 ($\text{M}^+\text{--Methyl methacrylate}$; 100), 280 (312-S; 100), 279 (280-H; 80). Found: C, 64.35; H, 5.10%. Calcd for $\text{C}_{22}\text{H}_{21}\text{O}_2\text{PS}_2$: C, 64.06; H, 5.13%.

(6d): Colorless needles; mp $168\text{--}169^\circ\text{C}$. IR (KBr) 2240 cm^{-1} (C $\equiv$ N); ^1H NMR $\delta=1.43$ (d, 3H, $J_{\text{HP}}=2.57$ Hz), 3.08 (dd, H-2, $J_{2,2'}=15.02$, $J_{\text{HP}}=15.02$ Hz), 3.59 (dd, H-2', $J_{2,2'}=15.02$, $J_{\text{HP}}=12.82$ Hz), 4.32 (dd, H-4, $J_{4,5}=4.58$, $J_{\text{HP}}=4.58$ Hz), 6.90 (d, H-7, $J_{\text{HP}}=31.14$ Hz), 6.95 (dd, H-5, $J_{4,5}=4.58$, $J_{\text{HP}}=38.47$ Hz), $7.39\text{--}7.59$ (m, 10H); ^{13}C NMR $\delta=27.4$ (dq, $J_{\text{CP}}=6.1$ Hz), 40.6 (d, C-3, $J_{\text{CP}}=6.1$ Hz), 48.7 (dd, C-4, $J_{\text{CP}}=18.3$ Hz), 49.3 (dt, C-2, $J_{\text{CP}}=47.6$ Hz), 117.1 (dd, C-7, $J_{\text{CP}}=15.9$ Hz), 122.3 (d, C $\equiv$ N, $J_{\text{CP}}=4.9$ Hz), 134.1 (d, C-8, $J_{\text{CP}}=2.4$ Hz), 140.2 (dd, C-5, $J_{\text{CP}}=4.9$ Hz), 144.8 (d, C-6, $J_{\text{CP}}=65.9$ Hz). MS m/z 379 (M^+ ; 10), 312 ($\text{M}^+\text{--Methacrylonitrile}$; 100), 280 (312-S; 93), 67 (Methacrylonitrile; 9). Found: C, 66.23; H, 4.70%. Calcd for $\text{C}_{21}\text{H}_{18}\text{NPS}_2$: C, 66.47; H, 4.78%.

(6e): Colorless Crystals; mp $158\text{--}159^\circ\text{C}$. MS m/z 312 ($\text{M}^+\text{--}\alpha\text{-Methylstyrene}$; 0.4), 280 (312-S; 95), 118 ($\alpha\text{-Methylstyrene}$; 100). Found: C, 72.66; H, 5.16%. Calcd for $\text{C}_{26}\text{H}_{23}\text{PS}_2$: C, 72.52; H, 5.38%. ^1H NMR (major adduct) $\delta=1.77$ (s, 3H), 3.35 (dd, H-2, $J_{2,2'}=14.84$, $J_{\text{HP}}=16.85$ Hz), 3.44 (dd, H-2', $J_{2,2'}=14.84$, $J_{\text{HP}}=12.49$ Hz), 4.77 (dd, H-4, $J_{4,5}=5.31$, $J_{\text{HP}}=5.31$ Hz), 6.68 (d, H-7, $J_{\text{HP}}=58.98$ Hz), 6.80 (dd, H-5, $J_{4,5}=5.31$, $J_{\text{HP}}=38.47$ Hz), $6.88\text{--}7.63$ (m, 15H).

5,9-Diphenyl-4-thia-3-phosphatetetracyclo[9.2.1.0^{2,10}.0^{3,7}]-tetradeca-5,7-diene 3-Sulfide (6f): Colorless crystals; mp $142\text{--}143^\circ\text{C}$. ^1H NMR $\delta=0.59\text{--}3.00$ (m, 9H), 3.15 (dd, H-2, $J_{2,10}=8.41$, $J_{\text{HP}}=14.02$ Hz), 4.71 (ddd, H-9, $J_{8,9}=5.28$, $J_{9,10}=5.28$, $J_{\text{HP}}=5.28$ Hz), 6.80 (d, H-6, $J_{\text{HP}}=31.01$ Hz),

7.05 (dd, H-8, $J_{8,9}=5.28$, $J_{HP}=35.63$ Hz), 7.16—7.61 (m, 10H); ^{13}C NMR (DEPT) $\delta=35.3$ (CH_2 , C-14), 42.7 (d-CH, C-9, $J_{CP}=26.8$ Hz), 51.6 (d-CH, C-10, $J_{CP}=7.3$ Hz), 55.3 (d-CH, C-2, $J_{CP}=40.3$ Hz), 117.3 (d-CH, C-6, $J_{CP}=14.6$ Hz), 140.9 (d-CH, C-8, $J_{CP}=3.7$ Hz), 144.8 (d-C, C-5, $J_{CP}=4.5$ Hz), 144.5 (d-C, C-7, $J_{CP}=58.6$ Hz); ^{31}P NMR $\delta=71.7$. MS m/z 406 (M^+ ; 67), 373 ($\text{M}^+-\text{S}-\text{H}$; 100), 312 ($\text{M}^+-\text{Norbornene}$; 29), 279 (312-S-H; 25). Found: C, 70.77; H, 5.52%. Calcd for $\text{C}_{24}\text{H}_{23}\text{PS}_2$: C, 70.90; H, 5.70%.

5,9-Diphenyl-4-thia-3-phosphatetracyclo[9.2.1.0^{2,10}.0^{3,7}]-tetradeca-6,8-diene 3-Sulfide (7f): Colorless crystals; mp 180—182 °C. ^1H NMR $\delta=1.22$ —2.78 (m, 8H), 2.86 (dd, H-2, $J_{2,10}=8.43$, $J_{HP}=16.12$ Hz), 3.52 (dd, H-10, $J_{2,10}=8.43$, $J_{HP}=12.82$ Hz), 5.95 (m, H-5), 6.39 (broad-d, H-6, $J_{HP}=43.00$ Hz), 6.73 (d, H-8, $J_{HP}=17.22$ Hz), 7.28—7.48 (m, 10H); ^{13}C NMR (DEPT) $\delta=34.8$ (CH_2 , C-14), 51.6 (d-CH, C-10, $J_{CP}=6.1$ Hz), 55.9 (d-CH, C-2, $J_{CP}=43.9$ Hz), 60.3 (d-CH, C-5, $J_{CP}=4.9$ Hz), 118.7 (d-CH, C-8, $J_{CP}=6.1$ Hz), 136.0 (d-CH, C-6, $J_{CP}=13.5$ Hz), 138.4 (d-C, C-7, $J_{CP}=61.1$ Hz), 149.0 (d-C, C-9, $J_{CP}=18.3$ Hz); ^{31}P NMR $\delta=84.3$. MS m/z 406 (M^+ ; 22), 312 ($\text{M}^+-\text{Norbornene}$; 100). Found: C, 71.00; H, 5.81%. Calcd for $\text{C}_{24}\text{H}_{23}\text{PS}_2$: C, 70.90; H, 5.70%.

3-Methoxycarbonyl-4,8-diphenyl-9-thia-1-phosphabicyclo[4.3.0]nona-5,7-diene 1-Sulfide (6g) and 3-Methoxycarbonyl-4,8-diphenyl-9-thia-1-phosphabicyclo[4.3.0]nona-4,6-diene 1-Sulfide (7g): Colorless crystals; mp 177—178 °C. IR (KBr) 1738 cm^{-1} . MS m/z 398 (M^+ ; 100), 366 (M^+-S , 96), 312 ($\text{M}^+-\text{Methyl acrylate}$; 35), 280 (312-S, 59). Found: C, 63.04; H, 4.78%. Calcd for $\text{C}_{21}\text{H}_{19}\text{O}_2\text{PS}_2$: C, 63.30; H, 4.81%. ^1H NMR (6g) $\delta=2.85$ —2.93 (m, H-2'), 3.52 (s, 3H), 4.10—4.17 (m, H-3), 4.47 (ddd, H-4, $J_{4,5}=4.40$, $J_{3,4}=6.23$, $J_{HP}=4.40$ Hz), 6.38 (dd, H-5, $J_{4,5}=4.40$, $J_{HP}=39.19$ Hz), 6.92 (d, H-7, $J_{HP}=26.37$ Hz), 7.08—7.62 (m, 10H); (7g) $\delta=3.01$ (ddd, H-2, $J_{2,3}=9.16$, $J_{2,2'}=13.19$, $J_{HP}=11.72$ Hz), 3.10 (ddd, H-2', $J_{2,3}=4.76$, $J_{2,2'}=13.19$, $J_{HP}=16.48$ Hz), 3.50 (s, 3H), 4.64—4.69 (m, H-3), 6.04 (m, H-8), 6.46 (broad-d, H-7, $J_{HP}=43.95$ Hz), 6.72 (d, H-5, $J_{HP}=21.61$ Hz), 7.08—7.62 (m, 10H).

(6h) and (7h): Colorless crystals; mp 172—173 °C; MS m/z 416 (M^+ ; 67), 312 ($\text{M}^+-\text{Styrene}$; 28), 280 (312-S, 88). Found: C, 72.13; H, 5.15%. Calcd for $\text{C}_{25}\text{H}_{21}\text{PS}_2$: C, 72.09; H, 5.08%. ^1H NMR (6h) $\delta=2.78$ (ddd, H-2, $J_{2,3}=2.20$, $J_{2,2'}=13.19$, $J_{HP}=17.21$ Hz), 3.09—3.16 (m, H-2'), 4.19 (ddd, H-4, $J_{3,4}=4.58$, $J_{4,5}=4.21$, $J_{HP}=4.40$ Hz), 4.44 (dddd, H-3, $J_{2,3}=2.20$, $J_{3,4}=4.58$, $J_{2,2'}=13.19$, $J_{HP}=15.38$ Hz), 6.52 (dd, H-5, $J_{4,5}=4.21$, $J_{HP}=39.92$ Hz), 7.00 (d, H-7, $J_{HP}=26.00$ Hz), 7.06—7.67 (m, 15H); (7h) $\delta=2.86$ (ddd, H-2, $J_{2,2'}=13.55$, $J_{2,3}=13.55$, $J_{HP}=11.35$ Hz), 3.09—3.16 (m, H-2'), 4.88—4.91 (m, H-3), 6.06 (m, H-8), 6.48 (broad-d, H-7, $J_{HP}=43.59$ Hz), 6.80 (d, H-5, $J_{HP}=21.98$ Hz), 7.06—7.67 (m, 15H).

(6i) and (7i): Colorless crystals; mp 176—177 °C. IR (KBr) 1718 cm^{-1} (C=O); MS m/z 382 (M^+ ; 48), 312 ($\text{M}^+-\text{Methyl vinyl ketone}$; 23), 280 (312-S; 19), 43 ($\text{CH}_3\text{C}=\text{O}^+$; 100). Found: C, 65.98; H, 5.19%. Calcd for $\text{C}_{21}\text{H}_{19}\text{OPS}_2$: C, 65.94; H, 5.01%. ^1H NMR (6i) $\delta=2.11$ (s, 3H), 2.86—2.93 (m, H-2, H-2'), 4.07—4.13 (m, H-3), 4.54 (ddd, H-4, $J_{3,4}=4.46$, $J_{4,5}=4.46$, $J_{HP}=6.56$ Hz), 6.39 (dd, H-5, $J_{4,5}=4.46$, $J_{HP}=39.02$ Hz), 6.89 (d, H-7, $J_{HP}=27.00$ Hz), 7.09—7.61 (m, 10H); (7i) $\delta=2.05$ (s, 3H), 2.81 (ddd, H-2, $J_{2,3}=11.54$, $J_{2,2'}=12.82$, $J_{HP}=9.52$ Hz), 3.00 (ddd, H-2', $J_{2,3}=4.95$, $J_{2,2'}=12.82$, $J_{HP}=16.48$ Hz), 4.74 (ddd, H-

3, $J_{2,3}=4.95$, $J_{2,3}=11.54$, $J_{HP}=5.12$ Hz), 6.04 (m, H-8), 6.48 (d, H-7, $J_{HP}=45.06$ Hz), 6.76 (d, H-5, $J_{HP}=21.25$ Hz), 7.09—7.61 (m, 10H).

2,3-Bis(methoxycarbonyl)-6-phenyl-4-styryl-4H-thiin (8j): Yellow oil; IR (neat) 1734 cm^{-1} (C=O); ^1H NMR $\delta=3.77$ (s, 3H), 3.85 (s, 3H), 4.46 (ddd, H-4, $J_{4,5}=6.7$, $J_{4,1'}=7.3$, $J_{4,2'}=0.5$ Hz), 6.05 (d, H-5, $J_{4,5}=6.7$ Hz), 6.15 (dd, H-1', $J_{4,1'}=7.3$, $J_{1',2'}=16.0$ Hz), 6.52 (dd, H-2', $J_{4,2'}=0.5$, $J_{1',2'}=16.0$ Hz), 7.01—7.57 (m, 10H); ^{13}C NMR $\delta=41.8$ (d, C-4), 52.6 (q), 53.2 (q), 118.9 (d, C-5), 124.9 (d, C-2'), 128.0 (s, C-6), 131.6 (d, C-1'), 133.2 (s, C-3), 134.8 (s, C-2), 165.1 (s, C=O), 165.8 (s, C=O); MS m/z 392 (M^+ ; 14), 360 (M^+-S , 45), 333 ($\text{M}^+-\text{CO}_2\text{CH}_3$, 100), 301 (333-S, 60). Found: m/z 392.1085. Calcd for $\text{C}_{23}\text{H}_{20}\text{O}_4\text{S}$; M, 392.1083.

2,3-Bis(ethoxycarbonyl)-6-phenyl-4-styryl-3,4-dihydro-2H-1,2,3-thiadiazine (8k): Orange oil. IR (neat) 1728 cm^{-1} (C=O); ^1H NMR $\delta=1.14$ (t, 3H, $J_{HH}=7.0$ Hz), 1.30 (t, 3H, $J_{HH}=7.0$ Hz), 4.12 (q, 2H, $J_{HH}=7.0$ Hz), 4.26 (q, 2H, $J_{HH}=7.0$ Hz), 5.60 (m, H-4), 6.00 (d, H-5, $J_{4,5}=5.0$ Hz), 6.27 (dd, H-1', $J_{4,1'}=5.5$, $J_{1',2'}=16.0$ Hz), 6.58 (d, H-2', $J_{1',2'}=16.0$ Hz), 7.10—7.54 (m, 10H); ^{13}C NMR $\delta=14.3$ (q, CH_3), 14.5 (q, CH_3), 55.1 (d, C-4), 63.1 (t, CH_2), 63.9 (t, CH_2), 154.6 (s, C=O), 155.0 (s, C=O); MS m/z 424 (M^+ ; 9), 250 (Thioketone; 92), 29 (CH_2CH_3 ; 100). Found: m/z 424.1449. Calcd for $\text{C}_{23}\text{H}_{24}\text{O}_4\text{N}_2\text{S}$; M, 424.1458.

8-(p-Methoxyphenyl)-5-styryl-2-thia-8-azabicyclo[4.3.0]non-3-ene-7,9-dione (8l): Colorless needles; mp 185—187 °C. IR (KBr) 1712 cm^{-1} (C=O); ^1H NMR $\delta=3.56$ (ddd, H-5, $J_{4,5}=5.13$, $J_{5,6}=5.13$, $J_{5,1'}=9.16$ Hz), 3.77 (s, 3H), 3.82 (dd, H-6, $J_{5,6}=5.13$, $J_{1,6}=9.52$ Hz), 4.34 (d, H-1, $J_{1,6}=9.52$ Hz), 4.49 (d, H-4, $J_{4,5}=5.13$ Hz), 6.69 (d, H-2', $J_{1',2'}=15.94$ Hz), 6.86—6.89 (m, 2H), 6.95 (dd, H-1', $J_{5,1'}=9.16$, $J_{1',2'}=15.94$ Hz), 6.98—7.01 (m, 2H), 7.24—7.60 (m, 10H); ^{13}C NMR (DEPT) $\delta=44.8$ (CH, C-6), 45.1 (CH, C-1), 50.9 (CH, C-5), 55.4 (CH_3), 128.8 (CH, C-4), 128.9 (CH, C-2'), 132.9 (CH, C-1'), 174.9 (C, C=O), 175.2 (C, C=O); MS m/z 453 (M^+ ; 33), 250 (Thioketone; 100), 203 (MI; 52). Found: C, 73.90; H, 5.05%. Calcd for $\text{C}_{28}\text{H}_{23}\text{O}_3\text{NS}$; C, 74.15; H, 5.11%.

The authors wish to express their thanks to Dr. Shū Kobayashi for measuring ^{31}P NMR spectra.

References

- 1) O. Tsuge, E. Wada, and S. Kanemasa, *Chem. Lett.*, **1983**, 239; O. Tsuge, S. Kanemasa, E. Wada, and H. Sakoh, *Yuki Gosei Kagaku Kyokai Shi*, **44**, 756 (1986).
- 2) S. Motoki, Y. Terauchi, and S. Kametani, *Chem. Lett.*, **1988**, 717; S. Motoki, Y. Matsuo, and Y. Terauchi, *Bull. Chem. Soc. Jpn.*, **63**, 284 (1990).
- 3) In the adduct **6**, stereoisomers with respect to the phenyl group at position 4 and the substituent at position 3 are possible depending on the *endo* and *exo* addition of the dienophiles to the thiaphosphole **1**. However, no stereoisomers were obtained in detectable amount except **6e** (inseparable mixture) and the ^1H NMR spectra showed that **6g**—**i** were 3,4-*cis* (*endo*) isomers.
- 4) (**2b**): Mp 152—153 °C, (**2c**): Mp 152—153 °C, (**6b**): Yellow oil, (**6c**): Mp 151—152 °C. Elementary analyses, IR, NMR, and mass spectra of these compounds gave satisfactory results.