

Preparation of New Optically Active Germylmethylamine Derivatives

Daiyo TERUNUMA,^{*,†} Hiroaki KIZAKI, Takashi SATO, Koujiro MASUO, and Hiroyuki NOHIRA

Department of Applied Chemistry, Faculty of Engineering, Saitama University, Shimo-ohkubo 255, Urawa, Saitama 338

(Received September 2, 1992)

Synopsis. New optically active organogermanium compounds, such as [methylphenyl(*o*-tolyl)-, benzylmethylphenyl-, benzylmethyl(*o*-tolyl)-, and benzyl(*o*-methoxyphenyl)-methylgermyl]methylamine, were prepared in high optical purity.

A few optically active organogermanium compounds have been reported.¹⁾ One of the problems in the preparation of asymmetric germanium compound is a difficulty to introduce different substituents on germanium. In the course of our interest concerning the preparation of optically active organometallic compounds,²⁾ new optically active organogermanium compounds were successfully obtained. Trichloro(chloromethyl)germane, prepared from the reaction of tetrachlorogermane with diazomethane, was treated stepwisely with Grignard reagents of aryl, benzyl, and methyl halides to give the corresponding (chloromethyl) germanium derivatives (**1a**, **1b**, **1c**, **1d**). Racemic organogermanium compounds, such as [methylphenyl(*o*-tolyl)germyl]methylamine (**2a**), (benzylmethyl phenylgermyl)methylamine (**2b**), [benzylmethyl(*o*-tolyl)germyl]methylamine (**2c**), and [benzyl(*o*-methoxyphenyl)methylgermyl]methylamine (**2d**), were prepared by treating the corresponding **1a**, **1b**, **1c** and **1d** with ammonia in an autoclave at 100°C (Scheme 1). The optical resolution of **2a** was carried out using 2-phenylpropanoic acid (PPA) as a resolving agent; **2b**, **2c**,

and **2d** were resolved using tartaric acid as a resolving agent in a similar manner as that described for the preparation of their silicon analogues.^{3,4)} The optical purities of these amines were determined by measuring the NMR spectra of the amides of the optically active **2a**, **2b**, and **2c** with Mosher agent (MTPA)⁵⁾ (Table 1). In the case of the amide of racemic **2d** with MTPA, no diastereotopic separation of the signals was observed in its NMR spectrum, and optical purity of **2d** was determined by NMR spectrum using (benzylmethylphenylsilyl)acetic acid as an optical purity-determining reagent.²⁾


Experimental

The melting points were uncorrected. The NMR spectra were recorded on a JEOL 60Si (60MHz), and a Bruker AM400 (400MHz) spectrometer in CDCl₃ using TMS as an internal standard. The optical rotations were measured with a JASCO DIP-360 polarimeter. All of the Grignard reactions were carried out under a dry-nitrogen atmosphere. Trichloro(chloromethyl)germane was prepared by the procedure described in the literature.⁶⁾

Dichloro(chloromethyl)phenylgermane. Phenylmagnesium bromide prepared from bromobenzene (17.7 g, 113 mmol) and magnesium (3.0 g, 124 mmol) in dry ether (120 ml) was added to a solution of trichloro(chloromethyl)germane (25.8 g, 113 mmol) in dry ether (120 ml) at 0°C with stirring. The reaction mixture was stirred for 1 h at 0°C and then refluxed for 1 h. After the addition of hexane (80 ml), the reaction mixture was kept at room temperature over night. Filtration, evaporation and distillation of the residue gave a colorless liquid. Yield 21.3 g (70%). Bp 140°C/12 Torr (1 Torr=133.322 Pa).

Dichloro(chloromethyl)(*o*-tolyl)germane and dichloro(chloromethyl)(*o*-methoxyphenyl)germane were prepared in the similar manner described above in 90% (bp 118°C/0.18 Torr) and 60% (bp 138°C/0.11 Torr) yield, respectively.

(Chloromethyl)methylphenyl(*o*-tolyl)germane (1a). The *o*-tolylmagnesium bromide prepared from *o*-bromotoluene (13.5 g, 79 mmol) and magnesium (2.1 g, 87 mmol) in dry ether (80 ml) was added to a mixture of dichloro(chloromethyl)phenylgermane (21.3 g, 79 mmol) in dry ether (50 ml) at 0°C with stirring. The mixture was stirred at room



Scheme 1.

[†]Present address: Department of Functional Materials Science and Engineering, Faculty of Engineering, Saitama University, Shimo-ohkubo 255, Urawa, Saitama 338.

Table 1. Optically Active Organogermanium Compounds

	Yield %	Specific rotation °(c 2, CH ₂ Cl ₂)	Optical purity % e.e.
2a	14	-7.1	99
2b	14	+10.0	99
2c	30	+6.0	98
2d	17	+21.1	97

temperature for 3 h. A Grignard reagent prepared from iodomethane (13.3 g, 87 mmol) and magnesium (2.3 g, 95 mmol) in dry ether (70 ml) was added to the reaction mixture at 0°C with stirring. The mixture was stirred at room temperature for 3 h, and then refluxed for 1 h. An aqueous HCl solution (3 M, 40 ml, 1 M = 1 mol dm⁻³) was added to the mixture at 0°C. The organic layer was separated and the aqueous layer were extracted with ether (30 ml × 2). The combined organic layer and extracts were dried (Na₂SO₄). Evaporation and distillation of the residue gave a colorless liquid. Yield 17.2 g (71%). Bp 141°C/0.18 Torr. ¹H NMR (CDCl₃) δ = 0.80 (3H, s, Ge-CH₃), 2.28 (3H, s, Ar-CH₃), 3.38 (2H, s, CH₂Cl), 7.0–7.6 (9H, m, C₆H₅, C₆H₄).

The compound **1b**, **1c**, and **1d** were prepared in the similar manner described above. **1b**: Yield 62%. Bp 146°C/0.18 Torr. ¹H NMR (CDCl₃) δ = 0.40 (3H, s, Ge-CH₃), 2.55 (2H, s, Ge-CH₂-Ar), 3.00 (2H, s, Ge-CH₂Cl), 6.6–7.6 (10H, m, Ph). **1c**: Yield 31%. Bp 128°C/0.15 Torr. ¹H NMR (CDCl₃) δ = 0.48 (3H, s, Ge-CH₃), 2.10 (3H, s, CH₃-Ar), 2.62 (2H, s, Ge-CH₂-Ar), 3.10 (2H, s, Ge-CH₂Cl), 6.6–7.6 (9H, m, C₆H₅, C₆H₄). **1d**: Yield 59%. Bp 140°C/0.14 Torr. ¹H NMR (CDCl₃) δ = 0.45 (3H, s, Ge-CH₃), 2.65 (2H, s, Ge-CH₂-Ar), 3.17 (2H, s, Ge-CH₂Cl), 3.79 (3H, s, OCH₃), 6.6–6.8 (9H, m, C₆H₅, C₆H₄).

Methylphenyl(*o*-tolyl)germylmethylamine (2a). In a stainless-steel autoclave, **1a** (17.2 g, 56 mmol) and liquid ammonia (96 g, 5.6 mol) were added under a nitrogen atmosphere. The autoclave was kept for 1.5 h at 100°C. The excess ammonia was liberated from the autoclave, and the contents were washed out with an aqueous alkali (6 M NaOH, 50 ml) and then benzene (100 ml). The organic layer was separated and the aqueous layer was extracted with ether (30 ml × 3). The combined organic layer and extracts were dried (Na₂SO₄). Evaporation and distillation gave a colorless liquid. Yield 13.0 g (80%). Bp 142°C/0.18 Torr. ¹H NMR (CDCl₃) δ = 0.75 (3H, s, Ge-CH₃), 1.10 (2H, s, NH₂), 2.30 (3H, s, Ar-CH₃), 2.95 (2H, s, Ge-CH₂N), 7.0–7.6 (9H, m, C₆H₅, C₆H₄).

The compound **2b**, **2c**, and **2d** were prepared in the similar manner described above. **2b**: Yield 84%. Bp 138°C/0.11 Torr. ¹H NMR (CDCl₃) δ = 0.31 (3H, s, Ge-CH₃), 0.80 (2H, s, NH₂), 2.50 (2H, s, Ge-CH₂-Ar), 2.61 (2H, s, Ge-CH₂-N), 6.7–7.7 (10H, m, Ph). **2c**: Yield 73%. Bp 149°C/0.3 Torr. ¹H NMR (CDCl₃) δ = 0.36 (3H, s, Ge-CH₃), 0.80 (2H, s, NH₂), 2.35 (3H, s, Ar-CH₃), 2.53 (2H, s, Ge-CH₂-Ar), 2.63 (2H, s, Ge-CH₂N), 6.6–7.6 (9H, m, C₆H₅, C₆H₄). **2d**:

Yield 68%. Bp 150°C/0.15 Torr. ¹H NMR (CDCl₃) δ = 0.25 (3H, s, Ge-CH₃), 0.75 (2H, s, NH₂), 2.50 (2H, s, Ge-CH₂-Ar), 2.56 (2H, s, Ge-CH₂N), 3.75 (3H, s, OCH₃), 6.5–7.5 (9H, m, C₆H₅, C₆H₄).

Optical Resolution of 2a. A solution of **2a** (10.0 g, 35 mmol) and (+)-PPA (5.25 g, 35 mmol) in benzene/hexane (1/1) mixture (122 ml) was kept overnight at room temperature. The crystalline precipitate was separated by filtration and recrystallized from benzene 10 times. Yield 2.5 g (16%). Mp 141–142°C. [α]_D +2.7° (c 1, MeOH). Found: C, 65.82; H, 6.65; N, 3.01%. Calcd for C₂₄H₂₉NGeO₂: C, 66.10; H, 6.70; N, 3.21%.

An aqueous solution of 10% NaOH was added to the crystals at 0°C. The liberated amine was extracted three times with ether (20 ml × 3). The combined extracts were dried (Na₂SO₄). Evaporation and distillation gave a colorless oil. Yield 1.3 g (14%). [α]_D -7.1° (c 2, CH₂Cl₂) (99% e.e.). The (+)-rich **2a** recovered from the mother liquor was treated with an equimolar amount of (-)-PPA in the similar manner described above to give white crystals. Yield 3.6 g (24%). Mp 140–141°C. [α]_D -2.7° (c 1, MeOH).

Liberation gave a colorless oil. Yield 2.0 g (20%). [α]_D +6.9° (c 2, CH₂Cl₂) (97% e.e.). The optical resolutions of **2b**, **2c**, and **2d** were carried out in the similar manner to that described in the literature²⁾ using (+)-tartaric acid as a resolving reagent. **2b**: Yield 14%. [α]_D +10.0° (c 2, CH₂Cl₂) (99% e.e.). **2c**: Yield 30%. [α]_D +6.0° (c 2, CH₂Cl₂) (98% e.e.). **2d**: Yield 17%. [α]_D +21.1° (c 2, CH₂Cl₂) (97% e.e.).

References

- 1) A. G. Brook and G. J. D. Peddle, *J. Am. Chem. Soc.*, **85**, 1869 (1963).
- 2) D. Terunuma, M. Kato, M. Kamei, H. Uchida, S. Ueno, and H. Nohira, *Bull. Chem. Soc. Jpn.*, **59**, 3581 (1986).
- 3) D. Terunuma, K. Murakami, M. Kokubo, K. Senda, and H. Nohira, *Bull. Chem. Soc. Jpn.*, **53**, 789 (1980).
- 4) D. Terunuma, N. Yamamoto, H. Kizaki, and H. Nohira, *Nippon Kagaku Kaishi*, **1990**, 451.
- 5) J. A. Dale and H. S. Mosher, *J. Am. Chem. Soc.*, **95**, 512 (1973).
- 6) D. Seyferth and E. G. Rochow, *J. Am. Chem. Soc.*, **77**, 907 (1955).