

Free Radical Reaction of 1,3-Dimethyluracils and Caffeine with Benzoyl Peroxide in γ -Butyrolactone

Toshio ITAHARA* and Naoko IDE

Institute of Chemistry, College of Liberal Arts, Kagoshima University, Korimoto, Kagoshima 890

(Received March 5, 1993)

Synopsis. Treatment of 1,3-dimethyluracil, 1,3-dimethylthymine, and caffeine with benzoyl peroxide in γ -butyrolactone at 80 °C resulted in a free radical alkylation.

Recently we reported that the treatment of 1,3-dimethyluracils (**1**) and caffeine (**2a**) with benzoyl peroxide in cyclohexane, acetonitrile, and alkyl acetates led to a free radical alkylation of **1** and **2a** with alkyl radicals formed by abstraction of hydrogen atom from the solvents.¹⁾ The previous work has also shown that alkyl acetates were favorable solvents for the free radical alkylation of **1** and **2a**, compared with the corresponding ethers and ketones, although the abstraction of hydrogen atom from ethers by the radicals derived from benzoyl peroxide is well-known.²⁾ Furthermore, several types of natural products³⁾ containing γ -butyrolactone ring such as canescin⁴⁾ and conocarpin⁵⁾ are known. These observations led us to examine a reaction of 1,3-dimethyluracil (**1a**), 1,3-dimethylthymine (**1b**), and **2a** with benzoyl peroxide in γ -butyrolactone, compared with that in tetrahydrofuran and cyclopentanone.

A solution of **1a** in γ -butyrolactone in the presence of benzoyl peroxide was heated at 80 °C under argon atmosphere. The reaction mixture was submitted to chromatography on silica gel. By elution of a mixture of hexane and ethyl acetate, two main products **1c** and **1d** were isolated together with a small amount of **1e**. Under similar conditions, the reaction of **1b** gave **1f** and **1g** and that of **2a** gave **2c** and **2d**. The results are summarized in Table 1. In order to compare with the reaction in γ -butyrolactone, treatment of **1a** and **2a** with benzoyl peroxide in tetrahydrofuran and cyclopentanone was further investigated. The reaction of **1a** and **2a** in tetrahydrofuran gave **1h** and **2h**, respectively, but almost no alkylation occurred in the case of the treatment of **1a** and **2a** in cyclopentanone (Chart 1). The results suggest that not only alkyl acetates¹⁾ but also γ -butyrolactone may be favorable solvents on free radical alkylation with benzoyl peroxide.

Experimental

Melting points were determined on a Yanagimoto micro melting point apparatus and uncorrected. The ¹H NMR (400 MHz) and ¹³C NMR (100 MHz) spectra were obtained with a JEOL GSX400 spectrometer using tetramethylsilane as an internal standard in CDCl₃. Mass spectra were obtained with a JEOL JMS-D300 spectrometer. The elemental analyses were performed by the Analytical Center of Kyoto University.

General Procedure for the Reaction of 1,3-Di-

methyluracil (**1a**), 1,3-Dimethylthymine (**1b**), and Caffeine (**2a**) with Benzoyl Peroxide. A solution of **1a**, **1b**, and **2a** (1.5 mmol) and benzoyl peroxide (3 mmol) in γ -butyrolactone (50 ml) or tetrahydrofuran (200 ml) was heated at 80 °C or at reflux temperature (in tetrahydrofuran) under argon atmosphere. The reaction mixture was evaporated under reduced pressure to give a residue, which was then chromatographed on silica gel (Fuji gel packed column NQ-2, 24 mm ϕ $\times$ 360 mm), eluted with a mixture of hexane and ethyl acetate, with low-pressure pump (Chemco Low-prep pump 81-M-2R) and UV-monitor (Shimadzu UV-detector SPC-6A, detected at 250 nm), to give the products and recovered substrate. The results are shown in Table 1. The spectral data of the products are given below.

5-(2-Oxotetrahydro-3-furyl)-1,3-dimethyluracil (1c): Mp 157–159 °C; ¹H NMR δ =7.26 (s, 1H, 6-H), 4.51 (ddd, 1H, J =8.8, 8.8, 2.5 Hz, 5'-H), 4.32 (ddd, 1H, J =8.8, 8.8, 6.8 Hz, 5'-H), 3.59 (t, 1H, J =10.0, 3'-H), 3.41 (s, 3H, Me), 3.35 (s, 3H, Me), 2.64 (m, 1H, 4'-H), 2.34 (m, 1H, 4'-H); ¹³C (¹³C-¹H COSY) NMR δ =176.84, 162.55, 151.43, 141.93 (δ 7.26), 109.29, 66.95 (δ 4.59 and 4.32), 39.60 (δ 3.59), 37.08 (δ 3.41), 28.77 (δ 2.64 and 2.34), 27.94 (δ 3.35); MS m/z (rel intensity, %) 225 (M^+ +1, 13), 224 (M^+ , 75), 180 (85), 179 (21), 166 (29), 165 (61), 94 (100). Calcd for C₁₀H₁₂N₂O₄:




Chart 1.

Table 1. Free Radical Reaction of 1,3-Dimethyluracils and Caffeine with Benzoyl Peroxide^{a)}

Substrate	Solvent	(ml)	Conv. %	Product /Isolated yield % ^{b)}
1a	γ -Butyrolactone	(50)	40	1c /56; 1d /20; 1e /2
1b	γ -Butyrolactone	(50)	38	1f /12; 1g /5
2a	γ -Butyrolactone	(50)	52	2c /52; 2d /24
1a	Tetrahydrofuran	(200)	15	1h /88
2a	Tetrahydrofuran	(200)	40	2h /85

a) Reaction conditions: substrate (1.5 mmol), benzoyl peroxide (3 mmol), 7h reaction at 80 °C in γ -butyrolactone and 15 h reaction at reflux temperature in tetrahydrofuran. b) Yield based on substrate consumed.

C, 53.57; H, 5.39; N, 12.50%. Found: C, 53.80; H, 5.37; N, 12.59%.

5-(5-Oxotetrahydro-2-furyl)-1,3-dimethyluracil (1d): Mp 110–111 °C; ¹H NMR δ =7.26 (s, 1H, 6-H), 5.38 (t, 1H, J =7.5 Hz, 2'-H), 3.42 (s, 3H, Me), 3.35 (s, 3H, Me), 2.75 (m, 1H, 3'-H), 2.55–2.68 (m, 2H, 4'-H), 2.11 (ddt, 1H, J =7.5, 12.4, 8.8 Hz, 3'-H); ¹³C (¹³C–¹H COSY) NMR δ =176.59, 161.75, 151.40, 139.27 (δ 7.26), 111.77, 75.96 (δ 5.38), 37.23 (δ 3.42), 28.48 (δ 2.75 and 2.11), 28.27 (δ 2.55–2.68), 27.82 (δ 3.35); MS m/z (rel intensity, %) 225 (M^+ +1, 6), 224 (M^+ , 39), 196 (97), 169 (97), 169 (100), 167 (56), 140 (72). Calcd for C₁₀H₁₂N₂O₄: C, 53.57; H, 5.39; N, 12.50%. Found: C, 53.36; H, 5.28; N, 12.24%.

5-Benzoyloxy-1,3-dimethyluracil (1e): Mp 150.5–152 °C; ¹H NMR δ =8.15 (dd, 2H, J =8.2, 1.0 Hz, Ph), 7.45–7.80 (m, 3H, Ph), 7.34 (s, 1H, 6-H), 3.44 (s, 3H, Me), 3.40 (s, 3H, Me); ¹³C NMR δ =164.53, 158.72, 150.84, 134.72, 134.07, 130.42, 128.64, 128.19, 126.57, 37.15, 28.41; MS m/z (rel intensity, %) 260 (M^+ , 3), 105 (100), 77 (38). Calcd for C₁₃H₁₂N₂O₄: C, 59.99; H, 4.65; N, 10.77%. Found: C, 59.94; H, 4.69; N, 11.06%.

5-[(2-Oxotetrahydro-3-furyl)methyl]-1,3-dimethyluracil (1f): Mp 77–78 °C; ¹H NMR δ =7.21 (s, 1H, 6-H), 4.32 (ddd, 1H, J =8.8, 8.8, 2.2 Hz, 5'-H), 4.17 (ddd, 1H, J =8.8, 8.8, 5.8 Hz, 5'-H), 3.40 (s, 3H, Me), 3.36 (s, 3H, Me), 2.92 (ddt, 1H, J =6.2, 5.9, 8.8 Hz, 3'-H), 2.80 (dd, 1H, J =14.0, 6.2 Hz, 5-CH₂), 2.61 (dd, 1H, J =14.0, 5.9 Hz, 5-CH₂), 2.38 (m, 1H, 4'-H), 2.02 (m, 1H, 4'-H); ¹³C (¹³C–¹H COSY) NMR δ =178.86, 163.84, 151.68, 141.89 (δ 7.21), 109.48, 66.53 (δ 4.32 and 4.17), 38.85 (δ 2.92), 36.99 (δ 3.40), 28.38 (δ 2.02 and 2.38), 28.04 (δ 3.36), 27.39 (δ 2.61 and 2.81); MS m/z (rel intensity, %) 239 (M^+ +1, 5), 238 (M^+ , 31), 192 (20), 165 (11), 154 (30), 153 (100), 140 (12). Calcd for C₁₁H₁₄N₂O₄: C, 55.45; H, 5.92; N, 11.76%. Found: C, 55.26; H, 5.87; N, 11.68%.

8-(2-Oxotetrahydro-3-furyl)caffeine (2c): Mp 211–213 °C; ¹H NMR δ =4.69 (ddd, 1H, J =8.8, 8.8, 2.2 Hz, 5'-H), 4.47 (ddd, 1H, J =8.8, 8.8, 7.6 Hz, 5'-H), 4.09 (s, 3H, Me), 3.99 (t, 1H, J =8.8 Hz, 3'-H), 3.54 (s, 3H, Me), 3.40 (s, 3H, Me), 3.12 (m, 1H, 4'-H), 2.67 (m, 1H, 4'-H); ¹³C (¹³C–¹H COSY) NMR δ =173.26, 155.29, 151.62, 147.63, 147.58, 108.52, 67.85 (δ 4.69 and 4.47), 38.37 (δ 3.99), 32.18 (δ 4.09), 29.72 (δ 3.54), 27.94 (δ 3.40), 27.23 (δ 3.12 and 2.67); MS m/z (rel intensity, %) 279 (M^+ +1, 18), 278 (M^+ , 100), 219 (48), 149 (68). Calcd for C₁₂H₁₄N₄O₄: C, 51.79; H, 5.07; N, 20.14%. Found: C, 51.55; H, 4.89; N, 20.06%.

8-(5-Oxotetrahydro-2-furyl)caffeine (2d): Mp

190–192 °C; ¹H NMR δ =5.61 (dd, 1H, J =6.0 and 7.0 Hz, 2'-H), 4.07 (s, 3H, Me), 3.55 (s, 3H, Me), 3.40 (s, 3H, Me), 3.10 (m, 1H, 3'-H), 2.85 (m, 1H, 4'-H), 2.58–2.70 (m, 2H, 3'-H and 4'-H); ¹³C (¹³C–¹H COSY) NMR δ =175.83, 155.48, 151.57, 148.24, 147.19, 108.98, 71.78 (δ 5.61), 32.26 (δ 4.07), 29.75 (δ 3.55), 28.00 (δ 3.40), 27.90 (δ 2.85 and 2.58–2.70), 25.13 (δ 3.10 and 2.58–2.70); MS m/z (rel intensity, %) 279 (M^+ +1, 16), 278 (M^+ , 100), 234 (16), 223 (22), 219 (27). Calcd for C₁₂H₁₄N₄O₄: C, 51.79; H, 5.07; N, 20.14%. Found: C, 51.79; H, 5.06; N, 20.02%.

5-(Tetrahydro-2-furyl)-1,3-dimethyluracil (1h): Mp 92–93 °C (lit.⁶⁾ 94–95 °C); ¹H NMR δ =7.24 (s, 1H), 4.79 (t, 1H, J =7 Hz), 3.97–4.03 (m, 1H), 3.86 (q, 1H, J =7 Hz), 3.41 (s, 3H), 3.34 (s, 3H), 2.37–2.46 (m, 1H), 1.90–2.00 (m, 1H), 1.63–1.72 (m, 1H); ¹³C NMR δ =162.50, 151.78, 138.25, 115.29, 74.71, 68.44, 37.01, 32.36, 27.72, 25.69.

8-(Tetrahydro-2-furyl)caffeine (2h): Mp 127–128 °C (lit.⁷⁾ 127–128 °C); ¹H NMR δ =5.03 (t, 1H, J =7 Hz), 4.04 (s, 3H), 3.91–3.97 (m, 2H), 3.56 (s, 3H), 3.39 (s, 3H), 2.53–2.63 (m, 1H), 2.23–2.33 (m, 1H), 2.12–2.22 (m, 1H), 2.00–2.10 (m, 1H); ¹³C NMR δ =155.51, 152.40, 151.73, 147.37, 108.37, 72.66, 68.92, 32.26, 29.72, 29.60, 27.88, 25.97.

References

- 1) T. Itahara and N. Ide, *Bull. Chem. Soc. Jpn.*, **65**, 2045 (1992).
- 2) a) W. E. Case, *J. Am. Chem. Soc.*, **69**, 500 (1947); b) P. D. Bartlett and K. Nozaki, *J. Am. Chem. Soc.*, **69**, 2299 (1947); c) D. B. Denny and G. Feig., *J. Am. Chem. Soc.*, **81**, 5322 (1959); d) R. L. Huang, H. H. Lee, and S. H. Ong, *J. Chem. Soc.*, **1962**, 3336.
- 3) "Handbook of Natural Occurring Compounds, Vol. 1, Acetogenins, Shikimates, and Carbohydrates," ed by T. K. Devon and A. I. Scott, Academic Press, New York (1975).
- 4) A. J. Birch, J. J. Wright, F. Gager, L. Mo, and A. Pelter, *Tetrahedron Lett.*, **1969**, 1519.
- 5) P. E. Kruger and G. W. Perold, *J. Chem. Soc. C*, **1970**, 2127.
- 6) U. Hacksell and G. D. Daves, Jr., *J. Org. Chem.*, **48**, 4144 (1983).
- 7) S. Jerumanis and A. Martel, *Can. J. Chem.*, **48**, 1716 (1970).