

Wagner–Meerwein Rearrangement of 4-Hydroxymethylpentacyclo[4.3.0.0^{2,5}.0^{3,8}.0^{4,7}]nonane to Pentacyclo[5.3.0.0^{2,6}.0^{3,9}.0^{4,8}]decane in Formic Acid

Takeshi HASEGAWA, Yoshiyuki KUWATANI, Hiroyuki HIGUCHI,[†] and Ikuo UEDA*

The Institute of Scientific and Industrial Research, Osaka University, Ibaraki, Osaka 567

[†]The Faculty of Science, Toyama University, Gofuku, Toyama 930

(Received May 10, 1993)

The cationic rearrangement of 1-bromo-4-[hydroxybis(4-methoxyphenyl)methyl]pentacyclo[4.3.0.0^{2,5}.0^{3,8}.0^{4,7}]nonane-9-spiro-2'-[1,3]dioxolane (**1**) in formic acid gave 4-bromo-1-formyloxy-10,10-bis(4-methoxyphenyl)pentacyclo[5.3.0.0^{2,6}.0^{3,9}.0^{4,8}]decane-5-spiro-2'-[1,3]dioxolane in 27% yield along with 5-bromo-9-formyloxy-10,10-bis(4-methoxyphenyl)pentacyclo[5.3.0.0^{2,5}.0^{3,9}.0^{4,8}]decane-6-spiro-2'-[1,3]dioxolane in 55% yield. The reaction of **1** with *p*-toluenesulfonic acid or hydrochloric acid in methanol also afforded 4-bromo-1-methoxy-10,10-bis(4-methoxyphenyl)pentacyclo[5.3.0.0^{2,6}.0^{3,9}.0^{4,8}]decane-5-spiro-2'-[1,3]dioxolane together with 5-bromo-9-methoxy-10,10-bis(4-methoxyphenyl)pentacyclo[5.3.0.0^{2,5}.0^{3,9}.0^{4,8}]decane-6-spiro-2'-[1,3]dioxolane. With respect to migration of the C₄–C₅ bond in the pentacyclo[4.3.0.0^{2,5}.0^{3,8}.0^{4,7}]nonane (homocubane) skeleton to pentacyclo[5.3.0.0^{2,6}.0^{3,9}.0^{4,8}]decane (*D*_{2h}-bishmocupane), a rearrangement into the bishomocupane system in driven by a concomitant release of strain; also, the stability of the 4-homocubane-methyl cation intermediate may play an important role in determining the course of the cage expansion into the *D*_{2h}-bishomocupane system. This reaction provides an attractive route for the synthesis of bishomocupane bridgehead alcohols and related compounds which are not easily accessible by other routes.

The principal structures of pentacyclo[4.2.0.0^{2,5}.0^{3,8}.0^{4,7}]octane (cubane), pentacyclo[4.3.0.0^{2,5}.0^{3,8}.0^{4,7}]nonane (homocubane), pentacyclo[5.3.0.0^{2,5}.0^{3,9}.0^{4,8}]decane (*C*₂-bishmocupane), and pentacyclo[5.3.0.0^{2,6}.0^{3,9}.0^{4,8}]decane (*D*_{2h}-bishmocupane) are depicted together with their relative strain energies (SE), obtained by force field calculations (Chart 1).¹⁾ The driving force in a skeletal rearrangement of cubane to homocubane or homocubane to bishomocupane is the relief of about 40 kcal mol⁻¹ of ring strain. In the regiospecific cationic rearrangement of the 4-hydroxymethylpentacyclo[4.3.0.0^{2,5}.0^{3,8}.0^{4,7}]nonane (4-homocubanemethanol)

system, force-field calculations show that migration of the C₃–C₄ bond (or the equivalent C₄–C₇ bond) to C₂-bishmocupane releases more strain energy than the migration of the central C₄–C₅ bond, which would lead *D*_{2h}-bishomocupane.²⁾ These theoretical calculations are supported by experimental results that the cationic rearrangement of 1-bromo-4-(hydroxydiphenyl)methylpentacyclo[4.3.0.0^{2,5}.0^{3,8}.0^{4,7}]nonane-9-spiro-2'-[1,3]dioxolane (**4**) and 1-bromo-4-(hydroxydimethyl)methylpentacyclo[4.3.0.0^{2,5}.0^{3,8}.0^{4,7}]nonane-9-spiro-2'-[1,3]dioxolane (**5**) leads to *C*₂-bishomocupane in good yields without giving *D*_{2h}-bishomocupane.^{2,3)} In this paper we describe a new example of the formation of 4-bromo-1-formyloxy-10,10-bis(4-methoxyphenyl)pentacyclo[5.3.0.0^{2,6}.0^{3,9}.0^{4,8}]decane-5-spiro-2'-[1,3]dioxolane (**2a**) and its analogs (**2b–e**) by a cationic rearrangement of 1-bromo-4-[hydroxybis(4-methoxyphenyl)methyl]pentacyclo[4.3.0.0^{2,5}.0^{3,8}.0^{4,7}]nonane-9-spiro-2'-[1,3]dioxolane (**1**) in a variety of acids, such as formic acid, *p*-toluenesulfonic acid (*p*-TsOH), and hydrochloric acid.

Results and Discussion

4-Homocubanemethanol **1** was easily accessible from methyl 1-bromopentacyclo[4.3.0.0^{2,5}.0^{3,8}.0^{4,7}]nonane-9-spiro-2'-[1,3]dioxolane-4-carboxylate.⁴⁾

First, reactions were carried out in formic acid, which was known to be a solvent having a high *Y* value.⁵⁾ The results are given in Table 1. When **1** was treated with formic acid, a mixture of **2a** and **3a** was obtained (Entry 1). The product ratio, 1/2.0 of **2/3**, was calculated from the yield of each compound. Interestingly, reaction of **1** with formic acid in dichloromethane gave rearranged chloro derivatives (**2c** and **3c**) along with rearranged formates (**2a** and **3a**) (Entry 2). Chloride ion

	SE(kcal/mol)
	166
	125
	82
	80

Chart 1.

Table 1. Cationic Rearrangement of **1**

Entry	Conditions ^{a)}		Product (% Yield) ^{e)}		Product ratio of 2/3 ^{g)}
	Acid/Solvent	Time/min			
1	HCOOH	15	2a (27),	3a (55)	1/2.0
2	HCOOH/CH ₂ Cl ₂ ^{b)}	15	2a (15), 2c (4)	3a (31), 3c (18)	1/2.6
3	HCOOH/MeOH ^{b)}	15	2a (4), 2b (9), 2d (4)	3a (31), 3b (2), 3d (20)	1/3.1
4 ^{d)}	<i>p</i> -TsOH/MeOH	15	2d (21), 2e (n.d.), ^{f)}	3d (60), 3e (8)	1/3.2
5 ^{d)}	<i>p</i> -TsOH/MeOH +CH ₂ Cl ₂ ^{c)}	15	2d (13), 2e (n.d.), ^{f)}	3d (69), 3e (10)	1/6.1
6 ^{d)}	HCl/MeOH	15	2c (n.d.), ^{f)} , 2d (14)	3c (14), 3d (60)	1/5.3

a) Reactions were carried out at room temperature. b) A (1:1) mixture of HCOOH and CH₂Cl₂ (or MeOH). c) A (1:1) mixture of MeOH and CH₂Cl₂. d) 0.3 Molar amount of acid was employed in each experiment. e) Isolated yield. f) n. d.; Not detected. g) Product ratio was calculated from isolated yields of **2** and **3**.

as a nucleophile arises from hydrochloric acid, which is formed by a reaction of formic acid and dichloromethane.⁶⁾ The product ratio of **2/3** was 1/2.6. When **1** was treated with formic acid in methanol, a mixture of **2a**, **2b**, **2d**, **3a**, **3b**, and **3d** was produced (Entry 3). The rearranged hydroxy derivatives (**2b** and **3b**), which were hydrolysis products of **2a** and **3a**, were also prepared in quantitative yields by the alkaline hydrolysis of **2a** and **3a**. The reaction of **1** with a 0.3 molar amount of *p*-TsOH in methanol gave a mixture of **2d**, **3d**, and **3e** (Entry 4). In Entries 3 and 4, the product ratio of **2/3** is shown to be about 1/3, respectively, resulting in an increase in the yield of **3**. Upon the treatment of **1** with a 0.3 molar amount of *p*-TsOH in a (1:1) mixture of methanol and dichloromethane, a mixture of **2d**, **3d**, and **3e** in a product ratio of 1/6.1 in **2/3** was given without the formation of **2e** (Entry 5). The reaction of **1** with a 0.3 molar amount of hydrochloric acid in methanol gave a mixture of **2d**, **3c**, and **3d** in the product ratio of 1/5.3 in **2/3** (Entry 6). In this series, it was shown that the reaction of **1** with formic acid gave **2** in the highest yield.

The cationic rearrangement of **4** and **5** in formic acid has been reinvestigated.⁷⁾ The reaction of **4** with formic acid gave a mixture of *D*_{2h}- and *C*₂-bishomocubane derivatives in a product ratio of 1/3.5. On the other hand, a similar treatment of **5** with formic acid gave only *C*₂-bishomocubane.

Structure of *D*_{2h}- and *C*₂-Bishomocubanes. Compound **2d** was isolated as colorless prisms by recrystallization from acetonitrile, and was found to have the formula C₂₇H₂₇BrO₅ based on elemental analysis and mass spectral data. In the proton nuclear magnetic resonance (¹H NMR) spectrum the symmetry in the structure of **2d** was characterized by two singlets for three methoxyl groups at δ=3.38 and 3.75 and symmetrical double-doublet protons for two benzene rings at δ=7.24 and 6.76. Four proton signals observed between δ=4.23 and 3.95 were assigned to the ethylene group of spiro[1,3]dioxolane ring. Two kinds of 2H multiplets at δ=3.02—3.00 and δ=2.98—2.96 were assigned to each proton of the 2- and 7-positions and the 3- and 8-positions.


Another two kinds of 1H multiplets at δ=3.70—3.68 and δ=2.80—2.79 were assigned to each proton of the 9- and 6-positions. From a comparison with the signals of aliphatic protons of the *C*₂-bishomocubane derivatives reported in the literature,^{2–4)} **2d** was shown to have a *D*_{2h}-bishomocubane skeleton. The structure of **2d** was unambiguously established as being 4-bromo-1-methoxy-10,10-bis(4-methoxyphenyl)pentacyclo-[5.3.0.0^{2,6}.0^{3,9}.0^{4,8}]decane-5-spiro-2'-[1,3]dioxolane by X-ray crystallography. X-Ray analytical data for **2d** are shown in Table 2, and its ORTEP drawing is given in Fig. 1. The structures of other derivatives **2** were determined based on their spectral data, compared with those of **2d**.

Compound **3d** was shown to have the composition C₂₇H₂₇BrO₅ from an elemental analysis and mass-spectral data. In the ¹H NMR spectrum, **3d**, which was characterized by three signals for three different methoxyl groups at δ=3.74, 3.73, and 3.62 and double-doublet protons for two benzene rings at δ=7.36 and 6.75 and at δ=7.28 and 6.73, was shown to have an unsymmetrical structure. Four proton signals observed between δ=4.26 and 3.84 were assigned to the ethylene group of the spiro[1,3]dioxolane ring. Two kinds of 2H multiplets at δ=3.44—3.42 and δ=3.09—3.04 were as-

Table 2. Single-Crystal X-Ray Analytical Data of **2d**

Formula	C ₂₇ H ₂₇ BrO ₅
Crystal system	Monoclinic
Space group	<i>P</i> 2 ₁ / <i>n</i>
Cell parameter	<i>a</i> =19.617(3) Å <i>b</i> = 7.726(1) Å <i>c</i> =16.813(2) Å <i>β</i> =109.77(1) ° <i>V</i> =2343 Å ³
<i>Z</i> value	4
<i>D</i> _{calcd}	1.450 g cm ⁻³
<i>R</i> (<i>R</i> _w)	0.055 (0.060)
No. of Ref.	
Measured	5971
Used ^{a)}	3011

a) |*F*_o|>3σ|*F*_o|.

Fig. 1. ORTEP drawing of **2d**.

signed to each proton of the 1- and 8-positions and 3- and 4-positions. Another two kinds of ¹H multiplets at $\delta=3.25\text{--}3.24$ and $\delta=2.17\text{--}2.15$ were assigned to each proton of the 2- and 7-positions. Thus, the structure of **3d** was assigned to 5-bromo-9-methoxy-10,10-bis(4-methoxyphenyl)pentacyclo[5.3.0.0^{2,5}.0^{3,9}.0^{4,8}]decane-6-spiro-2'-[1,3]dioxolane, compared with those spectral data of the *C*₂-bishomocubane derivatives reported in the literature.²⁻⁴ The structures of other derivatives **3** were determined in comparison with their spectral data with **3d**.

The cationic rearrangement of 4-homocubanemethanols to *D*_{2h}-bishomocubanes may be explained by the mechanism shown in Scheme 2.⁸ Formic acid is known to be a solvent having a high *Y* value. Reflecting its high ionizing power, 4-homocubanemethanol **1** produces, comparatively, a stable 4-homocubanyl cation (**A**) bearing a *p*-methoxyphenyl group on the α -carbon. This cation is converted into bishomocubanyl cations (**B** and **C**) by a Wagner-Meerwein rearrangement. Both of the cations react with external nucleophiles to give respective *D*_{2h}- and *C*₂-bishomocubane derivatives. Although the rearrangement into the bishomocubane system is driven by a concomitant release of strain, the stability of the free carbocation intermediate **A** may play an important role in determining the course of the cage expansion into the *D*_{2h}-bishomocubane system. A detailed mechanistic study of this reaction remains to be made at the present time.

In conclusion, it may be pointed out that the cationic rearrangement of 4-homocubanemethanol **1** in formic acid provides an attractive route for the synthesis of *D*_{2h}-bishomocubane bridgehead alcohols and related compounds. These compounds are not easily accessible by other routes.

With respect to the formation of the *D*_{2h}-bishomocubane skeleton, our findings may indicate

that: 1) the rearrangement into the bishomocubane system is mainly driven by a concomitant release of strain, and 2) the stability of the homocubanemethyl cation intermediate **A** plays an important role in determining the course of the cage expansion into the *D*_{2h}-bishomocubane system through a migration of the C₄-C₅ bond in the homocubane skeleton.

Experimental

The melting points were measured in a Gallenkamp melting-point apparatus, and are uncorrected. The IR spectra were recorded on a Hitachi 260-30 infrared spectrophotometer and ¹H NMR and ¹³C NMR spectra were measured on Hitachi R-90 (90 MHz), JEOL JNM-EX 270 (270 MHz) and Bruker AM 360 (360 MHz) spectrometers with tetramethylsilane used as an internal standard. The chemical shifts are reported in ppm (δ), and the signals are described as s (singlet), d (doublet), t (triplet), m (multiplet), q (quartet), quin (quintet) or br (broad). All spectra are consistent with the assigned structures. Mass spectra (MS) and high-resolution mass spectra (HRMS) were obtained on a JMS-DX 300 spectrometer operating at an ionization potential of 70 eV. Combustion analyses were performed on a Perkin-Elmer Model 240C elemental analyzer.


The solvents were dried over molecular sieves 4A overnight. The reagents employed in this study were commercially available.

1-Bromo-4-[hydroxybis(4-methoxyphenyl)methyl]pentacyclo[4.3.0.0^{2,5}.0^{3,8}.0^{4,7}]nonane-9-spiro-2'-[1,3]dioxolane (1**).** Methyl 1-bromopentacyclo[4.3.0.0^{2,5}.0^{3,8}.0^{4,7}]nonane-9-spiro-2'-[1,3]dioxolane-4-carboxylate⁴ (1.0 g, 3.19 mmol) was added to the Grignard reagent, prepared from 4-bromoanisole (1.3 g, 7.02 mmol) and Mg (171 mg, 7.02 mmol) in THF (20 ml). The mixture was refluxed for 1.5 h, poured into saturated aqueous NH₄Cl and extracted with Et₂O. The ethereal solution was dried over MgSO₄ and evaporated to give **1** as crystals, which were recrystallized from a mixture of CHCl₃ and hexane to give pure **1** as colorless prisms. Yield: 74%; mp 167.8–169.0 °C; ¹H NMR (360 MHz, CDCl₃) $\delta=7.15\text{--}6.83$ (8H, A₂B₂q, *J*=8.7 Hz), 4.22–4.18 (2H, m), 3.92–3.84 (2H, m), 3.79 (6H, s), 3.68–3.65 (1H, m), 3.59–3.52 (4H, m), 2.31–2.28 (1H, m), 1.96 (1H, s); ¹³C NMR (90 MHz, CDCl₃) $\delta=158.7$, 136.5, 128.4, 127.8, 124.1, 113.5, 66.0, 65.6, 58.3, 55.3, 46.2, 43.2, 43.1, 41.6; IR (CH₂ClCH₂Cl) 3600, 2960, 1615, 1515, 1450, 1425, 1315 cm⁻¹; MS *m/z* 496 (*M*⁺). Found: C, 62.72; H, 4.91; Br, 15.84%. Calcd for C₂₆H₂₅BrO₅: C, 62.78; H, 5.06; Br, 16.06%.


Reaction of 4-Homocubanemethanol **1 under a Variety of Conditions.** A typical example is given to illustrate the general procedure. A mixture of **1** (100 mg, 0.200 mmol) and formic acid (2 ml) was stirred for 15 min at room temperature, poured into ice-water (10 ml) and extracted twice with each of CH₂Cl₂ (20 ml). The CH₂Cl₂ layer was dried over MgSO₄ and evaporated to give oily products. The products were purified by column or preparative thin-layer chromatographies. The results are given in Table 1.

Products **2** and **3** are summarized along with the elemental analysis in Table 3 and the ¹H and ¹³C NMR spectral data of these compounds are given in Table 4.

Single-Crystal X-Ray Analysis of **2d.** Com-



Scheme 1.



Scheme 2.

Table 3. D_{2h} - and C_2 -Bishomocubanes (**2** and **3**)

Compd No.	Appearance	Mp $^{\circ}\text{C}$	Recryst. solvt. ^{a)}	Formula	Analysis			
					Calcd (Found)			
					C	H	Br	S
2a	Colorless needles	194.0—195.2	CH_2Cl_2 -Hex	$\text{C}_{27}\text{H}_{25}\text{BrO}_6$	61.72 (61.74)	4.80 4.52	15.21 15.52)	
2b	Pale yellow oil			$\text{C}_{26}\text{H}_{25}\text{BrO}_5$	HRMS 496.0884 ^{b)} (496.0870)			
2c	Colorless solids	190.5—191.5	CH_2Cl_2 -Hex	$\text{C}_{26}\text{H}_{24}\text{BrClO}_4$	60.54 (60.72)	4.69 4.54)		
2d	Colorless prisms	165.7—166.8	CH_3CN	$\text{C}_{27}\text{H}_{27}\text{BrO}_5$	63.42 (63.56)	5.32 5.19	15.63 15.60)	
3a	Colorless solids	183.2—184.2	CH_2Cl_2 -Hex	$\text{C}_{27}\text{H}_{25}\text{BrO}_6$	61.72 (61.96)	4.80 4.58	15.21 15.33)	
3b	Colorless needles	131.5—132.8	CHCl_3 -Hex	$\text{C}_{26}\text{H}_{25}\text{BrO}_5$	62.79 (62.83)	5.07 4.98	16.07 16.16)	
3c	Colorless solids	182.4—183.4	CH_2Cl_2 -Hex	$\text{C}_{26}\text{H}_{24}\text{BrClO}_4$	60.54 (60.73)	4.69 4.45)		
3d	Colorless needles	206.1—207.2	CH_3CN	$\text{C}_{27}\text{H}_{27}\text{BrO}_5$	63.42 (63.35)	5.32 5.32	15.63 15.75)	
3e	Colorless plates	224.8—225.8	CH_3CN	$\text{C}_{33}\text{H}_{31}\text{BrSO}_7$	60.83 (60.97)	4.80 4.59	12.26 12.46	4.92 4.84)

a) Hex; Hexane. b) High-resolution mass spectrum.

pound **2d** was recrystallized from acetonitrile at room temperature to give colorless prisms. The diffracted intensities were recorded at room temperature on a Rigaku AFC-5FOS four-circle diffractometer (ω - 2θ scan $2\theta < 55^{\circ}$, Mo $\text{K}\alpha$, $\lambda = 0.71069$ Å). The structure was solved by direct method (MULTAN-84),⁹⁾ and refined by a block-diagonal least-squared method.¹⁰⁾ The final R -factor was determined by a procedure described in the literature.^{11,12)} The results

are shown in Table 2 and an ORTEP drawing¹³⁾ of **2d** is depicted in Fig. 1.

We would like to express our thanks to the Material Analytical Center of the ISIR, Osaka University, for spectral measurements, microanalysis, and X-ray crystal analysis.

Table 4. ¹H and ¹³C NMR Spectral Data of **2** and **3**

	¹ H NMR spectra (360 MHz, CDCl ₃) (ppm)	¹³ C NMR spectra (90 MHz, CDCl ₃) (ppm)
2a	7.87 (1H, s), 7.14 (4H, d, <i>J</i> =9.0 Hz), 6.78 (4H, d, <i>J</i> =9.0 Hz), 4.20 (2H, m), 3.97 (2H, m), 3.79 (1H, m), 3.76 (6H, s), 3.26 (2H, dd, <i>J</i> =7.0, 5.0 Hz), 3.05 (2H, dd, <i>J</i> =7.0, 5.0 Hz), 2.95 (1H, m).	160.2, 158.4, 131.6, 129.4, 122.1, 113.9, 95.3, 66.2, 63.6, 63.5, 55.1, 50.9, 48.3, 45.6, 42.4.
2b^{a)}	7.23 (4H, d, <i>J</i> =9.0 Hz), 6.78 (4H, d, <i>J</i> = 9.0 Hz), 4.21—4.16 (2H, m), 3.97—3.92 (2H, m), 3.74 (6H, s), 3.75—3.70 (1H, m), 2.93 (2H, m), 2.88 (1H, m), 2.75 (2H, m), 2.39 (1H, s).	157.9, 132.8, 129.1, 122.4, 113.7, 90.1, 66.0, 64.4, 62.2, 55.1, 51.0, 47.7, 46.4, 42.2.
2c	7.21 (4H, d, <i>J</i> =9.0 Hz), 6.78 (4H, d, <i>J</i> = 9.0 Hz), 4.20 (2H, m), 3.98 (2H, m), 3.79 (1H, m), 3.77 (6H, s), 3.04 (5H, m).	158.2, 132.0, 129.6, 121.7, 113.4, 77.8, 66.1, 64.8, 63.7, 55.1, 52.8, 50.5, 48.9, 42.6.
2d	7.24 (4H, d, <i>J</i> =9.0 Hz), 6.76 (4H, d, <i>J</i> = 9.0 Hz), 4.23—4.20 (2H, m), 3.99—3.95 (2H, m), 3.75 (6H, s), 3.70—3.68 (1H, m), 3.38 (3H, s), 3.02—3.00 (2H, m), 2.98—2.96 (2H, m), 2.80—2.79 (1H, m).	157.9, 133.6, 129.6, 122.4, 113.4, 95.8, 66.1, 64.4, 62.6, 55.7, 55.1, 51.0, 48.0, 44.3, 43.4.
3a	8.29 (1H, s), 7.31 (2H, d, <i>J</i> =9.0 Hz), 7.12 (2H, d, <i>J</i> =9.0 Hz), 6.78 (2H, d, <i>J</i> = 9.0 Hz), 6.76 (2H, d, <i>J</i> =9.0 Hz), 4.23— 4.14 (2H, m), 3.98—3.87 (2H, m), 3.76 (3H, s), 3.75 (3H, s), 3.52 (2H, m), 3.41 (1H, m), 3.33 (1H, m), 2.98 (1H, m), 2.18 (1H, m).	159.1, 158.4, 157.9, 135.0, 132.4, 129.5, 129.0, 120.3 113.6, 113.5, 92.3, 66.2, 65.8, 65.6, 61.8, 55.2, 55.1, 50.5, 49.9, 49.1, 46.8, 45.0, 43.2.
3b	7.28 (2H, d, <i>J</i> =9.0 Hz), 7.26 (2H, d, <i>J</i> = 9.0 Hz), 6.78 (2H, d, <i>J</i> =9.0 Hz), 6.77 (2H, d, <i>J</i> =9.0 Hz), 4.24—4.13 (2H, m), 3.99—3.87 (2H, m), 3.76 (3H, s), 3.75 (3H, s), 3.41 (1H, m), 3.28 (1H, m), 3.19 (2H, m), 2.70 (1H, m), 2.24 (1H, bs), 2.16 (1H, m).	158.1, 157.7, 135.8, 133.4, 129.2, 128.9, 120.2, 113.7, 113.6, 88.4, 66.1, 65.7, 63.5, 62.8, 55.2, 52.1, 49.4, 49.2, 46.1, 44.1, 42.9.
3c	7.34 (2H, d, <i>J</i> =9.0 Hz), 7.16 (2H, d, <i>J</i> = 9.0 Hz), 6.78 (2H, d, <i>J</i> =9.0 Hz), 6.77 (2H, d, <i>J</i> =9.0 Hz), 4.24—4.13 (2H, m), 3.98—3.85 (2H, m), 3.77 (3H, s), 3.76 (3H, s), 3.45 (2H, m), 3.38 (1H, m), 3.29 (1H, m), 2.95 (1H, m), 2.18 (1H, m).	158.4, 157.9, 134.9, 131.9, 129.6, 129.3, 119.7, 113.4, 113.3, 75.0, 66.3, 66.2, 65.8, 61.6, 55.1, 53.5, 53.2, 50.6, 47.2, 46.5, 43.3.
3d	7.36 (2H, d, <i>J</i> =9.0 Hz), 7.28 (2H, d, <i>J</i> = 9.0 Hz), 6.75 (2H, d, <i>J</i> =9.0 Hz), 6.73 (2H, d, <i>J</i> =9.0 Hz), 4.26—3.84 (4H, m), 3.74 (3H, s), 3.73 (3H, s), 3.62 (3H, s), 3.44—3.42 (2H, m), 3.25—3.24 (1H, m), 3.09— 3.04 (2H, m), 2.17—2.15 (1H, m).	158.0, 157.5, 135.9, 133.7, 129.5, 129.2, 120.7, 113.3, 113.2, 94.4, 66.1, 65.7, 64.6, 62.7, 55.1, 55.0, 54.9, 51.6, 49.3, 46.7, 46.2, 44.1, 41.4.
3e	7.84 (2H, d, <i>J</i> =8.0 Hz), 7.32 (2H, d, <i>J</i> = 8.0 Hz), 7.18 (2H, d, <i>J</i> =9.0 Hz), 7.09 (2H, d, <i>J</i> =9.0 Hz), 6.72 (2H, d, <i>J</i> =9.0 Hz), 6.70 (2H, d, <i>J</i> =9.0 Hz), 4.21—3.84 (4H, m), 3.83—3.81 (1H, m), 3.75 (3H, s), 3.74 (3H, s), 3.37—3.30 (3H, m), 3.12—3.09 (1H, m), 2.44 (3H, s), 2.19—2.17 (1H, m).	158.3, 157.8, 144.8, 134.9, 134.3, 131.7, 130.0, 129.3, 128.9, 127.6, 120.1, 113.5, 113.4, 96.8, 66.2, 66.1, 65.7, 61.6, 55.1, 50.2, 50.0, 49.1, 46.8, 46.5, 44.8, 42.8, 21.6.

a) ¹H NMR; 270 MHz, ¹³C NMR; 67.5 MHz.

References

- 1) E. Osawa, K. Aigami, and Y. Inamoto, *J. Chem. Soc., Perkin Trans. 2*, **1979**, 181.
 - 2) A. J. H. Klunder and B. Zwanenburg, *Tetrahedron*, **29**, 161 (1973).
 - 3) T. C. W. Mak, Y. C. Yip, and T.-Y. Luh, *Tetrahedron*, **42**, 1981 (1986).
 - 4) N. B. Chapman, J. M. Key, and K. J. Toyne, *J. Org. Chem.*, **35**, 3860 (1970).
 - 5) a) E. Grunwald and S. Winstein, *J. Am. Chem. Soc.*, **70**, 846 (1948); b) W. Bentley and P. von R. Schleyer, *J. Am. Chem. Soc.*, **98**, 7658 (1976).
 - 6) Chloride ion as a nucleophile is produced from formic acid and dichloromethane: $\text{HCOOH} + \text{CH}_2\text{Cl}_2 \rightarrow \text{HCl} + \text{CH}_3\text{Cl} + \text{CO}_2$.
 - 7) T. Hasegawa, Ph. D. Dissertation, Osaka University, Osaka, Japan, 1993.
 - 8) J. P. Richard, "Simple Relationships between Carbocation Lifetime and the Mechanism for Nucleophilic Substitution at Saturated Carbon," in "Advances in Carbocation Chemistry," ed by X. Creary, JAI Press Inc., London, England (1989), Vol. 1, pp. 121—169.
 - 9) P. Main, G. Germain, and M. M. Woolfson, "A System of Computer Programs for the Automatic Solution of Crystal Structure from X-Ray Diffraction Data, MULTAN 84," Universities of York, England and Louvain, Belgium (1984).
 - 10) RASA software was employed for the entire structure analysis.
 - 11) "Atomic Scattering Factors from International Tables for X-Ray Crystallography," Kynoch Press, Birmingham (1974), Vol. 4.
 - 12) All calculations were made on a FACOM S3500 computer at Material Analytical Center, the Institute of Scientific and Industrial Research, Osaka University.
 - 13) ORTEP-II employed for drawing molecular structure: C. K. Johnson, "ORTEP Report ORNL-5138," Oak Ridge National Laboratory, Tennessee (1976).
-