

# Photocatalysis of $\text{RhCl}(\text{PCy}_3)_2$ for Cyclohexane Dehydrogenation: Thermal Dissociation of C–H Bond and Photoelimination of $\text{H}_2$


Hiroaki ITAGAKI, Hiroki MURAYAMA, and Yasukazu SAITO\*

Department of Industrial Chemistry, Faculty of Engineering, The University of Tokyo, Hongo, Bunkyo-ku, Tokyo 113

(Received December 6, 1993)

Catalytic cyclohexane dehydrogenation, yielding cyclohexene and dihydrogen, proceeded under photoirradiation on either a three-coordinated complex,  $\text{RhCl}(\text{PCy}_3)_2$  ( $\text{Cy}$ =cyclohexyl), or a dihydride complex,  $\text{RhClH}_2(\text{PCy}_3)_2$ , with almost the same turnover frequencies (8.4 or 8.6, respectively) attained by use of a cut-filter (UV-27) under refluxing conditions (354 K).  $\text{RhCl}(\text{PCy}_3)_2$  in cyclohexane gave stoichiometric amounts of cyclohexene and  $\text{RhClH}_2(\text{PCy}_3)_2$  at 354 K; the latter complex yielded little  $\text{H}_2$  even at 373 K in toluene. A photocatalysis cycle for cyclohexane dehydrogenation with  $\text{RhCl}(\text{PCy}_3)_2$  is proposed, where cyclohexene is yielded by thermal C–H bond dissociation and dihydrogen is photoeliminated from  $\text{RhClH}_2(\text{PCy}_3)_2$ , regenerating the original complex.

Functionalization of saturated hydrocarbons under mild conditions is one of the most challenging targets in catalytic chemistry. A lot of advances in the C–H bond activation have been made regarding saturated hydrocarbons with transition-metal complexes.<sup>1)</sup> Remarkable photocatalytic activities for alkane dehydrogenation (Eq. 1) were found with Vaska-type rhodium complexes  $\text{RhCl}(\text{CO})(\text{PR}_3)_2$ .<sup>2)</sup>



Flash photolysis studies revealed the role of a three-coordinated intermediate  $\text{RhCl}(\text{PR}_3)_2$  ( $\text{R}=\text{Ph}$ , *p*-tolyl, and Me), generated from  $\text{RhCl}(\text{CO})(\text{PR}_3)_2$  under photoirradiation.<sup>3)</sup> An electronic absorption band effective in CO dissociation was assigned by extended Hückel molecular orbital calculations to metal-to-ligand charge transfer (m.l.c.t.) from HOMO to a certain unoccupied orbital, antibonding with respect to the Rh–C bond.<sup>4)</sup> Moreover, thermocatalytic dehydrogenation of cyclooctane was achieved with the Wilkinson complexes  $\text{RhCl}(\text{PR}_3)_3$  ( $\text{R}=\text{Ph}$ ,  $\text{C}_6\text{H}_4\text{Me-}p$  or  $\text{R}_3=\text{MePh}_2$ ),<sup>5)</sup> for which the key role of  $\text{RhCl}(\text{PR}_3)_2$  was elucidated kinetically.<sup>6)</sup>

Formation of  $\text{RhClH}(\text{C}_6\text{H}_{11})(\text{PMe}_3)_2$  in cyclohexane<sup>3c,3d)</sup> or of  $\text{RhClH}_2(\text{P}^i\text{Pr}_3)_2$  and  $\text{H}_2[\text{Rh}(\text{P}^i\text{Pr}_3)_2\text{Cl}]_2$  in cyclooctane<sup>7)</sup> offers us pertinent suggestions on the reaction mechanism of photocatalytic alkane dehydrogenation. In the present study,  $\text{RhCl}(\text{PCy}_3)_2$  is reacted with cyclohexane under thermal and photocatalysis conditions, with attention being paid to C–H bond splitting and  $\text{H}_2$  elimination processes (Chart 1).

## Experimental

All manipulations were carried out under Ar or




Chart 1.

$\text{N}_2$  atmosphere. Stable complexes  $\text{RhCl}(\text{PCy}_3)_2$ <sup>8)</sup> and  $\text{RhClH}_2(\text{PCy}_3)_2$ <sup>9)</sup> were prepared by the published methods. Cyclohexane of reagent grade was dried and deaerated over metallic sodium and benzophenone by refluxing under  $\text{N}_2$  flow.

Photocatalytic dehydrogenation of cyclohexane was carried out under boiling and refluxing conditions (354 K) in a cylindrical quartz cell (diameter 45 mm, cell length 80 mm), which was irradiated by a xenon lamp (2 kW, external-type, Ushio) or an ultra high-pressure Hg lamp (500 W, external-type, Ushio) utilizing cut-off filters: UV-27, UV-29, UV-32, UV-36, and L-42 (Kenko), where the L-42 filter was specified to cut off wavelengths <420 nm (50% transmittance at 420 nm). The amount of gas evolved was measured by a gas buret (10 cm<sup>3</sup>). The reaction products were analyzed by gas chromatography using active carbon and PEG-20M columns for gas- and liquid-phase products, respectively. The UV/vis spectrum of  $\text{RhClH}_2(\text{PCy}_3)_2$  complex was recorded on a UV-365 (Shimadzu) spectrometer.

Thermal reactivities of  $\text{RhCl}(\text{PCy}_3)_2$  with cyclohexane and of  $\text{RhClH}_2(\text{PCy}_3)_2$  in toluene were investigated in situ in an NMR tube under an argon atmosphere. <sup>1</sup>H and <sup>31</sup>P{<sup>1</sup>H} spectra were recorded on JEOL GX 400 and JNM 90Q spectrometers, respectively, operating in the FT mode. <sup>31</sup>P chemical shifts were referred to an external standard of 85 %  $\text{H}_3\text{PO}_4$ , whereas <sup>1</sup>H spectra were referenced internally to the toluene-*d*<sub>8</sub> solvent and converted to the TMS scale.

## Results and Discussion

### Photocatalytic Dehydrogenation of Cyclohexane.

Photocatalytic cyclohexane dehydrogenation proceeded with  $\text{RhClH}_2(\text{PCy}_3)_2$  through a UV-29 filter, giving a total turnover number of 8.1 at the first (after 1.5 h) and 12.0 at the second (after 3 h) stage, respectively (Fig. 1). Since this complex decomposed gradually under reaction conditions (vide infra), the turnover frequency decreased from 10.8 h<sup>−1</sup> (the first stage initially) to 5.1 h<sup>−1</sup> (the first stage finally) and to 3.4 h<sup>−1</sup> (the second stage). Without photoirradiation, little dihydrogen evolved even at refluxing conditions (354 K).

**Wavelength Dependence of Photocatalytic Activity.** Figure 2 shows the electronic absorption




Fig. 1. Time-course plot for photocatalytic dehydrogenation of cyclohexane with  $\text{RhClH}_2(\text{PCy}_3)_2$ ; catalyst concentration  $10 \mu\text{mol}$  per  $100 \text{ cm}^3$  cyclohexane, reaction temperature  $354 \text{ K}$  (reflux), cut-filter UV-29, and ultra high-pressure Hg lamp ( $500 \text{ W}$ ) light source.




Fig. 2. UV-vis spectrum of  $\text{RhClH}_2(\text{PCy}_3)_2$  with reference to transmission characteristics of the cut-off filters; catalyst concentration  $5.0 \mu\text{mol}$  per  $100 \text{ cm}^3$  toluene and cell length  $10 \text{ mm}$ .

spectrum of  $\text{RhClH}_2(\text{PCy}_3)_2$  together with transmission characteristics of cut-filters used for the photoreaction. The photocatalytic reaction rates were dependent on wavelength and increased in the order of  $\text{L-42} < \text{UV-36} < \text{UV-32} < \text{UV-27}$  (Table 1). The UV-27 filter gave a high rate, whereas little  $\text{H}_2$  evolution was observed with the L-42 filter.

Wavelength dependence was investigated in a similar manner for the photocatalytic activity of  $\text{RhCl}(\text{PCy}_3)_2$ . Photoirradiation was indispensable for  $\text{RhCl}(\text{PCy}_3)_2$  to dehydrogenate cyclohexane catalytically, as observed for  $\text{RhClH}_2(\text{PCy}_3)_2$ . The photoreaction rates of these complexes were almost the same at each wavelength region (Table 1). It seems reasonable to assume, therefore, that both  $\text{RhClH}_2(\text{PCy}_3)_2$  and  $\text{RhCl}(\text{PCy}_3)_2$  lie on the same photoreaction cycle.

Stoichiometric coincidence of dihydrogen with cyclohexene was excellent. Neither cyclohexadiene nor benzene was ever detected by GC analysis. Little dehydrogenation from the  $\text{PCy}_3$  ligand itself under the

Table 1. Wavelength Dependence on the Rate of Cyclohexane Dehydrogenation with  $\text{RhCl}(\text{H})_2(\text{PCy}_3)_2$  and  $\text{RhCl}(\text{PCy}_3)_2$  <sup>a)</sup>

| Cut-filter | $\text{RhCl}(\text{H})_2(\text{PCy}_3)_2$ |                           | $\text{RhCl}(\text{PCy}_3)_2$ |                           |
|------------|-------------------------------------------|---------------------------|-------------------------------|---------------------------|
|            | $\text{H}_2$ <sup>b)</sup>                | $\text{C}_6\text{H}_{10}$ | $\text{H}_2$                  | $\text{C}_6\text{H}_{10}$ |
|            | $\mu\text{mol}$                           | $\mu\text{mol}$           | $\mu\text{mol}$               | $\mu\text{mol}$           |
| UV-27      | 86                                        | 74                        | 84                            | 72                        |
| UV-32      | 47                                        | 39                        | 48                            | 52                        |
| UV-36      | 34                                        | 20                        | 35                            | 34                        |
| L-42       | ca.0                                      | ca.0                      | ca.0                          | ca.0                      |

a)  $\text{RhCl}(\text{H})_2(\text{PCy}_3)_2$   $10 \mu\text{mol}$ ,  $\text{RhCl}(\text{PCy}_3)_2$   $10 \mu\text{mol}$ , cyclohexane  $100 \text{ cm}^3$ ,  $354 \text{ K}$  (reflux), Xe lamp ( $2 \text{ kW}$ ) light source, reaction period  $1 \text{ h}$ . b) Contribution of catalytic  $\text{H}_2$  evolution from  $\text{C}_6\text{H}_{12}$  with  $\text{RhCl}(\text{H})_2(\text{PCy}_3)_2$  can be estimated by subtracting the  $\text{H}_2$  amount of its dihydride ligand ( $10 \mu\text{mol}$ ) from the observed values.

reaction conditions was observed; this absence would be attributed at least partly to the low concentration of the  $\text{PCy}_3$ -coordinated complex ( $0.1 \text{ mM}$  ( $1 \text{ mM} = 1 \text{ mmol dm}^{-3}$ )).

#### Reactivity of $\text{RhCl}(\text{PCy}_3)_2$ with Cyclohexane.

Photocatalysis for cyclohexane dehydrogenation with these  $\text{PCy}_3$ -coordinated complexes should contain the processes of C-H bond splitting and  $\text{H}_2$  elimination.

When the cyclohexane solution of  $\text{RhCl}(\text{PCy}_3)_2$  was heated at  $354 \text{ K}$ , the color changed from violet to yellow, indicating a structural change of the complex. As shown in Fig. 3, the dihydride complex,  $\text{RhClH}_2(\text{PCy}_3)_2$  ( $\delta_{\text{P}} = 50.3 \text{ ppm}$ ,  $^1J_{\text{Rh-P}} = 114.6 \text{ Hz}$ ,  $\delta_{\text{H}} = -22.6 \text{ ppm}$ ,  $^1J_{\text{Rh-H}} = 27.4 \text{ Hz}$ ,  $^2J_{\text{P-H}} = 13.7 \text{ Hz}$ , identified by the authentic sample), was gradually formed as a sole product when  $\text{RhCl}(\text{PCy}_3)_2$  ( $\delta_{\text{P}} = 47.5 \text{ ppm}$ ,  $^1J_{\text{Rh-P}} = 196.5 \text{ Hz}$ ) in cyclohexane was heated at  $354 \text{ K}$ . In addition, a stoichiometric quantity of cyclohexene was detected by GC analysis from the cyclohexane solution of  $\text{RhCl}(\text{PCy}_3)_2$  after its refluxing for  $0.5 \text{ h}$ . A selective reaction process (Eq. 2) under thermal conditions was thus confirmed.



As observed in Fig. 3(c), free  $\text{PCy}_3$  ( $\delta_{\text{P}} = 10.0 \text{ ppm}$ ) was liberated by thermal decomposition of  $\text{PCy}_3$ -coordinated complexes; this release would be attributed at least partly to the gradual decrease of turnover frequency (Fig. 1).

A subtle difference between  $\text{RhCl}(\text{PCy}_3)_2$  and  $\text{RhCl}(\text{P}^i\text{Pr}_3)_2$  in cyclohexane thermolysis is to be noted, since the former could dehydrogenate solvent cyclohexane in a stoichiometric manner and, moreover,  $\text{RhClH}_2(\text{PCy}_3)_2$  was selectively formed during the reaction.<sup>10)</sup> Both selective performance of Eq. 2 and low reactivities of  $\text{PCy}_3$ -coordinated complexes would be correlated to the remarkable bulkiness of the  $\text{PCy}_3$  ligand.

**Reductive Elimination of  $\text{H}_2$  from  $\text{RhClH}_2(\text{PCy}_3)_2$ .** After the heat treatment of  $\text{RhClH}_2(\text{PCy}_3)_2$ ,





Fig. 3. Formation of  $\text{RhClH}_2(\text{PCy}_3)_2$  from  $\text{RhCl}(\text{PCy}_3)_2$  in cyclohexane solution;  $^{31}\text{P}\{^1\text{H}\}$  NMR spectra taken at 298 K for the samples before (a) and after the 0.5 h- (b) or 1.0 h- (c) heat treatment at 354 K.



Scheme 1.

( $\text{PCy}_3$ )<sub>2</sub> in toluene at 373 K for 0.5 h, a trace amount of dihydrogen was detected by GC analysis. The charged amount of  $\text{RhClH}_2(\text{PCy}_3)_2$  was recovered, as ascertained by  $^1\text{H}$  and  $^{31}\text{P}$  NMR analyses. Photoelimination of  $\text{H}_2$  from dihydrido complexes has frequently been pointed out,<sup>11)</sup> which is in conformity with our results

on photocatalytic dehydrogenation of cyclohexane with  $\text{RhClH}_2(\text{PCy}_3)_2$  or  $\text{RhCl}(\text{PCy}_3)_2$  under refluxing conditions (Fig. 1 and Table 1).

Thermal reductive elimination of  $\text{H}_2$  from  $\text{RhClH}_2(\text{PR}_3)_2$  was claimed to be rather facile for electron-withdrawing aryl-phosphine ligands, e.g., such as  $\text{PPh}_3$  and  $\text{P}(p\text{-tolyl})_3$ , in contrast to the cases with electron-donating alkyl-substituted phosphine ligands, e.g.,  $\text{PEtPh}_2$  and  $\text{PMePh}_2$ .<sup>6)</sup> The observed difficulty in thermal  $\text{H}_2$  elimination from  $\text{RhClH}_2(\text{PCy}_3)_2$  can be well interpreted in terms of the electron-donating property of  $\text{PCy}_3$ .

#### Proposal of Photocatalysis Cycle for Alkane Dehydrogenation.

A reaction mechanism for photocatalytic dehydrogenation of cyclohexane with  $\text{RhCl}(\text{PCy}_3)_2$  or  $\text{RhClH}_2(\text{PCy}_3)_2$  is now proposed (Scheme 1). Thermal steps consist of (1) C-H bond splitting of cyclohexane with  $\text{RhCl}(\text{PCy}_3)_2$ , (2) elimination of  $\beta$ -hydrogen from the cyclohexyl ligand, and (3) formation of  $\text{RhClH}_2(\text{PCy}_3)_2$  and cyclohexene. A photoreactive step follows: (4)  $\text{H}_2$  dissociation from  $\text{RhClH}_2(\text{PCy}_3)_2$  and regeneration of  $\text{RhCl}(\text{PCy}_3)_2$ .

#### References

- 1) R. H. Crabtree, *Chem. Rev.*, **85**, 245 (1985); R. G. Bergman, P. F. Seidler, and T. T. Wenzel, *J. Am. Chem. Soc.*, **107**, 4358 (1985); J. M. Buchanan, J. M. Stryker, and R. G. Bergman, *J. Am. Chem. Soc.*, **108**, 1537 (1986); R. A. Periana and R. G. Bergman, *J. Am. Chem. Soc.*, **108**, 7332 (1986); C. Bianchini, P. Barbaro, A. Meli, M. Peruzzini, A. Vacca, and F. Vizza, *Organometallics*, **12**, 2505 (1993), and references therein.
- 2) a) K. Nomura and Y. Saito, *J. Chem. Soc., Chem. Commun.*, **1988**, 161; b) T. Sakakura, T. Sodeyama, Y. Tokunaga, and M. Tanaka, *Chem. Lett.*, **1988**, 263; c) J. A. Maguire, W. T. Boese, and A. S. Goldman, *J. Am. Chem. Soc.*, **111**, 7088 (1989); d) K. Nomura and Y. Saito, *J. Mol. Catal.*, **54**, 57 (1989); e) T. Sakakura, T. Sodeyama, and M. Tanaka, *New J. Chem.*, **13**, 737 (1989); f) J. A. Maguire, W. T. Boese, M. E. Goldman, and A. S. Goldman, *Coord. Chem. Rev.*, **97**, 179 (1990).
- 3) a) D. A. Wink and P. C. Ford, *J. Am. Chem. Soc.*, **107**, 1794 (1985); b) D. A. Wink and P. C. Ford, *J. Am. Chem. Soc.*, **109**, 436 (1987); c) C. T. Spillett and P. C. Ford, *J. Am. Chem. Soc.*, **111**, 1932 (1989); d) P. C. Ford, T. L. Netzel, C. T. Spillett, and D. B. Pourreau, *Pure Appl. Chem.*, **62**, 1091 (1990).
- 4) A. Iwamoto, H. Itagaki, and Y. Saito, *J. Chem. Soc., Dalton Trans.*, **1991**, 1093.
- 5) T. Fujii and Y. Saito, *J. Chem. Soc., Chem. Commun.*, **1990**, 757.
- 6) T. Fujii, Y. Higashino, and Y. Saito, *J. Chem. Soc., Dalton Trans.*, **1993**, 517.
- 7) K. Shih and A. S. Goldman, *Organometallics*, **12**, 3390 (1993).
- 8) H. L. M. van Gaal and F. L. A. van den Bekerom, *J. Organomet. Chem.*, **134**, 237 (1977).
- 9) H. L. M. van Gaal, J. M. J. Verlaak, and T. Posno, *Inorg. Chim. Acta*, **23**, 43 (1977).

10) Heat treatment of  $\text{RhCl}(\text{P}^i\text{Pr}_3)_2$  (19 mM) in cyclohexane at 363 K for 3 h gave only a small amount of cyclohexene ( $<0.5$  mM) but a lot of product complexes, including  $\text{RhClH}_2(\text{P}^i\text{Pr}_3)_2$  (2.7 mM),  $\text{H}_2[\text{Rh}(\text{P}^i\text{Pr}_3)_2\text{Cl}]_2$  (0.9 mM),  $\text{RhCl}_2\text{H}(\text{P}^i\text{Pr}_3)_2$  (3.6 mM),  $[(^i\text{Pr}_2)\text{P}(\eta^2\text{-MeC=CH}_2)]\text{Rh}(\mu\text{-Cl}_2)\text{Rh}(\text{P}^i\text{Pr}_3)_2\text{H}_2$  (1.9 mM), and  $\text{H}_2(\text{P}^i\text{Pr}_3)_2\text{Rh}(\mu\text{-H}_2)\text{Rh}(\text{P}^i\text{Pr}_3)_2\text{HCl}$  (1.5 mM).<sup>7)</sup>

11) G. L. Geoffroy, G. S. Hammond, and H. B. Gray,

*J. Am. Chem. Soc.*, **97**, 3933 (1975); G. L. Geoffroy and R. Pierantozzi, *J. Am. Chem. Soc.*, **98**, 8054 (1976); G. L. Geoffroy and M. G. Bradley, *Inorg. Chem.*, **16**, 744 (1977); G. L. Geoffroy and M. G. Bradley, *Inorg. Chem.*, **17**, 2410 (1978); W. D. Jones and F. J. Feher, *J. Am. Chem. Soc.*, **106**, 1650 (1984); W. D. Jones and F. J. Feher, *J. Am. Chem. Soc.*, **108**, 4814 (1986); D. A. Wink and P. C. Ford, *J. Am. Chem. Soc.*, **108**, 4838 (1986).

---