

Facile Preparation of 1,6-Anhydro-2-azido-3-*O*-benzyl-2-deoxy- β -D-glucopyranose and Its 4-*O*-Substituted Derivatives

Nobuo SAKAIRI,* Shunya TAKAHASHI, Feng WANG, Yoshihito UENO, and Hiroyoshi KUZUHARA
The Institute of Physical and Chemical Research (RIKEN), Wako, Saitama 351-01

(Received December 8, 1993)

Synopsis. Treatment of 1,6-anhydro-2,3-*O*-endo-benzylidene- β -D-mannopyranose derivatives with trimethylamine–borane (1/1)–aluminium chloride resulted in highly regioselective fission of the cyclic acetals to give the corresponding 2-*O*-unprotected-3-*O*-benzyl derivatives. Trifluoromethanesulfonylation of these, followed by nucleophilic substitution, afforded 2-azido-2-deoxy derivatives in good yields.

1,6-Anhydro-2-azido-3-*O*-benzyl-2-deoxy- β -D-glucopyranose derivatives are versatile intermediates for the synthesis of D-glucosamine-containing complex oligosaccharides,¹⁾ and have been commonly prepared by nucleophilic ring opening of “Cerny epoxide” derived from 1,6-anhydro- β -D-glucopyranose.²⁾ Recently, a short-step preparation of these compounds has been developed employing D-mannopyranose as the starting material.³⁾ However, this procedure is sometimes inapplicable to substrates bearing such labile protecting groups as acetyl or *p*-methoxybenzyl groups because of substantial removal of the protecting groups during reductive ring opening of the 2,3-*O*-benzylidene intermediates with lithium aluminium hydride–aluminium chloride or sodium cyanoborohydride–hydrogen chloride. We now report a modified procedure for the preparation of D-glucosamine precursors by use of trimethylamine–borane (1/1)–aluminium chloride as the reducing agent.

Starting materials, 4-*O*-acetyl (**2**), 4-*O*-*p*-methoxybenzyl (**3**), and 4-*O*-benzyl derivatives (**4**) were derived from crystalline 1,6-anhydro-2,3-*O*-endo-benzylidene- β -D-mannopyranose (**1**)³⁾ by conventional ways. Compound **2** was prepared in 90% yield by acetylation of **1** with Ac₂O–pyridine, while **3** and **4** were prepared in 87 and 91% yields by alkylation of **1** with NaH-*p*-methoxybenzyl chloride–sodium iodide and NaH-benzyl bromide, respectively (Scheme 1). We employed **2** as the model compound to examine the reductive cleavage of the benzylidene group. The reaction with trimethylamine–borane (1/1) and aluminium chloride⁴⁾ was found to proceed regioselectively without any affection to the 4-*O*-acetyl group. Thus, treatment of **2** with an excess amount (5.8 mol equiv) of the reagent in tetrahydrofuran (THF) at room temperature for 5 h gave the 3-*O*-benzyl derivative (**5**) in 87% yield. In a similar way, the hydrogenolytic ring cleavage of the 4-*O*-*p*-methoxybenzyl and 4-*O*-benzyl derivative (**3** and **4**) afforded the 4-*O*-(*p*-methoxybenzyl)-3-*O*-benzyl (**6**) and 3,4-di-*O*-benzyl derivative (**7**) as the sole products (80 and 89% yields, respectively).


The next conversions of the 2-*O*-unprotected D-mannose derivatives into 2-azido-2-deoxy-D-glucose derivatives were carried out in a two-step manner: trifluoromethanesulfonylation and subsequent nucleophilic substitution.^{3,5,6)} Thus, **5** and **6** were treated with trifluoromethanesulfonic anhydride in pyridine–dichloromethane to give the corresponding 2-triflates, which were subjected to S_N2 reaction with lithium azide in *N,N*-dimethylformamide (DMF) at room temperature, giving the 2-azido-2-deoxy- β -D-glucopyranose derivatives (**8** and **9**) in 73 and 81% overall yields, respectively. Compound **7** was also converted into the 2-azido-2-deoxy derivative (**10**) in almost quantitative yield according to Dasgupta and Geregg.⁶⁾ The 2-azido-2-deoxy derivatives thus obtained were readily converted into both glucosamine acceptors and donors. Either Zemplen deacetylation of **8** or 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ) oxidation of **9** afforded known 4-hydroxy derivative (**11**) in almost quantitative yield. Furthermore, **9** was converted into the corresponding thioglycoside and glycosyl trichloroacetimidate, which were used as key intermediates for the synthesis of 2-amino-2-deoxy- β -cyclodextrin.¹¹⁾

In conclusion, D-glucosamine precursors having acid- or base-sensitive protecting groups at *O*-4 position were prepared from D-mannose in good overall yields, employing reductive ring opening of 2,3-*O*-benzylidene group with trimethylamine–borane (1/1)–aluminium chloride as the key reaction.

Experimental

General Procedures. Melting points were determined in a capillary with an Ishii melting-point apparatus and are uncorrected. Optical rotations were determined with a Perkin–Elmer Model 241MC polarimeter. ¹H NMR spectra were recorded at 270 MHz or 500 MHz with JEOL JNM-EX 270 or JEOL JNM-GX 500 spectrometers, using tetramethylsilane as the internal standard. Reactions were monitored by TLC on a precoated plate of silica gel 60F₂₅₄ (layer thickness, 0.25 mm; E. Merck, Darmstadt, Germany). Column chromatography was performed on silica gel 60 (70–230 or 230–400 mesh; E. Merck, Darmstadt, Germany). Molecular sieves 4A was activated at 180–200 °C under reduced pressure on diphosphorus pentaoxide prior to use.

4-*O*-Acetyl-1,6-anhydro-2,3-*O*-endo-benzylidene- β -D-mannopyranose (2**).** A solution of 1,6-anhydro-2,3-*O*-endo-benzylidene- β -D-mannopyranose (**1**)³⁾ (5.0 g, 20 mmol) in pyridine (50 cm³) and acetic anhydride (20 cm³) was stirred at room temperature for 1 d, evaporated under reduced pressure, and coevaporated with toluene sev-



Scheme 1. Reagents and conditions. a) **1**→**2**: Ac₂O–pyridine, r.t.; **1**→**3**: *p*-methoxybenzyl chloride–NaH–NaI, DMF, r.t.; **1**→**4**: benzyl bromide–NaH, DMF, r.t. b) BH₃·Me₃N–AlCl₃, THF, r.t. c) **5**→**8**, **6**→**9**, **7**→**10**: Tf₂O, pyridine–CH₂Cl₂ then LiN₃, DMF, r.t. d) MeONa, MeOH. e) DDQ, aq CH₂Cl₂, r.t.

eral times. Crystallization of the residual syrup with diethyl ether gave 5.3 g (90%) of the 4-acetate (**2**): Mp 190–191 °C (lit, 190–192 °C);⁷⁾ $[\alpha]_{\text{D}}^{23}$ –88° (c 0.20, CHCl₃) (lit, $[\alpha]_{\text{D}}^{20}$ –92° (c 0.7, CHCl₃)).⁷⁾ Found: C, 61.37; H, 5.53%. Calcd for C₁₅H₁₆O₆: C, 61.64; H, 5.52%.

1,6-Anhydro-2,3-O-endo-benzylidene-4-O-(*p*-methoxybenzyl)-β-D-mannopyranose (3). To an ice-cold solution of **1** (60 g, 0.24 mol) and sodium iodide (10 g) in DMF (400 cm³) was added 60% sodium hydride oil dispersion (25 g, 0.63 mol). Then the mixture was stirred at 0 °C for 2 h. *p*-Methoxybenzyl chloride (50 g, 0.32 mol) was added dropwise to the resulting mixture at 0 °C. The suspension was stirred at room temperature for 2 d, quenched by careful addition of MeOH, poured into ice cold aqueous ammonia (0.4 mol dm^{–3}, 2 dm³), and allowed to stand overnight. The precipitates were collected by filtration, and washed with water and EtOH. Recrystallization from EtOH gave 77 g (87%) of the 4-*p*-methoxybenzyl ether (**3**): Mp 105–106 °C; $[\alpha]_{\text{D}}^{27}$ –49° (c 0.21, CHCl₃); ¹H NMR (CDCl₃) δ=3.76 (1H, s, H-4), 3.79–3.83 (4H, m, H-6a, OMe), 3.94 (1H, d, *J*_{6a,6b}=7.3 Hz, H-6a), 4.24 (1H, dd, *J*_{1,2}=2.9 and *J*_{2,3}=7.0 Hz, H-2), 4.28 (1H, d, H-3), 4.60 (1H, d, *J*_{gem}=11.9 Hz, 1/2×CH₂Ar), 4.64 (1H, d, *J*_{5,6}=6.7 Hz, H-5), 4.68 (1H, d, 1/2×CH₂Ar), 5.52 (1H, br.s, H-1), and 5.76 (1H, s, CHPh). Found: C, 67.99; H, 5.98%. Calcd for C₂₁H₂₂O₆: C, 68.10; H, 5.99%.

1,6-Anhydro-4-O-benzyl-2,3-O-endo-benzylidene-β-D-mannopyranose (4). Treatment of **1** (45 g, 0.18 mmol) with 60% sodium hydride oil dispersion (15 g, 0.38 mol) and benzyl bromide (65 g, 0.38 mmol) in DMF (300 cm³) as described for the preparation of **3**, followed by crystallization from EtOH, gave 55.7 g (91%) of the 4-*O*-benzyl derivative (**4**): Mp 126.5–127 °C (lit, mp 127–128 °C);⁸⁾ $[\alpha]_{\text{D}}^{20}$ –51° (c 1.0, CHCl₃) (lit, $[\alpha]_{\text{D}}^{25}$ –49.7° (c 1, CHCl₃)).⁸⁾ ¹H NMR (CDCl₃) δ=3.78 (1H, br.s, H-4), 3.82 (1H, dd, *J*_{5,6}=6.3 and *J*_{6exo,6endo}=7.3 Hz, H-6), 3.95 (1H, dd, *J*_{5,6}=1.0 Hz, H-6), 4.25 (1H, dd, *J*_{1,2}=2.9 and *J*_{2,3}=6.9 Hz, H-2), 4.31 (1H, br.d, H-3), 4.67, 4.75 (1H, 2×d, *J*_{gem}=12.2 Hz, CH₂Ph), 4.66–4.68 (1H, m, H-5), 5.53 (1H, d, H-1), and 5.76 (1H, s, CHPh).

4-O-Acetyl-1,6-anhydro-3-O-benzyl-β-D-mannopyranose (5). To a suspension of **2** (5.0 g, 17 mmol), powdered molecular sieves 4A (5 g), and trimethylamine–borane (1/1) (7.3 g, 0.10 mmol) in THF (100 cm³) was added aluminium chloride (13.3 g, 0.10 mmol) by portions.

The mixture was stirred at room temperature for 5 h, filtered through a Celite pad, and washed with chloroform. The combined filtrate and washings were washed successively with 6% hydrochloric acid, aqueous sodium hydrogencarbonate, and brine, dried, and concentrated. Chromatographic purification on a silica gel column with toluene–EtOAc (3:1 v/v) gave 4.3 g (85%) of the 2-hydroxy derivative (**5**): $[\alpha]_{\text{D}}^{20}$ –61° (c 0.32, CHCl₃); ¹H NMR (CDCl₃) δ=2.13 (3H, s, OAc), 3.03 (1H, d, *J*_{2,OH}=11.6 Hz, OH), 3.63–3.71 (2H, m, H-2,3), 3.75 (1H, dd, *J*_{5,6a}=6.1 and *J*_{6a,6b}=7.6 Hz, H-6a), 4.17 (1H, dd, *J*_{5,6b}=1.0 Hz, H-6b), 4.52 (1H, dd, H-5), 4.58, 4.81 (1H, 2×d, *J*_{gem}=11.9 Hz, CH₂Ph), 4.97 (1H, s, H-4), and 5.36 (1H, br.s, H-1). Found: C, 61.10; H, 6.19%. Calcd for C₁₅H₁₈O₆: C, 61.22; H, 6.17%.

1,6-Anhydro-3-O-benzyl-4-O-(*p*-methoxybenzyl)-β-D-mannopyranose (6). To a suspension of **3** (2.0 g, 5.4 mmol), powdered molecular sieves 4A (5 g), and trimethylamine–borane (1/1) (2.9 g, 40 mmol) in THF (50 cm³) was added aluminium chloride (5.3 g, 40 mmol) by portions. The mixture was stirred at room temperature for 5 h, filtered through a Celite pad, and washed with chloroform. The combined filtrate and washings were washed successively with 10% hydrochloric acid, aqueous sodium hydrogencarbonate, and brine, dried, and concentrated. Column chromatography of the residue with benzene–EtOAc (19:1, v/v) as the eluant gave 1.6 g (80%) of the 2-hydroxy derivative (**6**): $[\alpha]_{\text{D}}^{27}$ –48° (c 0.60, CHCl₃); ¹H NMR (CDCl₃) δ=3.74 (1H, s, H-4), 3.80 (1H, dd, *J*_{6a,6b}=10.1 and *J*_{5,6a}=6.0 Hz, H-6a), 3.85 (3H, s, OMe), 3.94 (1H, d, H-6b), 4.29 (1H, m, H-2), 4.55–4.70 (5H, m, 2×ArCH₂, H-5), and 5.58 (1H, br.s, H-1). Found: C, 67.63; H, 6.56%. Calcd for C₂₁H₂₄O₆: C, 67.73; H, 6.50%.

1,6-Anhydro-3,4-di-O-benzyl-β-D-mannopyranose (7). Treatment of **4** (0.30 g, 0.88 mmol) with trimethylamine–borane (1/1) (0.37 g, 5.1 mmol), aluminium chloride (0.68 mg, 5.1 mmol), and powdered molecular sieves 4A (0.6 g) in a similar way as described for the preparation of **6** afforded 0.27 g (89%) of the 3,4-di-*O*-benzyl derivative (**7**): $[\alpha]_{\text{D}}^{20}$ –57° (c 1.0, CHCl₃) (lit, $[\alpha]_{\text{D}}^{23}$ –58.5° (c 0.76, CHCl₃)).⁶⁾ ¹H NMR (CDCl₃) δ=3.01 (1H, d, *J*_{2,OH}=11.0 Hz, OH), 3.50 (1H, br.s, H-2), 3.68–3.79 (3H, m, H-3,4, 6_{exo}), 4.07 (1H, br.d, *J*_{6exo,6endo}=6.8 Hz, H-6_{exo}), 4.47–4.70 (5H, m, 2×CH₂Ph, H-5), and 5.36 (1H, br.s, H-1).

4-*O*-Acetyl-1,6-anhydro-2-azido-3-*O*-benzyl-2-deoxy- β -D-glucopyranose (8). To a solution of trifluoromethanesulfonic anhydride (3.4 cm³, 20 mmol) and pyridine (2.2 cm³, 27 mmol) in CH₂Cl₂ (100 cm³) was added a solution of **5** (4.0 g, 14 mmol) in CH₂Cl₂ (20 cm³) at -10 °C. The mixture was stirred for 1 h at 0 °C, quenched with MeOH (1 cm³), poured into ice-water, and extracted with CH₂Cl₂. The organic layer was successively washed with 10% hydrochloric acid, aqueous sodium hydrogencarbonate, and brine, dried, and concentrated. To a solution of the residual syrup in DMF (100 cm³) was added LiN₃ (6.6 g, 0.14 mol). The solution was stirred for 30 min at room temperature, diluted with water (50 cm³), and extracted with diethyl ether (3×100 cm³). The combined extracts were successively washed with water (2×100 cm³) and brine (2×100 cm³), dried (MgSO₄), and concentrated in vacuo. The residue was purified by chromatography on a silica-gel column with toluene-EtOAc (4:1, v/v) to give 3.2 g (73%) of the 2-azido-2-deoxy derivative (**8**): $[\alpha]_D^{20} +71^\circ$ (*c* 0.26, CHCl₃); ¹H NMR (CDCl₃) δ =2.14 (3H, s, OAc), 3.23 (1H, br.s, H-2), 3.56 (1H, m, H-3), 3.77 (1H, dd, *J*_{5,6a}=6.0 and *J*_{6a,6b}=7.6 Hz H-6a), 4.19 (1H, dd, *J*_{5,6}=1.0 Hz, H-6b), 4.62 (1H, br.d, H-5), 4.69 (2H, ABq, CH₂Ph), 4.77 (1H, br.s, H-4), and 5.52 (1H, br.s, H-1). Found: C, 56.34; H, 5.37; N, 13.15%. Calcd for C₁₅H₁₇O₅N₃: C, 56.42; H, 5.37; N, 13.16%.

1,6-Anhydro-2-azido-3-*O*-benzyl-2-deoxy-4-*O*-(*p*-methoxybenzyl)- β -D-glucopyranose (9). To a solution of **6** (3.90 g, 10.5 mmol) in 8:1 CH₂Cl₂-pyridine (45 cm³) was dropwise added trifluoromethanesulfonic anhydride (3.8 cm³, 23 mmol) with stirring at -18 °C. The solution was stirred at -18 °C for 40 more min, then quenched with water (20 cm³), and extracted with CH₂Cl₂. The organic layer was successively washed with 10% hydrochloric acid, aqueous sodium hydrogencarbonate, and brine, dried, and concentrated. A solution of the residual syrup (5.1 g) and LiN₃ (2 g, 41 mmol) in DMF (20 cm³) was stirred at room temperature overnight, diluted with water, and extracted with diethyl ether. The extract was washed with brine, dried, and concentrated. Column chromatography with benzene-EtOAc (24:1, v/v) as the eluant gave 3.4 g (81%) of the 2-azido-2-deoxy derivative (**9**): $[\alpha]_D^{23} +45^\circ$ (*c* 0.40, CHCl₃); ¹H NMR (CDCl₃) δ =3.27 (1H, br.s, H-2), 3.63 (1H, br.s, H-3), 3.71 (1H, t, *J*_{5,6a}=*J*_{6a,6b}=6.1 Hz, H-6a), 3.80 (3H, s, OMe), 4.00 (1H, d, H-6b), 4.45-4.58 (5H, m, 2×ArCH₂, H-5), and 5.48 (1H, br.s, H-1). Found: C, 63.61; H, 5.85, N, 10.35%. Calcd for C₂₁H₂₃O₅N₃: C, 63.47; H, 5.38; N, 10.57%.

1,6-Anhydro-2-azido-3-*O*-benzyl-2-deoxy- β -D-glucopyranose (11). **Method A:** To a solution of **8** (0.32 g, 1.0 mmol) in MeOH (10 cm³) was added 1 mol dm⁻³ methanolic sodium methoxide (0.1 cm³). The mixture was stirred at room temperature overnight, then neu-

tralized with Dowex 50W-X8 (H⁺ form), and evaporated. The residue was chromatographed on a column of silica gel with CHCl₃-MeOH (97:3, v/v) as the eluant to give 0.27 g (96%) of the 2-hydroxy derivative (**11**): $[\alpha]_D^{21} -6.2^\circ$ (*c* 0.34, CHCl₃) (lit, $[\alpha]_D^{22} -5^\circ$ (*c* 1, CHCl₃)).¹⁰⁾

Method B. To a solution of **9** (37 g, 94 mmol) in CH₂Cl₂ (650 cm³) was added water (6 cm³) and 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (33 g, 145 mmol): then the mixture was stirred at room temperature for 2 h. Aqueous sodium thiosulfate was added to the orange-colored suspension, the precipitate was filtered off and washed with CH₂Cl₂. The combined filtrate and washings were washed with aqueous sodium thiosulfate three times, dried, and evaporated. The residue was purified by column chromatography with CHCl₃-MeOH (97:3, v/v) as the eluant to give 24.2 g (93%) of the 2-hydroxy derivative (**11**).

We express our gratitude to Ms. Mutsuko Yoshida and her collaborators for the elemental analyses, and to Dr. Jun Uzawa and Mrs. Tamiko Chijimatsu for recording and measuring the ¹H NMR spectra.

References

- 1) H. Paulsen, *Chem. Soc. Rev.*, **13**, 15 (1984); J. Banoub, P. Boullanger, and D. Lafont, *Chem. Rev.*, **92**, 1167 (1992).
- 2) T. Trnka and M. Cerny, *Coll. Czech. Chem. Commun.*, **36**, 2216 (1971); M. Cerny, J. Pacak, and J. Stanek, *Coll. Czech. Chem. Commun.*, **30**, 1151 (1965); H. Paulsen, H. Koebernick, W. Stenzel, and P. Nöll, *Tetrahedron Lett.*, **1975**, 1493.
- 3) M. Kloosterman, M. P. de Niji, and J. H. van Boom, *J. Carbohydr. Chem.*, **5**, 215 (1986).
- 4) M. Ek, P. J. Garegg, H. Hultberg, and S. Oscarson, *J. Carbohydr. Chem.*, **2**, 305 (1983).
- 5) Y. Ishido and N. Sakairi, *Carbohydr. Res.*, **97**, 151 (1981).
- 6) F. Dasgupta and P. J. Garegg, *Synthesis*, **1988**, 626.
- 7) F. H. Cano, C. Foces-Foces, J. Jimenez-Barbero, M. Bernabe, and M. Martin-Lomas, *Carbohydr. Res.*, **155**, 1 (1986).
- 8) T. Uryu, K. Hatanaka, Y. Sakamoto, K. Harima, S. Hagino, K. Ito, A. Funamoto, and K. Matsuzaki, *Nippon Kagaku Kaishi*, **1982**, 1638.
- 9) H. Hori, Y. Nishida, H. Orui, and H. Meguro, *J. Org. Chem.*, **54**, 1346 (1989).
- 10) H. Paulsen and W. Stenzel, *Chem. Ber.*, **111**, 2348 (1978).
- 11) N. Sakairi, L.-X. Wang, and H. Kuzuhara, *J. Chem. Soc., Chem. Commun.*, **1991**, 289.