

Asymmetric Synthesis of Functionalized Secondary Alcohols by Catalytic Ring-Cleavage Reactions of Cyclic Acetals Derived from (*R*)-1,3-Butanediol

Motoharu Kinugasa, Toshio Harada,* Takayuki Egusa, Katsuhiro Fujita, and Akira Oku

Department of Chemistry, Kyoto Institute of Technology, Sakyo-ku, Kyoto 606

(Received July 22, 1996)

In the presence of a catalytic amount (0.1—0.2 molar amount) of a 2-phenyl-1,3,2-oxazaborolidin-5-one, prepared by the reaction of dichlorophenylborane and *N*-(trifluoromethylsulfonyl)-*L*-phenylalanine, and enol silyl ethers ((*R*²)₂C=C-(*R*³)OTMS; *R*³ = *t*-BuS, EtS, EtO, Ph) (1.1–1.5 molar amount), chiral cyclic acetals **6** derived from (*R*)-1,3-butanediol and aldehydes undergo an efficient ring-cleavage reaction with the inversion of the stereochemistry at the acetal carbon to give the *anti* isomer of the corresponding products with high stereoselectivity. The reaction is applicable to acetals derived from aromatic, aliphatic, and α,β -unsaturated aldehydes. Not only enol silyl ethers, but also methyltrimethylsilane and allyltributyltin, can be employed as nucleophiles, leading to the selective formation of the *anti* isomer of the corresponding allylated ring-cleavage products. Removal of the chiral auxiliary from these ring-cleavage products by a two-step sequence ((i) PCC (ii) Bn₂NH₂(CF₃CO₂)) furnishes enantiomerically enriched β -hydroxy carbonyl compounds and homoallyl alcohols. A modest level of kinetic resolution is observed in the ring cleavage of a racemic acetal catalyzed by a phenylboron compound derived from *N*-mesyl-*L*-phenylalanine.


The Lewis acid-mediated cleavage of chiral cyclic acetals has been intensively investigated in recent years, and has proven to be a useful tool in asymmetric synthesis.¹⁾ A TiCl₄-mediated reaction of chiral acetals **1a**, derived from enantiomerically pure 2,4-pentanediol, proceeds with a stereoselective introduction of a variety of carbon nucleophiles in an invertive manner, affording a powerful method for asymmetric carbon–carbon bond formation (Scheme 1-a).^{2,3)} Acetals **1b** in which inexpensive and readily available 1,3-butanediol⁴⁾ is used as a chiral auxiliary also serve as substrates, exhibiting high stereoselectivity.⁵⁾ The TiCl₄-medi-

ated cleavage reaction is also utilized in an enantiodifferentiating transformation of prochiral diols (Scheme 1-b).⁶⁾ The cleavage of the equatorial C–O bond of *l*-menthone-derived spiroacetals **2** proceeds exclusively, leading to the formation of protected derivatives of the prochiral diols in an enantiodifferentiating manner.


In both types of ring-cleavage reactions, either a stoichiometric or excess amount of Lewis acids is required to obtain high diastereoselectivity.^{7,8)} It has been reported that the use of a 5.5 molar amount (abbrev. to mol. amt. hereafter) of “Ti-blend”, a 6 : 5 mixture of TiCl₄ and Ti(OPr^{*i*})₄, gave a higher selectivity in the reaction shown in Scheme 1-a.⁹⁾ A search for efficient Lewis acid complexes which catalyze the cleavage of cyclic acetals is important not only for preparative uses of the reaction, but also for the future development of an enantioselective ring cleavage without the aid of chiral auxiliaries.

With enol silyl ethers as nucleophiles, the Lewis acid (ML_{*n*}) mediated reaction of acetal **1** gives zwitterion **3** as an intermediate (Scheme 2). The migration of the silyl group to the oxygen atom of the ring-cleavage product with a regeneration of ML_{*n*} should be crucial for achieving an efficient catalytic cycle. It has recently been reported that 1,3,2-oxazaborolidin-5-ones derived from *N*-sulfonyl α -amino acids¹⁰⁾ act as effective catalysts for the mechanistically related asymmetric Mukaiyama aldol reaction.¹¹⁾ The facile migration of the silyl group observed there prompted us to investigate the ring cleavage mediated by the oxazaborolidinones.¹²⁾

We now report that oxazaborolidinones **5** catalyze the ring



Scheme 1.




Scheme 2.



cleavage of chiral acetals derived from (*R*)-1,3-butanediol in a stereoselective manner. A wide applicability of the catalytic reaction to a variety of acetals and nucleophilic reagents is described.¹³⁾

Results and Discussion

Organoboron compounds **5a–f** were prepared simply by treatment of the corresponding sulfonamides of L-amino acids (**4a–f**) with an equimolar amount of dichlorophenylborane in CH₂Cl₂ and a subsequent removal of produced HCl and the solvent in vacuo (Eq. 1). ¹H and ¹¹B NMR analyses of the residue showed a clean formation of **5a–f**. Alkylboron compounds derived from *N*-sulfonyl amino acids have been prepared by the dehydration of alkylboronic acids.¹⁴⁾ However, the reaction is sometimes sluggish and phenylboron compounds, in our hands, could not be prepared by this method. The reaction using dichlorophenylborane proceeded cleanly in a facile manner. Methylboron compound **5g** was also prepared by using dibromomethylborane (Eq. 2). Oxazaborolidinone **5h** was prepared by the reaction of BH₃–THF and **4d**.¹⁰⁾





4a; R¹ = H, R² = *p*-Tol
4b; R¹ = Bn, R² = *p*-Tol
4c; R¹ = Bn, R² = Me
4d; R¹ = Bn, R² = CF₃
4e; R¹ = *i*-Pr, R² = *p*-Tol
4f; R¹ = *i*-Pr, R² = CF₃



Ring-cleavage reaction of chiral acetal (2*S*,4*R*)-**6a** with

enol silyl ether **7a** (3 mol. amt.) derived from thioic *S*-ester was examined in the presence of oxazaborolidinones **5** in CH₂Cl₂ at rt (Eq. 3 and Table 1). In most reactions, a GC analysis of the crude reaction mixture showed the formation of silyl ether derivative **8a** as the sole product, suggesting a smooth migration of the silyl group. Because the hydrolysis of the silyl group took place in part during isolation by silica gel chromatography, the crude mixture was treated with Bu₄NF, and the product was isolated as alcohol **9a**.¹⁵⁾



(3)

The reaction using phenylboron compound **5a** derived from *N*-tosylglycine proceeded stereoselectively to give *anti*-**9a**, but in low yield (Entry 1). The reaction is also highly regioselective; no product derived from a C(2)–O(3) bond cleavage was detected. The yield of **9a** was improved considerably by using of valine- and phenylalanine-derived complexes **5b,e** (Entries 2 and 6). As shown in Entries 4–6, the structure of the sulfonamide moiety is also influential. The most satisfactory result was obtained when phenylboron compound **5d**, derived from *N*-(trifluoromethylsulfonyl)phenylalanine, was used: Even when a 10 mol. amt. of the catalyst was employed with a 1.5 mol. amt. of the nucleophile, the ring-cleavage product was obtained in high yield with a high stereoselectivity of 98 : 2 (Entry 4).

Although we used enantiomerically pure oxazaborolidinones prepared from L-amino acids, chirality of the catalyst complex is not an important factor for the *anti* : *syn* selectivity. Indeed, the reaction of racemic acetal *rac*-**6a** in the presence of **5d** afforded **9a** similarly with high stereoselectivity (Entry 5). It should be noted that the phenyl group attached to the boron atom is essential to the catalytic activity. Thus, only a stoichiometric or less efficient reaction took place when methylboron compound **5g** or *B*-unsubstituted compound **5h** was used, respectively (Entries 7 and 8). Dichlorophenylborane, itself, also did not catalyze the ring-cleavage reaction (Entry 9). For a comparison, we also examined the use of TMSOTf⁸⁾ as a catalyst (Entry 10). The nonstereoselective reaction observed here excludes the possibility of a ring cleavage involving a silicon-catalyzed mechanism.¹²⁾

Table 1. Reaction of Acetal (2*S*,4*R*)-**6a** and Enol Silyl Ether **7a** in the Presence of Oxazaborolidinones **5**^{a)}


Entry	Catalyst	Time (h)	Yield of 9a (%)	<i>anti</i> : <i>syn</i> ^{b)}	Recovery of 6a (%)
1 ^{c)}	5a	21	31	94 : 6	55
2	5b	21	52	96 : 4	32
3	5c	20	94	94 : 6	—
4 ^{d)}	5d	4	81	98 : 2	9
5 ^{c,d)}		1	80	97 : 3	—
6	5e	20	81	96 : 4	13
7	5g	20	15	93 : 7	76
8	5h	21	Trace	—	97
9 ^{c)}	PhBCl ₂	23	14	63 : 37	63
10 ^{c)}	TMSOTf	20	74	60 : 40	15

a) Unless otherwise noted, reactions were carried out by using 3 mol. amt. of **7a** and 0.2 mol. amt. of oxazaborolidinones **5** at rt. b) Ratios were determined by ¹H NMR (300 MHz) analyses. c) Racemic acetal *rac*-**6a** was used. d) 1.5 mol. amt. of **7a** and 0.1 mol. amt. of **5d** were used.

In the reaction using oxazaborolidinones **5b—e**, the ring-cleavage products existed as trimethylsilyl ether **8a** through the course of the reaction, judging from GC analyses of the reaction mixture. This observation implies that, in the present catalytic system, the intermediate **10** undergoes a facile silyl-group migration to give **8a**. On the other hand, when glycine-derived oxazaborolidinone **5a** was employed, the formation of **8a** as well as that of **9a** was observed during the early stage of the reaction. After 20 h of the reaction time, **8a** was the sole product and no **9a** was detected. The observed dependence of the rate of silyl-group migration on the structures of the catalysts can be rationalized if we assume that the migration proceeds via the trimethylsilyl ester derivative **11** (Scheme 3).¹⁶⁾ Thus, the substituent R originated from the amino acid derived moiety may enhance the rate for the regeneration of the catalyst along with a simultaneous release of the silylated product **8** by a Thorpe–Ingold effect.¹⁷⁾


We then examined the ring-cleavage reaction of several acetals and enol silyl ethers using a 0.1–0.3 mol. amt. of the optimal catalyst **5d** (Eq. 4 and Table 2). Not only the enol silyl ethers of thioic *S*-esters, but also those of an ester and a ketone, can be employed as nucleophiles. Thus, the ring cleavage of (2*S*,4*R*)-**6a** with **7c,d** proceeded with high stereoselectivity to give the corresponding products **9c,d** (Table 2,

Entries 3 and 4). Chiral acetals (2*S*,4*R*)-**6b—d**, derived from aliphatic and α,β -unsaturated aldehydes, also underwent a stereoselective ring-cleavage reaction (Entries 4–10). For these substrates, the stereoselectivities of the reaction at rt were lower than that of aromatic acetal (2*S*,4*R*)-**6a**. However, a satisfactory result was obtained when the reactions were carried out at –40 °C using a 0.2–0.3 mol. amt. of **5d**.










(4)

In order to extend the scope of the catalytic ring cleavage to the allylation of cyclic acetals, reactions using allylic silanes **12a,b** as nucleophiles were examined (Eq. 5). In the presence of **5d** (0.2 mol. amt.), the reaction of allylsilane **12a** and methallylsilane **12b** proceeded smoothly to give allylation products **13a** and **13b**, respectively, in high yields (Table 3, Entries 1 and 3). However, the level of the stereoselectivity was lower than those observed for the enol silyl ethers. A brief survey of other oxazaborolidinones as well as the reaction conditions showed that the most satisfactory results were obtained when catalyst **5d** was used in the reaction of **12b** at –50 °C (Entry 4). Tosyl and mesyl derivatives **5b,c** exhibited a higher stereoselectivity (Entries 8–11). However, these reactions did not reach completion. Triflates **5d,f** of higher Lewis acidity catalyzed the ring cleavage with **12b** at appreciable rate, even at –50 °C (Entries 4 and 12).



Scheme 3.

Table 2. Catalytic Ring-Cleavage of Acetals (2*S*,4*R*)-**6a**—**d** with Enol Silyl Ethers **7a**—**c**^{a)}

Entry	Acetal	Nucleophile	5d (mol. amt.)	Temp °C	Time h	Product	Yield %	<i>anti</i> : <i>syn</i> ^{b)}
1	6a	7a	0.1	R.T.	4		81	98 : 2
2		7b	0.1	R.T.	4		78	98 : 2
3		7c	0.2	-20	1		88	>99 : 1
4 ^{c)}		7d	0.1	R.T.	3		86	93 : 7
5	6b	7a	0.3	-40	22		88	88 : 12
6			0.2	R.T.	18		74	75 : 25
7	6c	7a	0.3	-40	4		90	93 : 7 ^{d)}
8			0.2	R.T.	4		88	79 : 21 ^{d)}
9	6d	7a	0.2	-40	2		71	96 : 4 ^{e)}
10			0.1	R.T.	4		80	93 : 7 ^{e)}








a) Unless otherwise noted, reactions were performed by using 1.5 mol. amt. of a nucleophile. b) Unless otherwise noted, ratios were determined by ¹H NMR (300 MHz) analyses. c) 1.1 mol. amt. of a nucleophile was used. d) The ratio was determined by a capillary GC analysis. e) The ratio was determined by the ¹H NMR (300 MHz) analysis of the aldehyde derivative **14g**.


Table 3. Reaction of Acetal **6a** and Allylic Silanes **12a,b** in the Presence of Oxazaborolidinones^{a)}

Entry	Catalyst	Allylic silane	Temp (°C)	Time (h)	Product	Yield (%)	<i>anti</i> : <i>syn</i> ^{b)}
1	5d	12a	R.T.	2	13b	56	64 : 36
2			-20	3		77	71 : 29
3		12b	R.T.	3		89	71 : 29
4			-50	3		91	83 : 17
5 ^{c)}	5b		-50	3		86	92 : 8
6 ^{d)}			-50	4		92	95 : 5
7 ^{e)}			-50	6		90	90 : 10
8			R.T.	21		20	91 : 9
9	5c		-50	6		31	92 : 8
10			R.T.	22		49	87 : 13
11	5f		-50	6		18	90 : 10
12			-50	3		90	85 : 15

a) Unless otherwise noted, reactions were carried out by using 0.2 mol. amt. of **5** in CH₂Cl₂. (2*S*,4*R*)-**6a** and *rac*-**6a** were used in Entries 1—4 and 5—12, respectively. b) Ratios were determined by ¹H NMR and capillary GC analyses for **13a** and **13b**, respectively. c) 1.0 mol. amt. of **5d** was used. d) **6a** was added in two portions (0.5 mol. amt. each) with time interval of 1 h. e) **6a** was added slowly by using syringe pump during 5 h and the mixture was stirred for 1 h.

Table 4. Catalytic Ring-Cleavage of Acetals **6** with Methallylsilane **12b**^{a)}

Entry	Acetal	Time (h)	Product	Yield (%)	<i>anti</i> : <i>syn</i> ^{b)}
1	<i>rac</i> - 6a	3	 13b	91	83 : 17 ^{c)}
2	(2 <i>S</i> ,4 <i>R</i>)- 6b	7	 13c	44	87 : 13
3	<i>rac</i> - 6c	8	 13d	62	87 : 13
4	(2 <i>S</i> ,4 <i>R</i>)- 6d	5	 13e	80	92 : 8
5 ^{d)}	(2 <i>S</i> ,4 <i>R</i>)- 6e	6	 13f	95	92 : 8
6	<i>rac</i> - 6f	3	 13g	90	82 : 18
7 ^{d)}	<i>rac</i> - 6g	6	 13h	81	94 : 6

a) Unless otherwise noted, reactions were carried out by using 0.2 mol. amt. of **5d** in CH₂Cl₂ at -50 °C.b) Unless otherwise noted, ratios were determined by ¹H NMR analyses. c) The ratio was determined by a capillary GC analysis. d) Acetal **6e** or **6g** was added slowly by using syringe pump during 5 h and the mixture was stirred for 1 h.

When a stoichiometric amount of **5d** was used in the reaction of **6a** and **12b** at -50 °C, the *anti* isomer of **13b** was obtained with higher stereoselectivity (*anti* : *syn* = 92 : 8) (Entry 5). Interestingly, a similar level of high stereoselectivity (95 : 5) was also observed when **6a** was added in two portions (0.5 mol. amt. each) with a time interval of 4 h to a solution of 0.5 mol. amt. of **5d** (Entry 6). Of preparative interest, a 90 : 10 selectivity was obtained by the slow addition of acetal **6a** using a syringe-pump for 5 h (Entry 7). These results suggest the concomitant formation of a less stereoselective catalyst during the reactions.¹⁸⁾ Although the cat-

alytic activity of this species would be lower, it may catalyze the nonstereoselective reaction if a relatively large amount of **6a** is present in the reaction mixture. By a portionwise or slow addition of **6a**, the undesirable pathway would be considerably retarded.

By using catalyst **5d** at -50 °C, a variety of acetals **6** underwent ring-cleavage reactions to give the allylation product **13** with *anti* selectivity (Eq. 6, Table 4). The reaction of acetals **6a,d–g**, derived from aromatic and α,β -unsaturated aldehydes, proceeded efficiently. On the other hand, the yields of the ring-cleavage products for aliphatic acetals **6b,c** were moderate.

















Table 5. Asymmetric Synthesis of Functionalized Alcohols **15** and **17**

Entry	Ring-cleavage product (<i>anti</i> : <i>syn</i>)	Aldehyde	Yield %	Product	Yield %	$[\alpha]_D^{25a)}$ (<i>c</i> , solvent)	Absolute configuration
1	9a (98 : 2)	14a	84	 15a	80	−43.4 (2.15, benzene)	<i>S</i>
2	9b (98 : 2)	14b	96	 15b	85	−60.7 (2.00, benzene)	<i>S</i>
3	9c (>99 : 1)	14c	81	 15c	96	−4.7 (0.96, MeOH) ^{b)}	<i>R</i>
4	9d (93 : 7)	14d	78	 15d	67	−27.9 (0.980, MeOH)	<i>S</i>
5	9e (88 : 12)	14e	82	 15e	69	−9.24 (2.51, benzene)	<i>R</i>
6	9f (93 : 7)	14f	79	 15f	84	−23.5 (1.50, CHCl ₃) ^{b)}	<i>S</i>
7	9g (96 : 4)	14g	83	 15g	71	+18 (0.57, MeOH) ^{b)}	<i>S</i>
8	13a (94 : 6)	16a	90	 17a	63	−41 (0.26, benzene)	<i>S</i>
9	13b (83 : 17)	16b	84	 17b	80	−38.1 (1.97, benzene)	<i>S</i>
10	13e (>99 : 1)	16c	66	 17c	89	−1.48 (2.43, Et ₂ O)	<i>S</i>
11	13f (92 : 8)	16d	76	 17d	98	−41.0 (2.01, benzene)	<i>S</i> ^{c)}
12	13h (94 : 6)	16e	56	 17e	77	—	<i>S</i> ^{c)}


a) Specific rotation of **15** or **17** unless otherwise noted. b) Specific rotation was measured after converting into a carboxylic acid derivative.
c) Tentative assignment.

In comparison with methallylsilane **12b**, allylsilane **12a** showed a lower stereoselectivity even when the reaction was performed at -20°C (Table 3, Entry 2). However, it was found that the ring cleavage with allylation was achieved with high stereoselectivity by using more nucleophilic allyl-tributyltin; the *anti* isomers of **13a,i** were obtained with high stereoselectivity (94 : 6) in the reaction of **6a,d** by using a 0.5 mol. amt. of **5d** (Eq. 7).



The chiral auxiliary of ring-cleavage products **9a–h** can be readily removed by a two-step sequence without any loss of stereochemical integrity (Eq. 8). Thus, pyridinium chlorochromate (PCC) oxidation of **9** and treatment of the result-


ing aldehydes **14** with dibenzylammonium trifluoroacetate in benzene¹⁹⁾ furnished the corresponding β -hydroxy carbonyl compounds **15** of high enantiomeric purities (Table 5, Entries 1–7). The ee value of **15a**, determined by converting into a (*S*)- α -methoxy- α -(trifluoromethyl)phenylacetic acid (MTPA) ester derivative, was 93%, being consistent with the *anti/syn* ratios of the ring-cleavage products **9a**. Similarly, homoallyl alcohols (*S*)-**17** were prepared starting from the allylation products **13** (Entries 8–14).



The absolute configurations of **15a,b,d,e**^{20,21)} and **17a,b,c**^{22,23)} were determined by a comparison with the literature values. For **15c,f,g**, the absolute configurations were determined after converting them into the known β -hydroxy carboxylic acids.^{24–26)}

The ring cleavage of chiral acetals (**2S,4R**)-**6a–d** took place exclusively at the less hindered C(2)–O(1) bonds with a stereoselective introduction of nucleophiles with an inversion of the C(2S) stereochemistry. These selectivities are well-consistent with the previously reported results on the TiCl₄-promoted ring-cleavage reaction. In spite of the use of chiral oxazaborolidinones **5** derived from enantiomerically pure L-amino acids, the chirality of the catalysts may not be an important factor for the *anti*:*syn* selectivity, because the selectivity is determined by the invertive vs retentive mode of the ring-cleavage reaction.

However, when racemic acetals are used as substrates and the reaction is quenched before a full conversion, the chirality of the catalysts should be of importance because one can expect differences in the reaction rates between enantiomeric acetals. Indeed, an appreciable degree of enantiodifferentiation was observed in the reaction of *rac*-**6a**. Thus, when the reaction of *rac*-**6a** and **7a** (3 mol. amt.) using oxazaborolidinone **5c** (0.2 mol. amt.) was quenched before a full conversion (r.t., 3 h), acetal (**2R,4S**)-**6a** of 51% ee was recovered in 27% yield together with **9a** (*anti*:*syn*=92:8, 69%) (Eq. 10).



In summary, we have shown that *B*-phenyl oxazaborolidinones **5**, derived from sulfonamides of α -amino acids serve as efficient catalysts for the cleavage of cyclic acetals with nucleophiles. Functionalized secondary alcohols **15** and **17** of high enantiomeric purities were obtained via the catalytic cleavage reaction of chiral acetals **6** derived from (*R*)-1,3-butanediol. Although the degree of selectivity was not satisfactory, the observed enantiodifferentiation of racemic acetal *rac*-**6a** suggests the possibility of catalytic asymmetric ring cleavage of achiral cyclic acetals using proper oxazaborolidinones.

Experimental

General. Unless otherwise noted, the ¹H, ¹¹B, and ¹³C NMR spectra were recorded in CDCl₃ at 300, 96.5, and 75.6 MHz, respectively. ¹¹B NMR chemical shifts were recorded in units of parts per million relative to the external standard BF₃·Et₂O. Microanalyses were performed at the Microanalysis Center of Kyoto University. Wakogel C-300 was used for flash chromatography. An enantiomeric excess of acetal **6a** was established by a capillary GC analysis using a chiral CP-Cyclodextrin-B-236-M-19 (25 m) column. Unless otherwise specified, all of the organic extracts were dried over Na₂SO₄. THF and CH₂Cl₂ were used after distillation from sodium benzophenone ketyl and from CaH₂, respectively. (*R*)-1,3-Butanediol was purchased from Wako Pure Chemical Industries

Ltd.. Enol silyl ethers **7a–d**²⁷⁾ and methallylsilane **12b**²⁸⁾ were prepared according to the literature.

Acetals **6a–d,f,g** were prepared by condensation of the corresponding aldehydes and (*R*)-1,3-butanediol.¹⁴⁾ Acetal **6e** was prepared by treatment of cinnamaldehyde with the bis(TMS) ether derivative of (*R*)-1,3-butanediol in the presence of TMSOTf.²⁹⁾ The spectral data of the acetals are as follows.

(2S,4R)-4-Methyl-2-phenyl-1,3-dioxane (6a): ¹H NMR δ = 1.33 (3H, d, *J* = 6.2 Hz), 1.54 (1H, m), 1.82 (1H, m), 3.91–4.04 (2H, m), 4.27 (1H, ddd, *J* = 1.3, 5.0, and 11.4 Hz), 5.52 (1H, s), 7.35 (3H, m), 7.54 (2H, m); ¹³C NMR δ = 21.71, 32.90, 66.94, 73.28, 101.22, 126.02, 128.10, 128.57, 138.77; IR (liquid film) 1100, 750, 695 cm^{−1}.

(2S,4R)-4-Methyl-2-(2-phenylethyl)-1,3-dioxane (6b): ¹H NMR δ = 1.25 (3H, d, *J* = 6.0 Hz), 1.44 (1H, m), 1.68 (1H, m), 1.94 (2H, m), 2.75 (2H, m), 3.67–3.78 (2H, m), 4.11 (1H, ddd, *J* = 1.4, 5.0, and 11.4 Hz), 4.51 (1H, t, *J* = 5.3 Hz), 7.16–7.31 (5H, m); ¹³C NMR δ = 21.7, 30.2, 33.1, 36.4, 66.6, 72.7, 101.1, 125.7, 128.3, 128.4, 141.8; IR (film) 1130, 700 cm^{−1}.

(2S,4R)-2-Cyclohexyl-4-methyl-1,3-dioxane (6c): ¹H NMR δ = 0.96–1.83 (15H, m including d (3H, *J* = 6.3 Hz) at 1.20), 3.63–3.74 (2H, m), 4.08 (1H, ddd, *J* = 1.2, 5.1, and 11.4 Hz), 4.21 (1H, d, *J* = 5.7 Hz); ¹³C NMR δ = 20.8, 24.8 (2C), 25.5, 26.4, 26.7, 32.3, 41.5, 65.4, 71.7, 104.1; IR (liquid film) 1120 cm^{−1}.

(2S,4R)-2-(2-phenylethenyl)-1,3-dioxane (6d): Bp (Kugelrohr) 97–100 °C/0.4 mmHg (1 mmHg = 133.322 Pa); ¹H NMR δ = 1.03 (3H, d, *J* = 6.0 Hz), 1.49 (1H, m), 1.74 (1H, m), 3.82–3.94 (2H, m), 4.19 (1H, ddd, *J* = 1.4, 5.0, and 11.4 Hz), 5.16 (1H, dd, *J* = 1.2 and 4.8 Hz), 6.21 (1H, dd, *J* = 4.8 and 16.2 Hz), 6.78 (1H, d, *J* = 16.2 Hz), 7.21–7.42 (5H, m); ¹³C NMR δ = 21.7, 33.0, 66.7, 73.0, 100.5, 125.9, 126.8, 128.0, 128.4, 133.2, 136.2; IR (film) 1670, 960, 970 cm^{−1}.

(2S,4R)-2-(4-Methoxyphenyl)-4-methyl-1,3-dioxane (6e): ¹H NMR δ = 1.30 (3H, d, *J* = 6.0 Hz), 1.52 (1H, m), 1.79 (1H, m), 3.79 (3H, s), 3.88–4.05 (2H, m), 4.23 (1H, ddd, *J* = 1.1, 5.0, and 11.3 Hz), 5.46 (1H, s), 6.88 (2H, m), 7.42 (2H, m); ¹³C NMR δ = 21.8, 32.9, 55.2, 67.0, 101.2, 113.6, 127.3, 131.4, 159.8; IR (liquid film) 1110, 1030, 830 cm^{−1}.

(2S,4R)-2-(2-Methoxyphenyl)-4-methyl-1,3-dioxane (6f): ¹H NMR δ = 1.30 (3H, d, *J* = 6.21 Hz), 1.52 (1H, m), 1.81 (1H, m), 3.83 (3H, s), 3.92–4.05 (3H, m), 4.22 (1H, br dd, *J* = 5.0 and 11.4 Hz), 5.90 (1H, s), 6.86 (1H, d, *J* = 8.3 Hz), 6.98 (1H, br t, *J* = ca. 8 Hz), 7.29 (1H, br t, *J* = ca. 8 Hz), 7.64 (1H, dd, *J* = 1.7 and 7.2 Hz); ¹³C NMR δ = 21.9, 33.1, 55.6, 67.2, 73.5, 96.4, 110.7, 120.8, 127.1, 127.3, 129.9, 156.4;

(2S,4R)-2-(2-Furyl)-4-methyl-1,3-dioxane (6g): ¹H NMR δ = 1.31 (3H, d, *J* = 6.3 Hz), 1.52 (1H, m), 1.85 (1H, m), 3.87–4.04 (2H, m), 4.23 (1H, ddd, *J* = 1.5, 5.0, and 11.6 Hz), 5.58 (1H, s), 6.35 (1H, dd, *J* = 1.8 and 3.2 Hz), 6.44 (1H, br d, *J* = 3.2 Hz), 7.39 (1H, dd, *J* = 0.8 and 1.8 Hz); ¹³C NMR δ = 21.6, 32.8, 67.0, 73.5, 95.9, 107.3, 110.0, 142.3, 151.1; IR (liquid film) 1100, 1060, 1030, 740 cm^{−1}.

N-Tosyl- and *N*-mesyl amino acids **4a–c,e** were prepared by the reaction of the corresponding amino acids with *p*-toluenesulfonyl chloride and methanesulfonyl chloride, respectively, under Schotten–Baumann conditions.^{10,30,31)}

***N*-Tosylglycine (4a):**³⁰⁾ Mp 149 °C (recrystallized from H₂O); ¹H NMR (acetone-*d*₆) δ = 2.39 (3H, s), 6.62 (1H, s), 3.75 (2H, d, *J* = 5.7 Hz), 7.36 (2H, d, *J* = 8.1 Hz), 7.74 (2H, d, *J* = 8.1 Hz).

***N*-Tosyl-L-phenylalanine (4b):**³⁰⁾ Mp 161–163 °C (recrystallized from MeOH and H₂O); ¹H NMR δ = 2.39 (3H, s), 3.00 (1H, dd, *J* = 6.3 and 13.9 Hz), 3.09 (1H, dd, *J* = 5.5 and 13.9

Hz), 4.17—4.24 (1H, m), 5.03 (1H, d, $J=8.5$ Hz), 7.05—7.09 (2H, m), 7.20—7.26 (5H, m), 7.59 (2H, d, $J=8.4$ Hz).

N-Mesyl-L-phenylalanine (4c):³¹⁾ Mp 104—106 °C (recrystallized from benzene and ethyl acetate); ¹H NMR $\delta=2.65$ (3H, s), 3.03 (1H, dd, $J=7.8$ and 13.8 Hz), 3.25 (1H, dd, $J=4.8$ and 13.8 Hz), 4.45 (1H, ddd, $J=4.8$, 7.8, and 9.3 Hz), 4.95 (1H, d, $J=9.3$ Hz), 7.23—7.37 (5H, m).

N-Tosyl-L-valine (4e):³⁰⁾ Mp 149 °C (MeOH and H₂O) (recrystallized from MeOH and H₂O); ¹H NMR $\delta=0.87$ (3H, d, $J=6.9$ Hz), 0.96 (3H, d, $J=6.9$ Hz), 2.05—2.15 (1H, m), 2.41 (3H, s), 3.79 (1H, dd, $J=4.5$ and 12.9 Hz), 5.07 (1H, m), 7.28 (2H, d, $J=8.1$ Hz), 7.72 (2H, d, $J=8.1$ Hz).

N-Trifluoromethylsulfonyl amino acids **4d,f** were prepared by the reaction of the methyl esters of the corresponding amino acids with trifluoromethanesulfonyl chloride in the presence of *i*-Pr₂NEt and subsequent saponification.^{10b)}

N-(Trifluoromethylsulfonyl)-L-phenylalanine (4d): Mp 121—122 °C (recrystallized from benzene); ¹H NMR $\delta=3.12$ (1H, dd, $J=5.7$ and 13.8 Hz), 3.26 (1H, dd, $J=5.4$ and 13.8 Hz), 4.57—4.64 (1H, m), 5.33 (1H, m), 7.17—7.20 (2H, m), 7.33—7.36 (3H, m).

N-(Trifluoromethylsulfonyl)-L-valine (4f): Mp 145—146 °C (recrystallized from ether); ¹H NMR (acetone-*d*₆) $\delta=0.98$ (3H, d, $J=6.9$ Hz), 1.05 (3H, d, $J=6.9$ Hz), 2.22—2.33 (1H, m), 3.99 (1H, d, $J=5.1$ Hz).

General Method for the Preparation of Oxazaborolidinones 5a—f. To a solution of sulfonamide **4a—f** (0.10 mmol) in CH₂Cl₂ (1 mL) at rt under nitrogen was added dichlorophenylborane (13 μ L, 0.10 mmol). After being stirred for 30 min the mixture was concentrated in vacuo to give phenylboron compounds **5a—f** as a white solid, which was then dissolved in CH₂Cl₂ (0.2 mL) and used for ring-cleavage reactions.

Oxazaborolidinone 5a: ¹H NMR $\delta=2.42$ (3H, s), 4.24 (2H, s), 7.31 (2H, d, $J=8.1$ Hz), 7.45—7.52 (2H, m), 7.57—7.63 (1H, m), 7.70 (2H, d, $J=8.1$ Hz), 8.16 (2H, d, $J=6.9$ Hz); ¹¹B NMR $\delta=46.5$.

Oxazaborolidinone 5b: ¹H NMR $\delta=2.39$ (3H, s), 3.37 (1H, dd, $J=2.7$ and 14.1 Hz), 3.69 (1H, dd, $J=5.4$ and 14.1 Hz), 4.55 (1H, dd, $J=2.7$ and 5.4 Hz), 7.20—7.39 (8H, m), 7.46—7.55 (4H, m), 7.78—7.82 (2H, m); ¹¹B NMR $\delta=42.9$.

Oxazaborolidinone 5c: ¹H NMR $\delta=3.01$ (3H, s), 3.43 (1H, dd, $J=3.0$ and 14.1 Hz), 3.57 (1H, dd, $J=5.4$ and 14.1 Hz), 4.85 (1H, dd, $J=3.0$ and 5.4 Hz), 7.20—7.33 (5H, m), 7.37—7.43 (2H, m), 7.51—7.57 (1H, m), 7.90—7.93 (2H, m); ¹¹B NMR $\delta=47.1$.

Oxazaborolidinone 5d: ¹H NMR $\delta=3.46$ (1H, dd, $J=2.4$ and 14.4 Hz), 3.62 (1H, dd, $J=5.7$ and 14.4 Hz), 4.89 (1H, qdd, $J=0.6$, 2.4, and 5.7 Hz), 7.21—7.38 (6H, m), 7.48—7.56 (2H, m), 7.70—7.74 (2H, m); ¹¹B NMR $\delta=48.3$.

Oxazaborolidinone 5e: ¹H NMR $\delta=0.89$ (3H, d, $J=6.9$ Hz), 1.23 (3H, d, $J=6.9$ Hz), 2.38 (3H, s), 2.59—2.75 (1H, m), 4.26 (1H, d, $J=3.6$ Hz), 7.20—7.26 (2H, m), 7.42—7.62 (5H, m), 8.06—8.10 (2H, m); ¹¹B NMR $\delta=37.4$.

Oxazaborolidinone 5f: ¹H NMR $\delta=0.97$ (3H, d, $J=6.9$ Hz), 1.27 (3H, d, $J=6.9$ Hz), 2.55—2.67 (1H, m), 4.53 (1H, d, $J=3.6$ Hz), 7.45—7.66 (3H, m), 8.05—8.08 (2H, m); ¹¹B NMR $\delta=36.9$.

Oxazaborolidinone 5g: The complex was prepared by the reaction of **4d** with MeBBR₂ according to a procedure similar to that described above. **5g:** ¹H NMR $\delta=0.60$ (3H, s), 3.35 (1H, dd, $J=2.4$ and 14.1 Hz), 3.54 (3H, dd, $J=5.7$ and 14.1 Hz), 4.65 (1H, dd, $J=2.4$ and 5.7 Hz), 7.17—7.20 (2H, m), 7.30—7.35 (3H, m); ¹¹B NMR $\delta=40.4$.

S-*t*-Butyl (3S)-3-[(1R)-3-Hydroxy-1-methylpropoxy]-3-phenylpropanethioate (9a) (Representative Procedure for Catalytic

Ring Cleavage of Acetals 6 with Enol Silyl Ethers 7): To a stirred solution of acetal **6a** (71.8 mg, 0.40 mmol) and **7a** (122 mg, 0.60 mmol) in CH₂Cl₂ (0.32 mL) at rt under argon was added a CH₂Cl₂ solution of **5d** (80 μ L, 0.5 M, 1 M=1 mol dm⁻³). After being stirred for 4 h, the reaction mixture was quenched by the addition of a CH₂Cl₂ solution of *i*-Pr₂NEt (1 M, 0.5 mL). A THF solution of Bu₄NF (1 mol dm⁻³ in THF, 0.5 mL) was added to the mixture and the whole was stirred at rt for 0.5 h. The mixture was diluted with Et₂O, poured into water, and extracted twice with Et₂O. Organic extracts were dried and concentrated in vacuo. Purification of the residue by flash chromatography (10% ethyl acetate/hexane) gave 101 mg (81% yield) of ring cleavage product **9a** (*anti*: *syn* 98:2); ¹H NMR $\delta=1.15$ (3H, d, $J=6.3$ Hz), 1.44 (9H, s), 1.57—1.72 (2H, m), 2.46 (1H, br s), 2.62 (1H, dd, $J=4.2$ and 14.4 Hz), 2.90 (1H, dd, $J=9.6$ and 14.4 Hz), 3.49—3.61 (2H, m), 3.68—3.73 (1H, m), 4.88 (1H, dd, $J=4.2$ and 9.6 Hz), 7.27—7.38 (5H, m) [a minor diastereomer resonated at $\delta=0.92$ (3H, d, $J=6.0$ Hz)]; ¹³C NMR $\delta=18.4$, 29.6, 39.1, 48.2, 52.8, 60.3, 71.4, 75.4, 126.8, 128.1, 128.6, 140.6, 197.3; IR (liquid film) 3400 (br), 1680, 760, 735, 695 cm⁻¹. Found: C, 65.64; H, 8.74; S, 10.54%. Calcd for C₁₇H₂₆O₃S: C, 65.77; H, 8.44; S, 10.33%.

When the reaction of **6a** and **7a** in the presence of **5d** (0.2 mol. amt.) was performed by a procedure like that described above, except that the initial reaction mixture was not treated with Bu₄NF, *S-t*-butyl (3S)-3-[(1R)-1-methyl-3-(trimethylsilyloxy)propoxy]-3-phenylpropanethioate (**8a**) was obtained in 86% yield (*anti*: *syn* 98:2). **8a:** ¹H NMR $\delta=0.02$ (9H, s), 1.13 (3H, d, $J=6.0$ Hz), 1.44 (9H, s), 1.48—1.59 (1H, m), 1.65—1.77 (1H, m), 2.60 (1H, dd, $J=4.5$ and 14.4 Hz), 2.90 (1H, dd, $J=9.3$ and 14.4 Hz), 3.38—3.50 (2H, m), 3.53—3.61 (1H, m), 4.84 (1H, dd, $J=4.5$ and 9.0 Hz), 7.25—7.33 (5H, m) [a minor diastereomer resonated at 0.98 (3H, d, $J=6.3$ Hz)]; ¹³C NMR $\delta=-0.6$, 19.0, 29.7, 40.4, 47.9, 53.2, 59.4, 70.0, 75.6, 126.7, 127.7, 128.3, 141.6, 197.5; IR (liquid film) 1680, 1090, 840, 750, 695 cm⁻¹.

Ring-cleavage products **9b—g** were prepared by a procedure similar to that described above. For the reaction conditions, see Table 2. The spectral data are as follows:

S-Ethyl (3S)-3-[(1R)-3-Hydroxy-1-methylpropoxy]-3-phenylpropanethioate (9b): ¹H NMR $\delta=1.16$ (3H, d, $J=6.0$ Hz), 1.23 (3H, t, $J=7.5$ Hz), 1.61—1.67 (2H, m), 2.39 (1H, br t, $J=ca.$ 5.5 Hz), 2.74 (1H, dd, $J=4.5$ and 14.7 Hz), 2.88 (2H, q, $J=7.5$ Hz), 3.02 (1H, dd, $J=9.3$ and 14.7 Hz), 3.49—3.64 (2H, m), 3.65—3.74 (1H, m), 4.93 (1H, dd, $J=4.5$ and 9.3 Hz), 7.28—7.39 (5H, m) [a minor diastereomer resonated at 0.93 (3H, d, $J=6.2$ Hz)]; ¹³C NMR $\delta=14.7$, 18.3, 23.5, 39.0, 52.3, 60.4, 71.4, 75.1, 126.8, 128.3, 128.7, 140.4, 196.7; IR (liquid film) 3400 (br), 1685, 760, 740, 700 cm⁻¹. Found: C, 63.52; H, 8.05; S, 11.63%. Calcd for C₁₅H₂₂O₃S: C, 63.80; H, 7.85; S, 11.35%.

Ethyl (3R)-3-[(1R)-3-Hydroxy-1-methylpropoxy]-2,2-dimethyl-3-phenylpropanoate (9c): ¹H NMR $\delta=1.02$ (3H, s), 1.10 (3H, d, $J=6.0$ Hz), 1.11 (3H, s), 1.26 (3H, t, $J=7.2$ Hz), 1.58—1.76 (2H, m), 2.68 (1H, br s), 3.52 (1H, dqd, $J=4.8$, 6.0, and 12.0 Hz), 3.65—3.77 (2H, m), 4.09 (1H, qd, $J=7.2$ and 10.8 Hz), 4.17 (1H, qd, $J=7.2$ and 10.8 Hz), 4.72 (1H, s), 7.28—7.34 (5H, m) [a minor diastereomer resonated at 4.66 (s, 1H)]; ¹³C NMR $\delta=14.1$, 17.9, 19.2, 22.8, 39.2, 47.6, 60.4, 60.6, 71.4, 82.7, 127.8, 128.0, 128.7, 137.4, 176.4; IR (liquid film) 3425 (br), 1730, 770, 750, 705 cm⁻¹. Found: C, 69.15; H, 9.09%. Calcd for C₁₇H₂₆O₄: C, 69.36; H, 8.90%.

(3S)-3-[(1R)-3-Hydroxy-1-methylpropoxy]-1,3-diphenyl-1-propanone (9d): ¹H NMR $\delta=1.16$ (3H, d, $J=6.0$ Hz), 1.61

(2H, q, $J=5.5$ Hz), 2.26 (1H, br t, $J=5.1$ Hz), 3.10 (1H, dd, $J=4.5$ and 16.2 Hz), 3.50—3.70 (4H including dd ($J=8.1$ and 16.2 Hz, 1H) at 3.59, m), 5.14 (1H, dd, $J=4.5$ and 8.1 Hz), 7.28—7.56 (8H, m), 7.91—7.96 (2H, m) [a minor diastereomer resonated at 0.94 (3H, d, $J=6.1$ Hz)]; ^{13}C NMR $\delta=18.5, 39.2, 47.1, 60.5, 71.3, 74.8, 126.9, 128.1, 128.2, 128.5, 128.7, 133.1, 137.3, 141.5, 197.5$; IR (liquid film) 3420 (br), 1680, 745, 695 cm^{-1} ; HRMS (CI) Calcd for $\text{C}_{19}\text{H}_{22}\text{O}_3$: ($\text{M}^+ + 1$), 299.1647. Found: m/z 299.1714.

***S*-*t*-Butyl (3*R*)-3-[(1*R*)-3-Hydroxy-1-methylpropoxy]-5-phenylpentanethioate (9e):** ^1H NMR (C_6D_6) $\delta=0.95$ (3H, d, $J=6.0$ Hz), 1.35 (9H, s), 1.35—1.41 (1H, m), 1.45—1.56 (1H, m), 1.63—1.71 (2H, m), 1.85 (1H, br s), 2.31 (1H, dd, $J=5.1$ and 14.7 Hz), 2.41—2.47 (2H, m), 2.52 (1H, dd, $J=7.5$ and 14.7 Hz), 3.39—3.50 (2H, m), 3.55—3.63 (1H, m), 3.78 (1H, qd, $J=5.3$ and 7.6 Hz), 6.98—7.14 (5H, m) [a minor isomer resonated at 0.87 (3H, d, $J=6.2$ Hz)]; ^{13}C NMR (C_6D_6) $\delta=20.5, 30.2, 31.9, 36.8, 40.2, 48.4, 50.7, 60.6, 73.1, 73.9, 126.7, 129.2, 129.2, 142.7, 198.1$; IR (liquid film) 3440 (br), 1675, 740, 695 cm^{-1} . Found: C, 67.25; H, 8.88; S, 9.72%. Calcd for $\text{C}_{19}\text{H}_{30}\text{O}_3\text{S}$: C, 67.42; H, 8.93; S, 9.47%.

***S*-*t*-Butyl (3*S*)-3-Cyclohexyl-3-[(1*R*)-3-hydroxy-1-methylpropoxy]propanethioate (9f):** ^1H NMR $\delta=1.10$ (3H, d, $J=6.3$ Hz), 0.86—1.77 (13H including s (9H) at 1.43, m), 2.46 (1H, dd, $J=4.5$ and 14.4 Hz), 2.54 (1H, dd, $J=7.8$ and 14.4 Hz), 2.59—2.63 (1H, m), 3.65—3.80 (4H, m) [a minor isomer resonated at 1.44 (9H, s)]; ^{13}C NMR $\delta=19.4, 26.3, 26.4, 26.6, 27.8, 28.8, 29.6, 39.0, 41.3, 47.0, 48.0, 60.5, 73.4, 77.8, 199.0$; IR (liquid film) 3420 (br), 2920, 1685 cm^{-1} . Found: C, 64.48; H, 10.01; S, 10.13%. Calcd for $\text{C}_{17}\text{H}_{32}\text{O}_3\text{S}$: C, 64.51; H, 10.19; S, 10.13%.

***S*-*t*-Butyl (3*S*,4*E*)-3-[(1*R*)-3-Hydroxy-1-methylpropoxy]-5-phenyl-4-pentenethioate (9g):** ^1H NMR $\delta=1.15$ (3H, d, $J=6.0$ Hz), 1.44 (9H, s), 1.70—1.73 (2H, m), 2.61 (1H, dd, $J=4.8$ and 14.1 Hz), 2.71 (1H, br s), 2.78 (1H, dd, $J=8.1$ and 14.1 Hz), 3.65—3.85 (3H, m), 4.45 (1H, dt, $J=4.8$ and 8.3 Hz), 6.05 (1H, dd, $J=8.4$ and 15.9 Hz), 6.58 (1H, d, $J=15.9$ Hz), 7.22—7.39 (5H, m); ^{13}C NMR $\delta=18.8, 29.6, 39.0, 48.2, 50.7, 60.7, 71.9, 74.6, 126.5, 128.0, 128.4, 128.6, 133.1, 136.1, 197.1$; IR (liquid film) 3440, 1680, 750, 695 cm^{-1} . Found: C, 67.82; H, 8.67; S, 9.36%. Calcd for $\text{C}_{19}\text{H}_{28}\text{O}_3\text{S}$: C, 67.82; H, 8.39; S, 9.53%.

Ring Cleavage Reaction with Allylic Silanes 12a,b. The reaction was carried out by a procedure similar to that described for enol silyl ethers 7. Spectral data of the ring cleavage products 13a—i are as follows.

(3*R*)-3-[(1*S*)-3-Phenyl-3-butenyloxy]-1-butanol (13a): ^1H NMR $\delta=1.16$ (3H, d, $J=6.3$ Hz), 1.66 (2H, q, $J=5.5$ Hz), 2.35—2.45 (1H, m), 2.49—2.59 (1H, m), 2.70 (1H, br s), 3.51—3.62 (2H, m), 3.67—3.74 (1H, m), 4.43 (1H, dd, $J=6.0$ and 7.8 Hz), 4.99—5.08 (2H, m), 5.74 (1H, tdd, $J=7.0, 10.2$ and 17.0 Hz), 7.26—7.39 (5H, m); [a minor isomer resonated at 0.99 (3H, d, $J=6.2$ Hz)]; ^{13}C NMR $\delta=18.8, 39.5, 42.8, 60.5, 71.2, 78.8, 116.9, 127.0, 127.8, 128.5, 134.7, 142.0$; IR (liquid film) 3400 (br), 920, 760, 705 cm^{-1} . Found: C, 76.12; H, 9.45%. Calcd for $\text{C}_{14}\text{H}_{20}\text{O}_2$: C, 76.33; H, 9.15%.

(3*R*)-3-[(1*S*)-3-Methyl-1-phenyl-3'-butenyloxy]-1-butanol (13b): ^1H NMR $\delta=1.16$ (3H, d, $J=6.1$ Hz), 1.62—1.69 (2H, m), 1.72 (13H, br s), 2.32 (1H, dd, $J=5.3$ and 13.5 Hz), 2.52 (1H, dd, $J=8.5$ and 13.5 Hz), 2.8 (1H, br), 3.51—3.66 (2H, m), 3.72 (1H, m), 4.55 (1H, dd, $J=5.3$ and 8.5 Hz), 4.70 (1H, m), 4.76 (1H, m) 7.26—7.39 (5H, m) [a minor diastereomer resonated at $\delta=0.96$ (3H, d, $J=6.2$ Hz)]; ^{13}C NMR $\delta=18.2, 22.5, 38.13, 38.8, 46.8, 60.4, 71.0, 77.0, 112.9, 126.8, 127.7, 128.4, 141.8, 142.0$; IR (liquid film) 3400 (br), 1650, 1080, 1050, 890, 755, 700 cm^{-1} ; MS m/z (rel intensity) 179 ($\text{M}^+ - \text{CH}_2 = \text{C}(\text{Me})\text{CH}_2$; 19), 145 (3), 107 (100), HRMS Calcd

for $\text{C}_{11}\text{H}_{15}\text{O}_2$: ($\text{M}^+ - \text{CH}_2 = \text{C}(\text{Me})\text{CH}_2$), 179.1072. Found: m/z 179.1067. Found: C, 77.04; H, 9.60%. Calcd for $\text{C}_{15}\text{H}_{22}\text{O}_2$: C, 76.88; H, 9.46%.

(3*R*)-3-[(1*S*)-3-Methyl-1-(2-phenylethyl)-3-butenyloxy]-1-butanol (13c): ^1H NMR $\delta=1.14$ (3H, d, $J=6.2$ Hz), 1.70—1.90 (5H, m, including br s (3H) at $\delta=1.74$), 2.18—2.30 (2H, m), 2.60—2.90 (3H, m), 3.58 (1H, quint, $J=\text{ca. } 6$ Hz), 4.74 (1H, m), 4.84 (1H, m) [a minor diastereomer resonated at $\delta=1.17$ (3H, d, $J=6.2$ Hz)]; ^{13}C NMR $\delta=19.8, 22.82, 31.3, 35.5, 39.0, 43.6, 60.7, 73.6, 74.6, 112.9, 125.7, 128.27, 128.33, 142.1, 142.6$; IR (liquid film) 3400 (br), 1650, 1050, 890, 745, 700 cm^{-1} ; MS (CI) m/z (rel intensity) 263 (MH^+ ; <1), 207 (55), 117 (98), 91 (100). HRMS Calcd for $\text{C}_{13}\text{H}_{19}\text{O}_2$: ($\text{M}^+ - \text{CH}_2 = \text{C}(\text{Me})\text{CH}_2$), 207.1385. Found: m/z 207.1376. Found: C, 77.60; H, 10.11%. Calcd for $\text{C}_{17}\text{H}_{26}\text{O}_2$: C, 77.82; H, 9.99%.

(3*R*)-3-[(1*S*)-1-Cyclohexyl-3-methyl-3-butenyloxy]-1-butanol (13d): ^1H NMR $\delta=0.95$ —1.45 (8H, m including d ($J=6.1$ Hz, 3H) at $\delta=1.197$), 1.45—1.85 (11H, m), 2.09 (1H, dd, $J=7.5$ and 4.0 Hz), 2.14 (1H, dd, $J=5.3$ and 14.0 Hz), 2.97 (1H, br s), 3.33 (1H, m), 3.65—3.85 (3H, m), 4.70 (1H, m), 4.76 (1H, m) [a minor diastereomer resonated at $\delta=1.122$ (3H, d, $J=6.2$ Hz)]; ^{13}C NMR (C_6D_6) $\delta=20.1, 23.1, 27.0, 27.1, 28.5, 28.8, 39.8, 40.4, 41.3, 60.3, 72.7, 79.7, 112.9, 143.5$; IR (liquid film) 3300 (br), 1685, 1090, 925 cm^{-1} ; MS m/z (rel intensity) 185 ($\text{M}^+ - \text{CH}_2 = \text{C}(\text{Me})\text{CH}_2$; 17), 113 (49), 107 (39), 95 (100). HRMS Calcd for $\text{C}_{11}\text{H}_{21}\text{O}_2$: ($\text{M}^+ - \text{CH}_2 = \text{C}(\text{Me})\text{CH}_2$), 185.1541. Found: m/z 185.1538. Found: C, 74.65; H, 11.87%. Calcd for $\text{C}_{15}\text{H}_{28}\text{O}_2$: C, 74.95; H, 11.74%.

(3*R*)-3-[(1*S*)-3-Methyl-1-[(*E*)-2-phenylethenyl]-3-butenyloxy]-1-butanol (13e): ^1H NMR $\delta=1.17$ (3H, d, $J=6.1$ Hz), 1.65—1.80 (5H, m, including br s (3H) at $\delta=1.77$), 2.27 (1H, dd, $J=5.8$ and 13.7 Hz), 2.39 (1H, dd, $J=7.7$ and 13.5 Hz), 2.98 (1H, br s), 3.65—3.85 (2H, m), 3.87 (1H, sextet, $J=6.1$ Hz), 4.14 (1H, ddd, $J=5.8, 7.7$, and 8.4 Hz), 4.76 (1H, m), 4.80 (1H, m), 6.05 (1H, dd, $J=8.4$ and 15.9 Hz), 6.53 (1H, d, $J=15.9$ Hz), 7.20—7.45 (5H, m); ^{13}C NMR $\delta=18.7, 22.9, 38.8, 44.8, 61.1, 71.6, 76.1, 113.1, 126.5, 127.8, 128.6, 130.1, 132.5, 136.4, 141.8$; IR (liquid film) 3420 (br), 1650, 1050, 965, 890, 750, 695 cm^{-1} ; MS m/z (rel intensity) 205 ($\text{M}^+ - \text{CH}_2 = \text{C}(\text{Me})\text{CH}_2$; 23), 170 (8), 155 (11), 133 (100); HRMS Calcd for $\text{C}_{13}\text{H}_{17}\text{O}_2$: ($\text{M}^+ - \text{CH}_2 = \text{C}(\text{Me})\text{CH}_2$), 205.1229. Found: m/z 205.1234. Found: C, 78.42; H, 9.30%. Calcd for $\text{C}_{17}\text{H}_{24}\text{O}_2$: C, 78.42; H, 9.29%.

(3*R*)-3-[(1*R*)-3-Methyl-1-[(*E*)-2-phenylethenyl]-3-butenyloxy]-1-butanol (a Minor Isomer): ^1H NMR $\delta=1.18$ (3H, d, $J=6.2$ Hz), 1.65—1.85 (5H, m, including br s (3H) at $\delta=1.79$), 2.24 (1H, dd, $J=7.4$ and 13.8 Hz), 2.83 (1H, dd, $J=7.7$ and 13.8 Hz), 2.50 (1H, br s), 3.65—3.90 (3H, m), 4.13 (1H, q, $J=\text{ca. } 6$ Hz), 4.80 (1H, m), 4.84 (1H, m), 6.13 (1H, dd, $J=7.4$ and 16.0 Hz), 6.52 (1H, d, $J=16.0$ Hz), 7.15—7.40 (5H, m); ^{13}C NMR $\delta=21.1, 22.8, 38.9, 44.9, 60.4, 73.9, 78.0, 113.6, 126.4, 127.6, 128.6, 130.8, 131.5, 136.7, 141.8$.

(3*R*)-3-[(1*S*)-3-Methyl-1-(4-methoxyphenyl)-3-butenyloxy]-1-butanol (13f): ^1H NMR $\delta=1.14$ (3H, d, $J=6.1$ Hz), 1.60—1.70 (2H, m), 1.70 (3H, br s), 2.31 (1H, dd, $J=5.6$ and 13.9 Hz), 2.51 (1H, dd, $J=8.3$ and 13.9 Hz), 2.85 (1H, br t, $J=\text{ca. } 5.5$ Hz), 3.45—3.65 (2H, m), 3.72 (1H, m), 3.80 (3H, s), 4.51 (1H, dd, $J=5.6$ and 8.3 Hz), 4.69 (1H, m), 4.75 (1H, m), 6.89 (2H, m), 7.24 (2H, m) [a minor diastereomer resonated at $\delta=0.94$ (3H, d, $J=6.3$ Hz) and 1.73 (3H, br s)]; ^{13}C NMR $\delta=18.3, 22.6, 38.8, 46.8, 55.2, 60.8, 71.0, 76.5, 112.9, 113.8, 128.1, 133.8, 142.2, 159.2$; IR (liquid film) 3420 (br), 3075, 1610, 1245, 890, 835 cm^{-1} ; MS m/z (rel intensity) 209 ($\text{M}^+ - \text{CH}_2 = \text{C}(\text{Me})\text{CH}_2$; 21), 175 (3), 137 (100); HRMS Calcd for $\text{C}_{12}\text{H}_{17}\text{O}_3$: ($\text{M}^+ - \text{CH}_2 = \text{C}(\text{Me})\text{CH}_2$), 209.1178. Found: m/z

209.1174. Found: C, 72.89; H, 9.25%. Calcd for $C_{16}H_{24}O_3$: C, 72.69; H, 9.15%.

(3R)-3-[(1S)-3-Methyl-1-(2-methoxyphenyl)-3-butenyloxy]-1-butanol (13g): 1H NMR δ =1.16 (3H, d, J =6.1 Hz), 1.60—1.75 (5H, m including br s (3H) at J =1.76), 2.37 (1H, dd, J =5.3 and 13.8 Hz), 2.43 (1H, dd, J =8.2 and 13.8 Hz), 3.07 (1H, br s), 3.50—3.80 (2H, m), 3.83 (3H, s), 4.68 (1H, m), 4.74 (1H, m), 5.08 (1H, dd, J =5.3 and 8.2 Hz), 6.87 (1H, dd, J =1.0 and 8.2 Hz), 6.98 (1H, dt, J =1.0 and 7.5 Hz), 7.25 (1H, dt, J =1.7 and 7.8 Hz), 7.37 (1H, dd, J =1.7 and 7.5 Hz) [a minor diastereomer resonated at δ =0.98 (3H, d, J =6.3 Hz) and 4.99 (1H, dd, J =4.9 and 8.1 Hz)]; ^{13}C NMR δ =18.5, 22.5, 38.9, 45.5, 55.2, 60.7, 70.5, 71.5, 110.4, 112.4, 120.7, 127.3, 128.4, 130.1, 142.7, 157.1; IR (liquid film) 3420 (br), 1600, 1240, 890, 755 cm^{-1} ; MS m/z (rel intensity) 209 (M^+ — CH_2 =C(Me)CH₂; 23), 175 (4), 137 (100); HRMS Calcd for $C_{12}H_{17}O_3$: (M^+ — CH_2 =C(Me)CH₂), 209.1178. Found: m/z 209.1184. Found: C, 72.99; H, 9.43%. Calcd for $C_{16}H_{24}O_3$: C, 72.69; H, 9.15%.

(3R)-3-[(1S)-1-(2-Furyl)-3-methyl-3-butenyloxy]-1-butanol (13h): 1H NMR δ =1.16 (3H, d, J =6.1 Hz), 1.55—1.75 (5H, m including br s (3H) at J =1.69), 2.49 (1H, dd, J =6.6 and 13.9 Hz), 2.59 (1H, dd, J =7.7 and 13.9 Hz), 2.60 (1H, br), 3.45—3.85 (3H, m), 4.56 (1H, dd, J =6.6 and 7.7 Hz), 4.70 (1H, m), 4.76 (1H, m), 6.26 (1H, dd, J =0.6 and 3.3 Hz), 6.33 (1H, dd, J =1.8 and 3.3 Hz), 7.40 (1H, dd, J =0.6 and 1.8 Hz) [a minor diastereomer resonated at δ =0.97 (3H, d, J =6.2 Hz)]; ^{13}C NMR δ =18.9, 22.5, 38.9, 43.0, 60.8, 70.4, 71.9, 108.0, 110.0, 113.0, 141.5, 142.2, 154.2; IR (liquid film) 3410 (br), 1650, 1050, 890, 740 cm^{-1} ; MS m/z (rel intensity) 169 (M^+ — CH_2 =C(Me)CH₂; 20), 135 (4), 97 (100); HRMS Calcd for $C_9H_{13}O_3$: (M^+ — CH_2 =C(Me)CH₂), 169.0865. Found: m/z 169.0877. Found: C, 69.41; H, 9.08%. Calcd for $C_{13}H_{20}O_3$: C, 69.61; H, 8.99%.

(3R)-3-[(1S)-1-(E)-2-Phenylethenyl]-3-butenyloxy]-1-butanol (13i) (Ring Cleavage Reaction with Allyltributyltin). The reaction was carried out by a procedure similar to that described for the enol silyl ethers 7.

13i: 1H NMR δ =1.17 (3H, d, J =6.0 Hz), 1.65—1.80 (2H, m), 2.25—1.45 (2H, m), 2.92 (1H, br s), 3.65—3.92 (3H, m including sextet (1H, J =ca. 6 Hz) at δ =3.85), 4.04 (1H, td, J =6.3 and 8.4 Hz), 5.05—5.17 (2H, m), 5.83 (1H, tdd, J =7.0, 10.2, and 17.1 Hz), 6.08 (1H, dd, J =8.4 and 15.9 Hz), 6.54 (1H, d, J =15.9 Hz), 7.20—7.45 (5H, m) [a minor diastereomer resonated at δ =1.18 (3H, d, J =6.2 Hz)]; ^{13}C NMR δ =19.0, 38.9, 40.8, 61.1, 71.9, 77.3, 117.3, 126.5, 127.9, 128.6, 129.8, 132.7, 134.2, 136.3; IR (liquid film) 3420 (br), 1055, 970, 915, 750, 690 cm^{-1} ; MS m/z (rel intensity) 205 (M^+ — CH_2 =CHCH₂; 22), 133 (100); HRMS Calcd for $C_{13}H_{17}O_2$: (M^+ — CH_2 =CHCH₂), 205.1229. Found: m/z 205.1232. Found: C, 77.75 H, 9.20%. Calcd for $C_{16}H_{22}O_2$: C, 78.01; H, 9.00%.

S-*t*-Butyl (3S)-3-[(1S)-1-Methyl-2-formylethoxy]-3-phenylpropanethioate (14a) (Representative Procedure for PCC Oxidation). To a solution of **9a** (93 mg, 0.30 mmol) in CH_2Cl_2 (1 mL) was added PCC (215 mg, 1.0 mmol). After being stirred for 5 h, the mixture was diluted with Et_2O and filtered through Cellite. The filtrate was concentrated in vacuo. Purification of the residue by flash chromatography (3% ethyl acetate/hexane) gave 77 mg (84% yield) of **14a**: 1H NMR δ =1.22 (3H, d, J =6.3 Hz), 1.44 (9H, s), 2.35 (1H, ddd, J =1.8, 5.1, and 15.9 Hz), 2.48 (1H, ddd, J =3.0, 6.9, and 15.9 Hz), 2.59 (1H, dd, J =4.2 and 14.7 Hz), 2.90 (1H, dd, J =9.6 and 14.7 Hz), 3.79—3.89 (1H, m), 4.88 (1H, dd, J =4.2 and 9.6 Hz), 7.25—7.36 (5H, m), 9.55 (1H, dd, J =1.8 and 3.0 Hz) [a minor diastereomer resonated at δ =9.79 (1H, dd, J =1.8 and 5.7 Hz)]; ^{13}C NMR δ =19.0, 29.6, 48.1, 50.7, 52.8, 68.1, 75.7, 126.6, 128.1, 128.6, 140.7, 197.3, 201.7; IR (liquid film) 2720, 1720, 1680, 760,

740, 700 cm^{-1} .

Aldehydes **14b—g** and **16a—e** were prepared by a procedure similar to that described above. The spectral data are as follows:

S-Ethyl (3S)-3-[(1S)-1-Methyl-2-formylethoxy]-3-phenylpropanethioate (14b): 1H NMR δ =1.22 (3H, t, J =6.0 Hz), 1.23 (3H, t, J =7.5 Hz), 2.34 (1H, ddd, J =1.8, 5.1, and 15.9 Hz), 2.49 (1H, ddd, J =3.0, 7.2, and 15.9 Hz), 2.71 (1H, dd, J =4.2 and 15.0 Hz), 2.87 (2H, q, J =7.5 Hz), 3.02 (1H, dd, J =9.3 and 15.0 Hz), 3.89—3.79 (1H, m), 4.93 (1H, dd, J =4.2 and 9.3 Hz), 7.27—7.38 (5H, m), 9.55 (1H, dd, J =1.8 and 3.0 Hz) [a minor diastereomer resonated at δ =9.76 (1H, dd, J =1.2 and 3.3 Hz)]; ^{13}C NMR δ =14.7, 18.9, 23.4, 50.7, 52.4, 68.1, 75.5, 126.7, 128.3, 128.6, 140.6, 196.7, 201.6; IR (liquid film) 2720, 1720, 1680, 765, 740, 700 cm^{-1} .

Ethyl (3R)-2,2-Dimethyl-3-[(1R)-1-methyl-2-formylethoxy]-3-phenylpropanoate (14c): 1H NMR δ =0.96 (3H, s), 1.11 (3H, s), 1.16 (3H, d, J =6.3 Hz), 1.25 (3H, t, J =7.2 Hz), 2.37 (1H, ddd, J =2.1, 5.1, and 15.9 Hz), 2.49 (1H, ddd, J =3.0, 6.9, and 15.9 Hz), 3.86—3.76 (1H, m), 4.08 (1H, qd, J =7.2 and 10.8 Hz), 4.16 (1H, qd, J =7.2 and 10.8 Hz), 4.76 (1H, s), 7.23—7.35 (5H, m), 9.66 (1H, dd, J =2.1 and 3.0 Hz); ^{13}C NMR δ =14.2, 18.7 (2C), 22.5, 47.8, 50.9, 60.5, 68.1, 82.8, 127.8, 128.0, 128.7, 137.5, 176.4, 201.5; IR (liquid film) 2725, 1720, 770, 745, 710 cm^{-1} .

(3S)-3-[(1R)-1-Methyl-2-formylethoxy]-1,3-diphenyl-1-propanone (14d): 1H NMR δ =1.21 (3H, d, J =6.3 Hz), 2.31 (1H, ddd, J =1.8, 4.8, and 15.9 Hz), 2.43 (1H, ddd, J =3.0, 7.5, and 15.9 Hz), 3.00 (1H, dd, J =4.2 and 16.2 Hz), 3.60 (1H, dd, J =8.7 and 16.2 Hz), 3.88 (1H, dqd, J =4.8, 6.3, and 7.5 Hz), 5.14 (1H, dd, J =4.2 and 8.7 Hz), 7.30—7.57 (8H, m), 7.91—7.95 (2H, m), 9.51 (1H, dd, J =1.8 and 3.0 Hz) [a minor diastereomer resonated at 9.72 (1H, dd, J =1.8 and 2.7 Hz)]; ^{13}C NMR δ =18.9, 47.1, 50.8, 67.8, 75.0, 126.9, 128.1, 128.2, 128.5, 128.6, 133.1, 137.2, 141.5, 197.7, 201.8; IR (liquid film) 2725, 1720, 1690, 750, 700, 695 cm^{-1} .

S-*t*-Butyl (3R)-3-[(1R)-1-Methyl-2-formylethoxy]-5-phenylpentanethioate (14e): 1H NMR δ =1.19 (3H, d, J =6.3 Hz), 1.46 (9H, s), 1.78—1.86 (2H, m), 2.45 (1H, ddd, J =1.8, 5.4, and 16.5 Hz), 2.54—2.66 (4H, m), 2.70 (2H, dd, J =7.2 and 14.4 Hz), 3.87—3.95 (1H, m), 3.97—4.07 (1H, m), 7.15—7.30 (5H, m), 9.79 (1H, dd, J =1.8 and 2.1 Hz) [a minor diastereomer resonated at 9.76 (1H, t, J =2.1 Hz)]; ^{13}C NMR δ =20.5, 29.7, 31.2, 36.1, 48.2, 49.9, 50.6, 69.2, 73.7, 125.9, 128.2, 128.4, 141.6, 198.1, 201.2; IR (liquid film) 2710, 1720, 1680, 735, 700 cm^{-1} .

S-*t*-Butyl (3S)-3-Cyclohexyl-3-[(1R)-1-methyl-2-formylethoxy]propanethioate (14f): 1H NMR δ =1.16 (3H, d, J =6.0 Hz), 1.44 (9H, s), 0.89—1.75 (11H, m), 2.43 (1H, ddd, J =2.1, 5.4, and 16.2 Hz), 2.50—2.63 (3H, m), 3.66—3.71 (1H, m), 3.93—4.02 (1H, m), 9.78 (1H, dd, J =2.1 and 2.7 Hz) [a minor diastereomer resonated at δ =9.75 (1H, dd, J =1.8 and 2.7 Hz)]; ^{13}C NMR δ =20.4, 26.2, 26.3, 26.5, 27.9, 28.8, 29.7, 41.6, 47.0, 48.0, 50.7, 69.4, 78.2, 199.1, 201.7; IR (liquid film) 2770, 1720, 1680 cm^{-1} .

S-*t*-Butyl (3S,4E)-3-[(1R)-1-Methyl-2-formylethoxy]-5-phenyl-4-pentenethioate (14g): 1H NMR δ =1.21 (3H, d, J =6.0 Hz), 1.44 (9H, s), 2.43 (1H, ddd, J =1.8, 5.1, and 16.2 Hz), 2.58 (1H, ddd, J =2.7, 7.5, and 16.2 Hz), 2.59 (1H, dd, J =5.1 and 14.4 Hz), 2.78 (1H, dd, J =8.7 and 14.4 Hz), 4.03—4.13 (1H, m), 4.44—5.51 (1H, m), 6.01 (1H, dd, J =8.1 and 15.9 Hz), 6.57 (1H, d, J =15.9 Hz), 7.22—7.39 (5H, m), 9.73 (1H, dd, J =1.8 and 2.7 Hz) [a minor diastereomer resonated at 9.78 (1H, dd, J =2.1 and 2.4 Hz)]; ^{13}C NMR δ =19.3, 29.7, 48.2, 50.7, 50.8, 67.9, 74.9, 126.6, 128.0, 128.6 (2C), 132.9, 136.1, 197.1, 201.4; IR (liquid film) 2725, 1720, 1680, 750, 695 cm^{-1} .

(3R)-3-[(1S)-1-Phenyl-3-butenyloxy]butanal (16a): 1H NMR δ =1.23 (3H, d, J =6.0 Hz), 2.31—2.40 (1H, m), 2.47—2.58 (1H,

m), 3.80—3.90 (1H, m), 4.38 (1H, dd, $J=5.7$ and 7.8 Hz), 4.97—5.06 (2H, m), 5.67—5.81 (1H, m), 7.25—7.37 (5H, m), 9.57 (1H, dd, $J=1.8$ and 3.0 Hz); ^{13}C NMR $\delta=19.2, 42.7, 50.9, 67.7, 79.2, 116.8, 126.9, 127.8, 128.4, 134.8, 141.9, 201.7$; IR (film) 2720, 1720, 760, 700 cm^{-1} .

(3R)-3-[(1S)-3-Methyl-1-phenyl-3-butenyloxy]butanal (16b): ^1H NMR $\delta=1.22$ (3H, d, $J=6.3$ Hz), 1.71 (3H, br s), 2.22—2.40 (2H, m), 2.24—2.55 (2H, m), 3.83 (1H, m), 4.51 (1H, dd, $J=5.7$ and 8.1 Hz), 4.67 (1H, m), 4.74 (1H, m), 7.24—7.37 (5H, m), 9.57 (1H, dd, $J=\text{ca. } 2$ and 3 Hz) [a minor diastereomer resonated at $\delta=1.06$ (3H, d, $J=6.3$ Hz) and 9.82 (1H, t, $J=2$ Hz)]; IR (film) 2720, 1725, 890, 755, 700 cm^{-1} .

(3R)-3-[(1S)-3-Methyl-1-[(E)-2-phenylethenyl]-3-butenyloxy]butanal (16c): ^1H NMR $\delta=1.22$ (3H, d, $J=6.3$ Hz), 1.75 (3H, br s), 2.24 (1H, dd, $J=6.0$ and 13.8 Hz), 2.37 (1H, dd, $J=7.2$ and 13.8 Hz), 2.45 (1H, ddd, $J=1.8, 4.8,$ and 16.2 Hz), 2.58 (1H, ddd, $J=\text{ca. } 3, 7.5,$ and 16.2 Hz), 4.06—4.17 (2H, m), 4.75 (1H, m), 4.79 (1H, m), 6.02 (1H, dd, $J=8.3$ and 15.9 Hz), 6.51 (1H, d, $J=15.9$ Hz), 7.22—7.41 (5H, m), 9.74 (dd, $J=\text{ca. } 2$ and 3 Hz); IR (film) 2725, 1725, 890, 750, 695 cm^{-1} .

(3R)-3-[(1S)-3-Methyl-1-(4-methoxyphenyl)-3-butenyloxy]butanal (16d): ^1H NMR $\delta=1.20$ (3H, d, $J=6.0$ Hz), 1.69 (1H, br s), 2.23—2.37 (2H, m), 2.42—2.54 (2H, m), 3.81 (3H, s), 3.82 (1H, m), 4.46 (1H, dd, $J=5.9$ and 8.0 Hz), 4.65 (1H, m), 4.73 (1H, m), 6.76 (2H, m), 7.19 (2H, m), 9.55 (1H, dd, $J=\text{ca. } 2$ and 3 Hz) [a minor isomer resonated at $\delta=1.05$ (3H, d, $J=6.0$ Hz) and 9.81 (1H, t, $J=\text{ca. } 2.5$ Hz)].

(3R)-3-[(1S)-3-Methyl-1-(2-furyl)-3-butenyloxy]butanal (16e): ^1H NMR $\delta=1.22$ (3H, d, $J=6.3$ Hz), 1.69 (3H, br s), 2.34 (1H, ddd, $J=2.0, 6.0,$ and 13.8 Hz), 2.42—2.62 (3H, m), 3.92 (1H, m), 4.55 (1H, t, $J=7.2$ Hz), 4.70 (1H, m), 4.75 (1H, m), 6.25 (1H, dd, $J=0.5$ and 3.3 Hz), 6.32 (1H, dd, $J=\text{ca. } 2.0$ and 3.3 Hz), 7.38 (1H, dd, $J=0.5$ and 2.0 Hz), 9.55 (1H, dd, $J=2.0$ and 3.2 Hz) [a minor isomer resonated at $\delta=1.05$ (3H, t, $J=6.3$ Hz) and 9.82 (3H, t, $J=2.2$ Hz)].

S-*t*-Butyl (S)-3-Hydroxy-3-phenylpropanethioate (15a) (Representative Procedure for the Liberation of Enantiomerically Enriched Secondary Alcohols). To a solution of aldehyde **14a** (71 mg, 0.23 mmol) in benzene (3 mL) at rt was added dibenzylammonium trifluoroacetate (78 mg, 0.25 mmol). After being stirred for 3 h, the mixture was poured into water and extracted twice with ethyl acetate. Organic extracts were dried and concentrated in vacuo. Purification of the residue by flash chromatography (5% ethyl acetate/hexane) gave 44 mg (80% yield) of **15a**: $[\alpha]_{\text{D}}^{25} -43.4$ (c 2.15, benzene) (lit.²⁰) $[\alpha]_{\text{D}}^{25} -41.4$). (S)-MTPA ester of **15a**: ^1H NMR $\delta=1.39$ (9H, s), 2.86 (1H, dd, $J=4.2$ and 16.2 Hz), 3.21 (1H, dd, $J=9.6$ and 16.2 Hz), 3.41—3.43 (3H, m), 6.45 (1H, dd, $J=4.2$ and 9.6 Hz), 7.27—7.42 (10H, m) [a minor diastereomer resonated at $\delta=1.44$ (9H, s)].

The following compounds were obtained by a procedure similar to that described above starting from **14b—g** and **16a—e**:

S-Ethyl (S)-3-Hydroxy-3-phenylpropanethioate (15b): $[\alpha]_{\text{D}}^{25} -60.7$ (c 2.00, benzene) (lit.²⁰) $[\alpha]_{\text{D}}^{25} -56.3$).

Ethyl (R)-2,2-Dimethyl-3-hydroxy-3-phenylpropanoate (15c): ^1H NMR $\delta=1.10$ (3H, s), 1.13 (3H, s), 1.26 (3H, t, $J=7.2$ Hz), 3.19 (1H, br s), 4.17 (2H, q, $J=7.2$ Hz), 4.88 (1H, s), 7.29—7.32 (5H, m); ^{13}C NMR $\delta=14.1, 19.1, 23.0, 47.5, 60.9, 78.7, 127.7$ (3C), 140.0, 177.7; IR (liquid film) 3450 (br), 2980, 1720, 770, 745, 705 cm^{-1} . Treatment of **15c** with aq NaOH in THF afforded (R)-2,2-dimethyl-3-hydroxy-3-phenylpropanoic acid: $[\alpha]_{\text{D}}^{25} -4.7$ (c 0.96, MeOH) (lit.²⁴) $[\alpha]_{\text{D}}^{25} +4.5$ for the S enantiomer).

(S)-3-Hydroxy-1,3-diphenyl-1-propanone (15d): $[\alpha]_{\text{D}}^{25} -27.9$

(c 0.980, MeOH) (lit.²¹) $[\alpha]_{\text{D}}^{25} -31.97$ for the R enantiomer).

S-*t*-Butyl (R)-3-Hydroxy-5-phenylpentanethioate (15e): $[\alpha]_{\text{D}}^{25} -9.24$ (c 2.51, benzene) (lit.²⁰) $[\alpha]_{\text{D}}^{25} -7.5$).

S-*t*-Butyl (S)-3-Cyclohexyl-3-hydroxypropanethioate (15f): ^1H NMR $\delta=0.96$ —1.40 (6H, m), 1.46 (9H, s), 1.63—1.84 (5H, m), 2.55 (1H, dd, $J=9.0$ and 15.6 Hz), 2.65 (1H, dd, $J=3.0$ and 15.6 Hz), 2.72 (1H, d, $J=3.9$ Hz), 3.75—3.82 (1H, m); ^{13}C NMR $\delta=26.0, 26.1, 26.4, 28.1, 28.8, 29.7, 43.1, 48.2, 48.4, 72.8, 201.1$; IR (liquid film) 3450 (br), 2920, 1680 cm^{-1} . Treatment of **15f** with aq NaOH in THF afforded (S)-3-cyclohexyl-3-hydroxypropanoic acid: $[\alpha]_{\text{D}}^{25} -23.5$ (c 1.50, CHCl_3) (lit.²⁵) $[\alpha]_{\text{D}}^{25} -31.8$ for the R enantiomer).

S-*t*-Butyl (S)-3-Hydroxy-5-phenyl-4-pentenoate (15g): ^1H NMR $\delta=1.47$ (9H, s), 2.71—2.82 (2H, m), 2.97 (1H, br s), 4.71—4.77 (1H, m), 6.18 (1H, dd, $J=6.3$ and 15.6 Hz), 6.64 (1H, dd, $J=0.9$ and 15.6 Hz), 7.21—7.38 (5H, m); ^{13}C NMR $\delta=29.7, 48.7, 50.9, 69.5, 126.5, 127.7, 128.5, 129.9, 130.7, 136.5, 199.6$; IR (liquid film) 3450, 2960, 1680, 755, 700 cm^{-1} . Treatment of **15g** with aq NaOH in THF afforded (S)-3-hydroxy-5-phenyl-4-pentenoic acid: $[\alpha]_{\text{D}}^{25} +18$ (c 0.57, MeOH) (lit.²⁶) $[\alpha]_{\text{D}}^{25} -16.2$).

(S)-1-Phenyl-3-buten-1-ol (17a): $[\alpha]_{\text{D}}^{25} -41$ (c 0.26, benzene) (lit.²²) $[\alpha]_{\text{D}}^{25} -44.92$.

(S)-3-Methyl-1-phenyl-3-buten-1-ol (17b): $[\alpha]_{\text{D}}^{21} -38.1$ (c 1.97, benzene) (lit.²³) $[\alpha]_{\text{D}}^{25} -46.6$.

(S)-5-Methyl-1-phenyl-1,5-hexadien-3-ol (17c):³² $[\alpha]_{\text{D}}^{22} -1.48$ (c 2.43, Et_2O) (lit.²³) $[\alpha]_{\text{D}}^{25} -1.69$. The ee of **17c** was determined to be 92% by ^1H NMR analysis of the (S)-MTPA ester. (S)-MTPA ester of **17c**: ^1H NMR $\delta=1.73$ (3H, br s), 2.39 (1H, dd, $J=5.6$ and 14.1 Hz), 2.54 (1H, ddd, $J=0.6, 8.1,$ and 14.1 Hz), 3.54 (3H, q, $J=1.2$ Hz), 4.71 (1H, m), 4.77 (1H, m), 5.85 (1H, m), 6.20 (1H, dd, $J=7.8$ and 15.9 Hz), 6.73 (1H, d, $J=15.9$ Hz), 7.25—7.43 (8H, m), 7.54 (2H, m) [a minor diastereomer resonated at $\delta=1.80$ (3H, br s), 3.57 (3H, q, $J=1.2$ Hz), 4.83 (1H, m), 4.88 (1H, m), 6.06 (1H, dd, $J=7.8$ and 16.0 Hz), 6.62 (1H, d, $J=16.0$ Hz)].

(S)-1-(4-Methoxyphenyl)-3-methyl-3-buten-1-ol (17d):³² $[\alpha]_{\text{D}}^{25} +41.0$ (c 2.01, benzene). The ee of **17d** was determined to be 81% by ^1H NMR analysis of the (S)-MTPA ester. (S)-MTPA ester of **17d**: ^1H NMR $\delta=1.78$ (3H, br s), 2.45 (1H, dd, $J=5.4$ and 14.3 Hz), 2.70 (1H, dd, $J=8.9$ and 14.3 Hz), 3.41 (3H, br s), 3.81 (3H, s), 4.64 (1H, m), 4.71 (1H, m), 6.12 (1H, dd, $J=5.5$ and 8.6 Hz), 6.88 (2H, m), 7.2—7.45 (7H, m) [a minor diastereomer resonated at $\delta=1.76$ (3H, br s), 3.51 (3H, br s), 4.79 (1H, m), 4.85 (1H, m)].

(S)-1-(2-Furyl)-3-methyl-3-buten-1-ol (17e):³² The ee of **17e** was determined to be 77% by ^1H NMR analysis of the (S)-MTPA ester. (S)-MTPA ester of **17e**: ^1H NMR $\delta=1.69$ (3H, br s), 2.64 (1H, dd, $J=5.8$ and 14.3 Hz), 2.75 (1H, dd, $J=8.6$ and 14.3 Hz), 3.44 (3H, q, $J=1.2$ Hz), 4.67 (1H, m), 4.74 (1H, m), 6.25 (1H, dd, $J=5.9$ and 8.6 Hz), 6.35 (1H, dd, $J=2.0$ and 3.1 Hz), 6.43 (1H, br d, $J=3.1$ Hz), 7.28—7.45 (6H, m) [a minor diastereomer resonated at $\delta=1.76$ (3H, br s), 3.51 (3H, q, $J=1.2$ Hz), 4.81 (1H, m), 4.86 (1H, m)].

Reaction of Racemic Acetal rac-6a and Enol Silyl Ether 7a Catalyzed by Phenyl Boron Complex 5c. The reaction of **rac-6a** and **7a** was carried out at r.t. for 3 h using a 0.2 mol. amt. of **5c** according to a procedure similar to that described in the representative procedure for a catalytic ring-cleavage reaction. The purification of crude products by flash chromatography (5—10% ethyl acetate/hexane) gave **9a** (*anti*: *syn*=92:8, 69% yield) and (2R,4S)-**6a** (27% yield), whose ee was determined to be 51% by a capillary GC analysis.

This work was supported partially by a Grant-in-Aid for

Scientific Research No. 30135628 from the Ministry of Education, Science and Culture.

References

- 1) Review: a) A. Alexakis and P. Mangency, *Tetrahedron Asymmetry*, **1**, 477 (1990); b) D. Seebach, R. Imwinkelreid, and T. Weber, in "Modern Synthetic Methods," ed by R. Scheffold, Springer Verlag, Berlin (1986), Vol. 4, p. 125. For leading references, see also: c) K. Ishihara, N. Hanaki, and H. Yamamoto, *J. Am. Chem. Soc.*, **115**, 10695 (1993); d) S. E. Denmark and N. G. Almstead, *J. Am. Chem. Soc.*, **113**, 8089 (1991).
- 2) a) J. M. McNamara and Y. Kishi, *J. Am. Chem. Soc.*, **104**, 7371 (1982); b) H. Sekizaki, M. Jung, J. M. McNamara, and Y. Kishi, *J. Am. Chem. Soc.*, **104**, 7372 (1982); c) P. A. Bartlett, W. S. Johnson, and J. D. Elliott, *J. Am. Chem. Soc.*, **105**, 2088 (1983).
- 3) For recent mechanistic studies, see Ref. 1d and a) S. E. Denmark and T. M. Willson, *J. Am. Chem. Soc.*, **111**, 3475 (1989); b) I. Mori, K. Ishihara, L. A. Flippin, K. Nozaki, H. Yamamoto, P. A. Bartlett, and C. H. Heathcock, *J. Org. Chem.*, **55**, 6107 (1990); c) S. E. Denmark and N. G. Almstead, *J. Org. Chem.*, **56**, 6458 (1991); d) T. Sammakia and R. S. Smith, *J. Org. Chem.*, **57**, 2997 (1992); e) T. Sammakia and R. S. Smith, *J. Am. Chem. Soc.*, **114**, 10998 (1992).
- 4) a) D. Seebach and M. Zuger, *Helv. Chim. Acta*, **65**, 495 (1982); b) S. J. Danishefsky, D. W. Armistead, F. E. Wincott, H. G. Selnick, and R. Hungate, *J. Am. Chem. Soc.*, **109**, 8117 (1987).
- 5) a) V. M. Choi, J. D. Elliott, and W. S. Johnson, *Tetrahedron Lett.*, **25**, 591 (1984); b) I. R. Silverman, C. Edington, J. D. Elliott, and W. S. Johnson, *J. Org. Chem.*, **52**, 180 (1987).
- 6) a) T. Harada, T. Hayashiya, I. Wada, N. Iwa-ake, and A. Oku, *J. Am. Chem. Soc.*, **109**, 527 (1987); b) T. Harada, T. Shintani, and A. Oku, *J. Am. Chem. Soc.*, **117**, 12346 (1995); c) For a review, see: T. Harada and A. Oku, *Synlett*, **1994**, 95.
- 7) A reductive cleavage of chiral cyclic acetals catalyzed by an aluminium catalyst has been reported: K. Ishihara, N. Hanaki, and H. Yamamoto, *Synlett*, **1993**, 127, and Ref. 1c.
- 8) For Lewis acid-catalyzed cleavage acyclic acetals, see: a) T. Tsunoda, M. Suzuki, and R. Noyori, *Tetrahedron Lett.*, **21**, 71 (1980); b) H. Sakurai, K. Sasaki, and A. Hosomi, *Tetrahedron Lett.*, **22**, 745 (1981); c) T. Mukaiyama, H. Nagaoka, M. Murakami, and M. Ohshima, *Chem. Lett.*, **1985**, 977.
- 9) W. S. Johnson, P. H. Crackett, J. D. Elliott, J. J. Jagodzinski, S. D. Lindell, and S. Natarajan, *Tetrahedron Lett.*, **25**, 3951 (1984).
- 10) a) Takasu and H. Yamamoto, *Synlett*, **1990**, 194; b) D. S. Sartor, J. Saffrich, and G. Helmchen, *Synlett*, **1990**, 197.
- 11) a) S. Kiyooka, Y. Kaneko, M. Komura, H. Matsuo, and M. Nakano, *J. Org. Chem.*, **56**, 2276 (1991); b) E. R. Parmee, O. Tempkin, S. Masamune, and A. Abiko, *J. Am. Chem. Soc.*, **113**, 9365 (1991); c) E. R. Parmee, Y. Hong, O. Tempkin, and S. Masamune, *Tetrahedron Lett.*, **33**, 1729 (1992); d) S. Kiyooka, Y. Kaneko, and K. Kume, *Tetrahedron Lett.*, **33**, 4927 (1992); e) E. J. Corey, C. L. Cywin, and T. D. Roper, *Tetrahedron Lett.*, **33**, 6907 (1992).
- 12) For mechanistic studies on the silyl group migration, see: a) E. M. Carreira and R. A. Singer, *Tetrahedron Lett.*, **35**, 4323 (1994); b) S. E. Denmark and C.-T. Chen, *Tetrahedron Lett.*, **35**, 4327 (1994); c) T. K. Hollis and B. Bosnich, *J. Am. Chem. Soc.*, **117**, 4570 (1995).
- 13) Preliminary communication of this work has appeared: M. Kinugasa, T. Harada, K. Fujita, and A. Oku, *Synlett*, **1996**, 43.
- 14) a) E. J. Corey and T.-P. Loh, *J. Am. Chem. Soc.*, **113**, 8966 (1991); b) E. J. Corey and T.-P. Loh, *Tetrahedron Lett.*, **34**, 3979 (1993).
- 15) For the reaction using **5d**, purification of the crude mixture without treatment with Bu₄NF by flash chromatography afforded **8a** (*anti*:*syn*=98:2) in 86% yield.
- 16) A similar mechanism has been proposed for the mechanistically related Mukaiyama aldol reaction.^{11c)}
- 17) R. M. Beesley, C. K. Ingold, and J. F. Thorpe, *J. Chem. Soc.*, **1915**, 1080.
- 18) A silicon-catalyzed process is a possible candidate.¹²⁾
- 19) W. S. Johnson, A. B. Kelson, and J. D. Elliott, *Tetrahedron Lett.*, **29**, 3757 (1988).
- 20) S. Kobayashi, H. Uchiro, Y. Fujishita, I. Shiina, and T. Mukaiyama, *J. Am. Chem. Soc.*, **113**, 4247 (1991).
- 21) S. H. Mashraqui and R. M. Kellogg, *J. Org. Chem.*, **49**, 2513 (1984).
- 22) H. C. Brown and P. K. Jadhav, *J. Am. Chem. Soc.*, **105**, 2092 (1983).
- 23) N. Minowa and T. Mukaiyama, *Bull. Chem. Soc. Jpn.*, **60**, 3697 (1987).
- 24) A. S. Kende, K. Kawamura, and M. J. Orwat, *Tetrahedron Lett.*, **30**, 5821 (1989).
- 25) R. O. Duthaler, P. Herold, W. Lottenbach, K. Oertler, and M. Riediker, *Angew. Chem.*, **101**, 490 (1989).
- 26) H. H. Meyer, *Liebigs Ann. Chem.*, **1979**, 484.
- 27) Y. Kita, J. Segawa, J. Haruta, H. Yasuda, and Y. Tamura, *J. Chem. Soc., Perkin Trans. 1*, **1982**, 1099.
- 28) G. Hagen and H. Mayr, *J. Am. Chem. Soc.*, **113**, 4954 (1991).
- 29) T. Tsunoda, M. Suzuki, and R. Noyori, *Tetrahedron Lett.*, **21**, 1357 (1980).
- 30) E. W. McChesney and W. K. Wann, Jr., *J. Am. Chem. Soc.*, **59**, 1116 (1937).
- 31) B. Helferich and H. Grünert, *Liebigs Ann. Chem.*, **545**, 178 (1940).
- 32) G. E. Keck, D. Krishnamurthy, and M. C. Grier, *J. Org. Chem.*, **58**, 6543 (1993).