

Manganese-Catalyzed Silylmagnesiation of Acetylenes and 1,3-Dienes

Jun Tang, Hiroshi Shinokubo, and Koichiro Oshima*

Department of Material Chemistry, Graduate School of Engineering, Kyoto University, Sakyo-ku, Kyoto 606-01

(Received August 7, 1996)


The treatment of 4-benzyloxy-1-butyne with $\text{PhMe}_2\text{SiMgMe}$ in the presence of a catalytic amount of MnCl_2 gave a monosilylated product, $(E)\text{-PhCH}_2\text{OCH}_2\text{CH}_2\text{CH}=\text{CHSiMe}_2\text{Ph}$, selectively after an aqueous workup. Meanwhile, the reaction of isoprene with $\text{PhMe}_2\text{SiMgMe}$ under manganese catalysis and the subsequent addition of iodomethane afforded 4-dimethylphenylsilyl-2,3-dimethyl-1-butene in good yield.

We have reported¹⁾ on the preparation of tris(trialkylsilyl)manganate $(\text{R}_3\text{Si})_3\text{MnMgMe}$ and its reaction with acetylenes or 1,3-dienes. For instance, the treatment of 4-benzyloxy-1-butyne (**1a**) with $(\text{PhMe}_2\text{Si})_3\text{MnMgMe}$, derived from PhMe_2SiLi , MeMgI , and MnCl_2 , afforded a disilylated product, 4-benzyloxy-1,2-bis(dimethylphenylsilyl)-1-butene (**2**), in 51% yield as a stereoisomeric mixture ($Z/E = 42/58$). Meanwhile, a treatment of isoprene (**3**) with $(\text{PhMe}_2\text{Si})_3\text{MnMgMe}$, followed by the addition of iodomethane, gave a mixture of (E) -3,3-dimethyl-1-dimethylphenylsilyl-1-butene (**4**) and 4-dimethylphenylsilyl-2-methyl-2-pentene (**5**) (**4**:**5** = 1:2) in 90% combined yield (Scheme 1). Here, we wish to report that the reaction of acetylenes or 1,3-dienes with $\text{PhMe}_2\text{SiMgMe}$ in the presence of a catalytic amount of manganese(II) chloride has provided different types of products from those obtained by means of a stoichiometric reagent, $(\text{PhMe}_2\text{Si})_3\text{MnMgMe}$.


(1) Manganese(II) Chloride Catalyzed Silylmagnesiation of Acetylenes. We have reported on the simultaneous cis addition of a silyl group and a metal to acetylenes with regioselectivity using PhMe_2SiLi and metal compounds, such as MeMgI , Et_2AlCl , ZnBr_2 , and Et_3B , in the presence of a transition-metal catalyst.^{2,3)} Platinum- or cop-

per-catalyzed silylmagnesiation ($\text{PhMe}_2\text{SiMgMe}$) of 1-dodecyne, followed by aqueous quenching, provided (E) -1-dimethylphenylsilyl-1-dodecene exclusively, which has the same regiochemistry as those products prepared from stoichiometric silylcupration⁴⁾ or silyltitanation.⁵⁾ On the other hand, palladium-catalyzed silylaluminum ($\text{PhMe}_2\text{SiAlEt}_2$) afforded the regioisomer, 2-dimethylphenylsilyl-1-dodecene, with high regioselectivity (Scheme 2).

The treatment of 4-benzyloxy-1-butyne (**1a**, 1.0 mmol) with $\text{PhMe}_2\text{SiMgMe}$, prepared from PhMe_2SiLi (1.6 mmol)⁶⁾ and MeMgI (1.6 mmol) in situ, in the presence of a catalytic amount of MnCl_2 (0.08 mmol)^{7,8)} at 0 °C, gave a mixture of monosilylated products, **6a** and **7a**, in 90% combined yield after an aqueous workup.⁹⁾ The reaction proceeded with high regio- and stereoselectivities. (E) -4-Benzyloxy-1-dimethylphenylsilyl-1-butene (**6a**) was obtained as the major product, and only a small amount of the regioisomer, 4-benzyloxy-2-dimethylphenylsilyl-1-butene (**7a**), was detected in the reaction mixture (**6a**/**7a** > 95/5) (Scheme 3). The stereoisomer, (Z) -4-benzyloxy-1-dimethylphenylsilyl-1-butene, could not be observed in the reaction mixture. The use of tetrahydropyranyl ether of 3-butyne-1-ol (**1b**) in place of **1a** provided a regioisomeric mixture of **6b** and **7b** in 90%



Scheme 1.




Scheme 2.

presence of MnCl_2 catalyst under an argon atmosphere, followed by the addition of iodomethane, afforded 4-dimethylphenylsilyl-2,3-dimethyl-1-butene (**18a**) as a main product, along with **4** and **5** (Scheme 6).

The representative results of the reaction of 1,3-butadiene and 2,3-dimethyl-1,3-butadiene as well as isoprene are summarized in Table 1. Several comments are worth making: (1) The distribution of the products heavily depended on the reaction temperature. For instance, when iodomethane was added at -78°C to silyl-substituted allylic magnesium compound, prepared from isoprene and $\text{PhMe}_2\text{SiMgMe}$ at 0°C in the presence of MnCl_2 catalyst, 4-dimethylphenylsilyl-2,3-dimethyl-1-butene (**18a**) was obtained exclusively (Entry 1). In contrast, the addition of iodomethane at 0°C provided 3,3-dimethyl-1-dimethylphenylsilyl-1-butene (**4**) and 4-dimethylphenylsilyl-2-methyl-2-pentene (**5**) in addition to **18a** (**18a**:**4**:**5** = 88:6:6) (Entry 2). Moreover, the addition at 25°C afforded 4-dimethylphenylsilyl-2-methyl-2-pentene (**5**) as the main product in 37% yield in addition to **18a** (14% yield) (Entry 3). To prevent the formation of a complex isomeric mixture, other electrophiles, such as allyl bromide and carbonyl compounds, were added at -78°C (Entry 4—15 except 13). (2) The addition of aldehyde or

ketone at -78°C to the intermediary magnesium compound, derived from manganese-catalyzed silylmagnesiation of isoprene, afforded homoallylic silane derivatives **18e**, **18f**, and **18g** exclusively. In contrast, quenching with CH_3COOH , even at -78°C , gave a mixture of 4-dimethylphenylsilyl-2-methyl-1-butene (**18c**, 70%) and 1-dimethylphenylsilyl-3-methyl-2-butene (**19c**, 8%) (Entry 5). The addition of D_2O at -78°C also provided an isomeric mixture of **18d** and **19d** (**18d**/**19d** = 77/23). (3) Manganese salts, such as $\text{Mn}_2(\text{CO})_{10}$ and $\text{CH}_3\text{C}_5\text{H}_4\text{Mn}(\text{CO})_3$, were also effective as MnCl_2 . For example, the treatment of isoprene (1.0 mmol) with silylmagnesium reagent (1.6 mmol) in the presence of $\text{Mn}_2(\text{CO})_{10}$ and $\text{CH}_3\text{C}_5\text{H}_4\text{Mn}(\text{CO})_3$ (0.08 mmol each) provided a mixture of 4-dimethylphenylsilyl-2-methyl-1-butene (**18c**) and 1-dimethylphenylsilyl-3-methyl-2-butene (**19c**) in 90% (**18c**/**19c** = 62/38) and 70% (**18c**/**19c** = 72/28) combined yields, respectively, after quenching with H_2O at -78°C . (4) The reagent PhMe_2SiLi or $\text{PhMe}_2\text{SiMgMe}$ did not react with isoprene without manganese(II) chloride, which suggests that the manganese salts played a critical role for the silylmagnesiation reaction. The addition of MeMgI was also essential for a successful reaction. The treatment of isoprene with PhMe_2SiLi instead of $\text{PhMe}_2\text{SiMgMe}$ in the presence




Scheme 6.


Table 1. Manganese-Catalyzed Silylmagnesiation of 1,3-Dienes^{a)}

$\begin{array}{c} \text{R}^1 \\ \diagup \\ \text{C} \\ \diagdown \\ \text{R}^2 \end{array} \xrightarrow[2) \text{ Electrophile}]{1) \text{ SiMgMe}/\text{cat. MnCl}_2} \begin{array}{c} \text{R}^1 \\ \diagup \\ \text{C} \\ \diagdown \\ \text{R}^2 \end{array} \begin{array}{c} \text{R}^2 \\ \diagup \\ \text{C} \\ \diagdown \\ \text{E} \end{array} \text{Si} + \begin{array}{c} \text{R}^1 \\ \diagup \\ \text{C} \\ \diagdown \\ \text{E} \end{array} \begin{array}{c} \text{R}^2 \\ \diagup \\ \text{C} \\ \diagdown \\ \text{Si} \end{array} + \begin{array}{c} \text{R}^1 \\ \diagup \\ \text{C} \\ \diagdown \\ \text{E} \end{array} \begin{array}{c} \text{R}^2 \\ \diagup \\ \text{C} \\ \diagdown \\ \text{Si} \end{array} + \begin{array}{c} \text{R}^1 \\ \diagup \\ \text{C} \\ \diagdown \\ \text{E} \end{array} \begin{array}{c} \text{R}^2 \\ \diagup \\ \text{C} \\ \diagdown \\ \text{Si} \end{array}$ $\text{Si} = \text{SiMe}_2\text{Ph}$				Product / %			
Entry	R ¹	R ²	Electrophile	18	19	20	21
1	Me	H	MeI	83 (18a)	0	0	0
2			MeI ^{b)}	86	0	6 (4)	6 (5)
3			MeI ^{c)}	14	0	2	37
4			$\text{CH}_2=\text{CHCH}_2\text{Br}$	83 (18b)	0	0	0
5			CH_3COOH	70 (18c)	8 (19c)	0	0
6			D_2O	74 (18d)	22 (19d)	0	0
7			<i>n</i> -BuCHO	88 (18e)	0	0	<2
8			PhCHO	95 (18f)	0	0	<2
9			CH_3COCH_3	90 (18g)	0	0	<2
10	H	H	MeI	63 (18h)	0	0	10 (21h)
11			CH_3COCH_3	80 (18i)	0	0	0
12			$\text{D}_2\text{O}^{\text{d)}$	48 (18j)	47 (19j)	0	0
13			$\text{H}_2\text{O}^{\text{e)}$	7 (18k)	19k ^{f)}	39 (20k)	21k ^{f,g)}
14	Me	Me	MeI	53 (18l)	0	0	0
15			CH_3COCH_3	47 (18m)	0	0	0

a) $\text{PhMe}_2\text{SiMgMe}$ and 1,3-diene were stirred at 0°C in the presence of MnCl_2 catalyst for 10 min and then electrophiles were added at -78°C unless otherwise noted. b) MeI was added at 0°C . c) The reaction between $\text{PhMe}_2\text{SiMgMe}$ and isoprene was performed at 25°C for 0.5 h and then MeI was added at 25°C . d) D_2O was added at -78°C . e) H_2O was added at 0°C . f) **19k**+**21k** = 54%. g) *E/Z* = 70/30.



Scheme 7.



Scheme 8.

of a catalytic amount of MnCl_2 gave 4-dimethylphenylsilyl-2-methyl-1-butene in only < 20% yield after quenching with water. (5) 1,3-Butadiene was equally reactive toward a MnCl_2 -catalyzed silylmagnesiation reaction, and 2,3-dimethyl-1,3-butadiene was less reactive than 1,3-butadiene and isoprene. 1,3-Pentadiene and 1,3-cyclooctadiene were too sluggish to react with $\text{PhMe}_2\text{SiMgMe}$ in the presence of a MnCl_2 catalyst.

We are tempted to assume the following reaction mechanism for stoichiometric silylmanganation and manganese-catalyzed silylmagnesiation reactions of isoprene (Scheme 7). In the stoichiometric reaction with $(\text{PhMe}_2\text{Si})_3\text{MnMgMe}$, regioselective silylmanganation of isoprene occurs to give silylated allylmanganese compound **22** in the first step of the reaction sequence. Then, a β -elimination of Mn-H from **22**, followed by hydromanganation (1,4-fashion), generates a more stable allylmanganese species **23**. The addition of iodomethane to **23** affords two isomeric products, **4** and **5**. Meanwhile, the catalytic reaction proceeds as follows. Manganese-catalyzed silylmagnesiation takes place regioselectively to introduce a silyl group at the least-hindered position, giving an allylmagnesium compound **24**. The allylic metal species **24** reacts at a secondary carbon with electrophiles, such as iodomethane and carbonyl compounds, regioselectively at -78°C to give **18**. However, at a higher temperature, such as 25°C , a partial manganese-

catalyzed isomerization of **24** into **25** competes with the reaction between **24** and iodomethane, and a complex isomeric mixture containing **18**, **20**, and **21** is obtained upon a treatment with electrophiles. Heating a solution of **24** accelerated the isomerization, which was exemplified by the following experiment. The mixture of isoprene and $\text{PhMe}_2\text{SiMgMe}$ containing a MnCl_2 catalyst was stirred for 10 min at 0°C , and then heated at 80°C for 30 min. The addition of iodomethane to the resulting mixture at 25°C provided a mixture of three isomers, **18a**, **4**, and **5** (**18a**:**4**:**5** = 11:12:77), in 40% combined yield.

The manganese-catalyzed isomerization by β -elimination and a subsequent readdition of Mn-H took place more easily in the adduct of 1,3-butadiene with $\text{PhMe}_2\text{SiMgMe}$ compared to the adduct derived from isoprene and $\text{PhMe}_2\text{SiMgMe}$. For instance, when D_2O was added at -78°C to allylic magnesium compound **26**, prepared from butadiene and $\text{PhMe}_2\text{SiMgMe}$ at 0°C in the presence of MnCl_2 catalyst, a mixture of **18j** and **19j** (**18j**:**19j** = 1:1) was obtained (Entry 12). In contrast, an addition of H_2O at 0°C afforded **20k** and **21k** in addition to **18k** and **19k** (**18k**:**19k**+**21k**:**20k** = 7:54:39). Moreover, two stereoisomers could be prepared separately by changing the reaction temperature without any contamination by the other isomer in the reaction of 1,3-butadiene and acetone. The addition of acetone to the reaction mixture, derived from 1,3-butadiene and silylmagnesium in

the presence of MnCl_2 at -78°C , afforded 2-methyl-3-[(dimethylphenylsilyl)methyl]-4-penten-2-ol (**18i**) as a single product (Entry 11 in Table 1). In contrast, 2,3-dimethyl-5-dimethylphenylsilyl-4-penten-2-ol (**20i**) was obtained exclusively upon the addition of acetone at 25°C (Scheme 8).

Experimental

Distillation of the products was performed using Kugelrohr (Büchi); the boiling points are indicated by the air-bath temperature values without any correction. The NMR spectra (^1H and ^{13}C) were recorded on a Varian GEMINI 300 spectrometer in CDCl_3 ; tetramethylsilane (TMS) was used as an internal standard. The IR spectra were determined on a JASCO IR-810 spectrometer. The analyses were carried out at the Elemental Analysis Center of Kyoto University.

General Procedure for the Reaction of Acetylene with $\text{PhMe}_2\text{SiMgMe}$ in the Presence of a Catalytic Amount of MnCl_2 . The reaction of 3-decyn-1-ol benzyl ether (**8**) with $\text{PhMe}_2\text{SiMgMe}$ is representative. A tetrahydrofuran (THF) solution of PhMe_2SiLi (0.7 M, 2.2 mL, 1.6 mmol, $M = \text{mol dm}^{-3}$) and MeMgI (1.0 M, ether solution, 1.6 mL, 1.6 mmol) was added to a suspension of MnCl_2 (10 mg, 0.08 mmol) in THF (2 mL) at 0°C under an argon atmosphere. After being stirred for 30 min at 0°C , a THF solution of **8** (0.24 g, 1.0 mmol) was added, and the resulting mixture was stirred for 2 h at 0°C and warmed up to 25°C . The mixture was poured into ice-cooled water and extracted with ethyl acetate (20 mL $\times$ 3). The combined organic layer was dried over Na_2SO_4 and concentrated in vacuo. The residual oil was submitted to silica-gel column chromatography (hexane/ethyl acetate = 20/1) to give a mixture of (*E*)-1-benzyloxy-4-dimethylphenylsilyl-3-decene (**9**) and its regioisomer (**10**) in 95% yield (0.36 g, **9/10** = 73/27).

(*E*)-1-Benzyloxy-4-dimethylphenylsilyl-3-decene (9**) and (*E*)-1-Benzyloxy-3-dimethylphenylsilyl-3-decene (**10**) (73 : 27):** IR (neat) 2952, 2924, 2854, 1110, 812, 730, 697 cm^{-1} ; ^1H NMR (CDCl_3) δ = 0.30 (s, 1.62H), 0.32 (s, 4.38H), 0.82 (t, J = 6.9 Hz, 2.19H), 0.85 (t, J = 6.9 Hz, 0.81H), 1.06—1.35 (m, 8H), 2.09 (m, 2H), 2.46 (dt, J = 6.9, 6.9 Hz, 2H), 3.22 (t, J = 8.1 Hz, 0.54H), 3.49 (t, J = 6.9 Hz, 1.46H), 4.36 (s, 0.54H), 4.51 (s, 1.46H), 5.80 (t, J = 6.9 Hz, 0.73H), 5.87 (t, J = 6.6 Hz, 0.27H), 7.32 (m, 8H), 7.48 (m, 2H); ^{13}C NMR (CDCl_3 , major isomer) δ = -2.79, 13.95, 22.49, 29.22, 29.54, 29.89, 30.02, 31.53, 69.83, 72.85, 127.59, 127.67, 128.44, 128.80, 134.07, 133.77, 138.64, 139.16, 141.61. Found: C, 78.60; H, 9.56%. Calcd for $\text{C}_{25}\text{H}_{36}\text{OSi}$: C, 78.88; H, 9.53%.

(*E*)-1-Benzyloxy-3-deuterio-4-dimethylphenylsilyl-3-decene (9-d**) and (*E*)-1-Benzyloxy-4-deuterio-3-dimethylphenylsilyl-3-decene (**10-d**) (73 : 27):** IR (neat) 2952, 2924, 2852, 1598, 1248, 1109, 830, 814, 771, 731, 697 cm^{-1} ; ^1H NMR (CDCl_3) δ = 0.30 (s, 1.62H), 0.32 (s, 4.38H), 0.82 (t, J = 6.9 Hz, 2.19H), 0.85 (t, J = 6.9 Hz, 0.81H), 1.06—1.35 (m, 8H), 2.09 (m, 2H), 2.46 (t, J = 7.2 Hz, 2H), 3.22 (t, J = 8.1 Hz, 0.54H), 3.49 (t, J = 6.9 Hz, 1.46H), 4.36 (s, 0.54H), 4.51 (s, 1.46H), 7.32 (m, 8H), 7.48 (m, 2H); ^{13}C NMR (CDCl_3 , major isomer) δ = -2.78, 13.94, 22.49, 29.13, 29.55, 29.90, 30.00, 31.54, 69.84, 72.86, 127.59, 127.68, 128.45, 128.80, 134.08, 138.67, 139.19, 141.48. Found: C, 78.68; H+D, 9.91%. Calcd for $\text{C}_{25}\text{H}_35\text{DOSi}$: C, 78.68; H+D, 9.69%.

4-[(2-Benzyloxy)ethyl]-5-dimethylphenylsilyl-1,4-undecadiene (12**) and 7-Benzyloxy-5-dimethylphenylsilyl-4-hexyl-1,4-heptadiene (**13**) (80 : 20):** IR (neat) 2952, 2924, 2852, 1455, 1428, 1249, 1109, 832, 814, 771, 729, 698 cm^{-1} ; ^1H NMR (CDCl_3) δ = 0.340 (s, 1.2H), 0.344 (s, 4.8H), 0.86 (t, J = 6.9 Hz, 3H), 1.23 (m, 8H), 2.13 (m, 2H), 2.49 (t, J = 7.5 Hz, 1.6H), 2.54 (t, J = 7.5

Hz, 0.4H), 2.73 (d, J = 6.3 Hz, 2H), 3.50 (t, J = 7.5 Hz, 1.6H), 3.52 (t, J = 7.5 Hz, 0.4H), 4.44 (s, 0.4H), 4.50 (s, 1.6H), 4.85 (m, 1H), 4.89 (m, 1H), 5.45 (m, 1H), 7.30 (m, 8H), 7.47 (m, 2H); ^{13}C NMR (CDCl_3 , major isomer) δ = -0.29, 13.97, 22.59, 29.64, 31.05, 31.63, 32.35, 41.75, 69.15, 72.75, 116.25, 127.56, 127.66, 127.75, 128.43, 128.64, 133.83, 134.11, 135.69, 136.80, 138.68, 140.65. Found: C, 79.73; H, 9.71%. Calcd for $\text{C}_{26}\text{H}_{36}\text{SiO}_2$: C, 79.95; H, 9.51%.

4-[(2-Benzyloxy)ethyl]-3-hexyl-2,2-dimethyl-5-phenyl-1-oxa-2-silacyclopent-3-ene (16a**) and 3-[(2-Benzyloxy)ethyl]-4-hexyl-2,2-dimethyl-5-phenyl-1-oxa-2-silacyclopent-3-ene (**17a**) (80 : 20):** IR (neat) 3026, 2952, 2922, 2852, 1455, 1249, 1098, 1079, 1054, 1027, 863, 833, 779, 733, 697 cm^{-1} ; ^1H NMR (CDCl_3) δ = 0.21 (s, 0.6H), 0.29 (s, 2.4H), 0.29 (s, 0.6H), 0.37 (s, 2.4H), 0.88 (t, J = 6.9 Hz, 2.4H), 0.95 (m, 0.6H), 1.15—1.50 (m, 8H), 1.50—1.65 (m, 0.2H), 2.08 (ddd, J = 6.6, 8.5, 14.1 Hz, 0.8H), 2.13 (m, 0.2H), 2.25 (t, J = 7.5 Hz, 1.6H), 2.47 (ddd, J = 6.0, 8.5, 14.1 Hz, 0.8H), 2.55—2.65 (m, 0.4H), 3.13 (dt, J = 6.0, 8.5 Hz, 0.8H), 3.32 (dt, J = 6.6, 8.5 Hz, 0.8H), 3.55 (t, J = 6.6 Hz, 0.4H), 4.38 (s, 1.6H), 4.55 (s, 0.4H), 5.45 (s, 0.2H), 5.49 (m, 0.8H), 7.19—7.60 (m, 10H); ^{13}C NMR (CDCl_3 , major isomer) δ = 0.72, 1.54, 13.97, 22.54, 28.03, 28.16, 29.49, 30.34, 31.63, 68.37, 72.6, 86.79, 127.32, 127.52, 127.56, 127.86, 128.39, 128.52, 137.41, 138.42, 142.29, 151.33. Found: C, 76.41; H, 8.87%. Calcd for $\text{C}_{26}\text{H}_{36}\text{SiO}_2$: C, 76.42; H, 8.87%.

4-[(2-Benzyloxyethyl)]-3-hexyl-5-(4-methoxyphenyl)-2,2-dimethyl-1-oxa-2-silacyclopent-3-ene (16b**) and 3-[(2-Benzyloxy)ethyl]-3-hexyl-5-(4-methoxyphenyl)-2,2-dimethyl-1-oxa-2-silacyclopent-3-ene (**17b**) (70 : 30):** IR (neat) 2954, 2926, 2854, 1613, 1513, 1456, 1362, 1304, 1249, 1173, 1132, 1100, 1077, 1038, 1010, 864, 827, 780, 734, 697 cm^{-1} ; ^1H NMR (CDCl_3) δ = 0.20 (s, 0.9H), 0.28 (s, 3H), 0.35 (s, 2.1H), 0.78—0.90 (m, 0.9H), 0.90 (t, J = 6.6 Hz, 2.1H), 1.10—1.48 (m, 8H), 1.58—1.70 (m, 0.3H), 2.03—2.18 (m, 0.3H), 2.08 (ddd, J = 6.6, 8.4, 14.7 Hz, 0.7H), 2.25 (t, J = 8.4 Hz, 1.4H), 2.46 (ddd, J = 6.3, 8.7, 14.7 Hz, 0.7H), 2.50—2.61 (m, 0.6H), 3.16 (dt, J = 6.3, 9 Hz, 0.7H), 3.53 (t, J = 6.6 Hz, 0.6H), 3.32 (dt, J = 6.3, 9 Hz, 0.7H), 3.77 (s, 0.9H), 3.79 (s, 2.1H), 4.39 (s, 1.4H), 4.52 (s, 0.6H), 5.41 (s, 0.3H), 5.45 (s, 0.7H), 6.82 (m, 0.6H), 6.84 (m, 1.4H), 7.13 (m, 1.4H), 7.25 (m, 0.6H), 7.22—7.38 (m, 5H); ^{13}C NMR (CDCl_3 , major isomer) δ = 0.70, 1.61, 13.99, 22.56, 28.07, 28.23, 29.52, 30.37, 31.65, 55.21, 68.40, 72.61, 86.30, 113.90, 127.55, 127.58, 128.40, 128.54, 134.59, 137.33, 138.46, 151.46, 159.34. Found: C, 73.74; H, 8.62%. Calcd for $\text{C}_{27}\text{H}_{38}\text{SiO}_3$: C, 73.94; H, 8.73%.

The physical data for **6a**, **6b**, **6c**, **7a**, **7b**, **7c**, and **11** were identical with those which have been described in the literature.^{3b)}

General Procedure for the Reaction of Silylmagnesium Reagent with 1,3-Dienes.

A THF solution of PhMe_2SiLi (0.7 M, 2.2 mL, 1.6 mmol) and MeMgI (1.0 M ether solution, 1.6 mL, 1.6 mmol) were added to a suspension of MnCl_2 (0.08 mmol) in THF (2 mL) at 0°C under an argon atmosphere. After being stirred for 30 min at 0°C , isoprene (THF solution, 1.0 mmol) was added and the whole was stirred for 10 min. The reaction flask was cooled to -78°C in a dry ice bath, and iodomethane (3.2 mmol) was added. The resulting mixture was stirred for 30 min at -78°C , and then warmed up to 25°C . An extractive workup followed by silica-gel column chromatography gave 4-dimethylphenylsilyl-2,3-dimethyl-1-butene (**18a**) in 83% yield: Bp 111—115 $^\circ\text{C}$ (3 Torr, 1 Torr = 133.322 Pa); IR (neat) 2870, 1644, 1427, 1373, 1248, 1112, 887, 812, 697 cm^{-1} ; ^1H NMR (CDCl_3) δ = 0.29 (s, 3H), 0.30 (s, 3H), 0.84 (dd, J = 7.5, 15.9 Hz, 1H), 0.97 (dd, J = 7.5, 15.9 Hz, 1H), 1.00 (d, J = 6.9 Hz, 3H), 1.64 (s, 3H), 2.36 (tq, J = 7.5, 6.9

Hz, 1H), 4.60 (s, 1H), 4.68 (s, 1H), 7.35 (m, 3H), 7.52 (m, 2H); ^{13}C NMR (CDCl_3) $\delta = -2.59, -2.36, 18.44, 22.51, 22.95, 37.38, 108.46, 127.74, 128.80, 133.65, 139.99, 152.36$. Found: C, 76.77; H, 10.18%. Calcd for $\text{C}_{14}\text{H}_{22}\text{Si}$: C, 77.05; H, 10.08%.

3-[(Dimethylphenylsilyl)methyl]-2-methyl-1,5-hexadiene (18b): Bp 125–127 °C (3 Torr); IR (neat) 2904, 1642, 1427, 1373, 1248, 1112, 991, 889, 698 cm^{-1} ; ^1H NMR (CDCl_3) $\delta = 0.27$ (s, 3H), 0.29 (s, 3H), 0.92 (d, $J = 7.8$ Hz, 2H), 1.57 (m, 3H), 2.06 (m, 2H), 2.29 (tt, $J = 7.2, 7.2$ Hz, 1H), 4.66 (s, 2H), 4.93 (m, 2H), 5.64 (m, 1H), 7.35 (m, 3H), 7.52 (m, 2H); ^{13}C NMR (CDCl_3) $\delta = -2.87, -2.29, 17.83, 20.19, 41.24, 43.23, 111.09, 115.31, 127.72, 128.81, 133.66, 137.49, 139.86, 148.87$. Found: C, 78.54; H, 10.11%. Calcd for $\text{C}_{16}\text{H}_{24}\text{Si}$: C, 78.67; H, 9.83%.

4-Dimethylphenylsilyl-2-methyl-1-butene (18c): Bp 104–106 °C (3 Torr); IR (neat) 2888, 1649, 1427, 1374, 1248, 1171, 1113, 883, 811, 771, 697 cm^{-1} ; ^1H NMR (CDCl_3) $\delta = 0.28$ (s, 6H), 0.90 (m, 2H), 1.71 (s, 3H), 2.00 (m, 2H), 4.66 (s, 1H), 4.69 (s, 1H), 7.35 (m, 3H), 7.52 (m, 2H); ^{13}C NMR (CDCl_3) $\delta = -3.28, 13.63, 22.13, 31.71, 108.48, 127.83, 128.93, 133.66, 139.35, 148.51$. Found: C, 76.46; H, 10.09%. Calcd for $\text{C}_{13}\text{H}_{20}\text{Si}$: C, 76.45; H, 9.79%.

3-Deuterio-4-dimethylphenylsilyl-2-methyl-1-butene (18d) and 1-Deuterio-4-dimethylphenylsilyl-2-methyl-2-butene (19d) (78 : 22): Bp 104–106 °C (3 Torr); IR (neat) 2956, 2910, 1649, 1427, 1248, 1113, 873, 837, 698 cm^{-1} ; ^1H NMR (CDCl_3) $\delta = 0.24$ (s, 1.32H), 0.26 (s, 4.68H), 0.87 (d, $J = 9.3$ Hz, 1.56H), 1.46 (m, 0.44H), 1.61 (d, $J = 8.4$ Hz, 0.44H), 1.67 (s, 0.66H), 1.69 (s, 2.34H), 1.97 (t, $J = 9.3$ Hz, 0.78H), 4.65 (s, 0.78H), 4.68 (s, 0.78H), 5.14 (t, $J = 8.4$ Hz, 0.22H), 7.35 (m, 3H), 7.50 (m, 2H); ^{13}C NMR (CDCl_3 , major isomer) $\delta = -3.28, 13.53, 22.10, 31.35$ (t, $J = 19.5$ Hz), 108.49, 127.83, 128.93, 133.66, 139.36, 148.51. Found: C, 75.96; H+D, 10.45%. Calcd for $\text{C}_{13}\text{H}_{19}\text{DSi}$: C, 76.08; H+D, 10.23%.

3-[(Dimethylphenylsilyl)methyl]-2-methyl-1-octen-4-ol (erythrolthreo=1 : 1 mixture) (18e): Bp 160 °C (0.5 Torr, decomp); IR (neat) 3410, 2954, 2858, 1643, 1427, 1248, 1112, 837, 730, 698 cm^{-1} ; ^1H NMR (CDCl_3) $\delta = 0.24$ (s, 1.5H), 0.25 (s, 1.5H), 0.26 (s, 1.5H), 0.28 (s, 1.5H), 0.83–1.59 (m, 15H), 2.13 (m, 1H), 3.34 (m, 1H), 4.72 (s, 0.5H), 4.77 (s, 0.5H), 4.78 (s, 0.5H), 4.87 (m, 0.5H), 7.33 (m, 3H), 7.47 (m, 2H); ^{13}C NMR (CDCl_3) $\delta = -3.18, -2.96, -2.31, -2.05, 13.95, 13.99, 14.03, 15.78, 18.73, 20.57, 22.59, 22.74, 27.74, 28.18, 33.72, 34.13, 48.80, 49.61, 73.63, 74.18, 112.93, 115.20, 127.72, 127.76, 128.83, 128.94, 133.64, 133.68, 139.44, 139.78, 146.55, 147.17$. Found: C, 74.40; H, 10.41%. Calcd for $\text{C}_{18}\text{H}_{30}\text{OSi}$: C, 74.50; H, 10.34%.

2-[(Dimethylphenylsilyl)methyl]-3-methyl-1-phenyl-3-buten-1-ol (18f): Bp 160 °C (0.5 Torr, decomp); IR (neat) 3404, 2896, 1492, 1427, 1248, 1181, 1112, 999, 814, 698 cm^{-1} ; ^1H NMR (CDCl_3) $\delta = 0.15$ (s, 1.5H), 0.17 (s, 1.5H), 0.20 (s, 1.5H), 0.21 (s, 1.5H), 0.52 (dd, $J = 3, 15$ Hz, 0.5H), 0.81 (dd, $J = 12.3, 15$ Hz, 0.5H), 1.00 (dd, $J = 1.5, 15$ Hz, 0.5H), 1.14 (dd, $J = 3.0, 15$ Hz, 0.5H), 1.55 (s, 1.5H), 1.66 (s, 1.5H), 1.96 (s, 0.5H), 2.21 (s, 0.5H), 2.45 (m, 1H), 4.31 (d, $J = 9$ Hz, 0.5H), 4.54 (d, $J = 6$ Hz, 0.5H), 4.69 (m, 0.5H), 4.77 (m, 0.5H), 4.92 (m, 0.5H), 4.98 (m, 0.5H), 7.29 (m, 3H), 7.37 (m, 2H); ^{13}C NMR (CDCl_3) $\delta = -3.34, -2.90, -2.33, -2.03, 13.37, 15.14, 17.83, 21.46, 50.26, 51.67, 76.67, 77.08, 113.34, 116.25, 126.35, 127.22, 127.58, 127.66, 127.67, 127.83, 128.12, 128.36, 128.77, 128.87, 133.60, 133.65, 139.03, 139.56, 142.42, 143.04, 146.15, 146.51$. Found: C, 77.59; H, 8.67%. Calcd for $\text{C}_{20}\text{H}_{26}\text{OSi}$: C, 77.42; H, 8.38%.

3-[(Dimethylphenylsilyl)methyl]-2,4-dimethyl-4-penten-2-ol (18g): Bp 135–137 °C (1 Torr); IR (neat) 3444, 2894, 1638, 1427, 1375, 1248, 1111, 945, 818, 728, 698 cm^{-1} ; ^1H NMR (CDCl_3)

$\delta = 0.26$ (s, 3H), 0.29 (s, 3H), 1.02 (m, 2H), 1.15 (s, 6H), 1.51 (s, 1H), 1.66 (m, 3H), 2.16 (m, 1H), 4.76 (m, 1H), 4.85 (m, 1H), 7.35 (m, 3H), 7.52 (m, 2H); ^{13}C NMR (CDCl_3) $\delta = -3.26, -2.23, 14.54, 21.83, 27.61, 27.75, 53.28, 72.89, 114.95, 127.72, 128.87, 133.66, 139.56, 146.87$. Found: C, 73.27; H, 10.19%. Calcd for $\text{C}_{16}\text{H}_{26}\text{OSi}$: C, 73.27; H, 9.91%.

4-Dimethylphenylsilyl-3-methyl-1-butene (18h): Bp 99–100 °C (3 Torr); IR (neat) 2874, 1638, 1248, 1113, 908, 786, 697 cm^{-1} ; ^1H NMR (CDCl_3) $\delta = 0.29$ (s, 3H), 0.30 (s, 3H), 0.83 (dd, $J = 7.2, 14.7$ Hz, 1H), 0.93 (dd, $J = 7.2, 14.7$ Hz, 1H), 1.00 (d, $J = 6.6$ Hz, 3H), 2.32 (m, 1H), 4.82 (m, 1H), 4.90 (m, 1H), 5.74 (ddd, $J = 7.2, 9.9, 17.4$ Hz, 1H), 7.35 (m, 3H), 7.52 (m, 2H); ^{13}C NMR (CDCl_3) $\delta = -2.30, -2.02, 23.77, 23.82, 34.15, 111.09, 127.76, 128.80, 133.65, 136.90, 146.98$. Found: C, 76.22; H, 9.93%. Calcd for $\text{C}_{13}\text{H}_{20}\text{Si}$: C, 76.48; H, 9.80%.

3-[(Dimethylphenylsilyl)methyl]-2-methyl-4-penten-2-ol (18i): Bp 133–135 °C (1 Torr); IR (neat) 3552, 3424, 3066, 2968, 2886, 1638, 1427, 1372, 1249, 1156, 1113, 1000, 914, 832, 728, 699 cm^{-1} ; ^1H NMR (CDCl_3) $\delta = 0.25$ (s, 3H), 0.27 (s, 3H), 0.75 (dd, $J = 12, 14.7$ Hz, 1H), 1.03 (dd, $J = 2.4, 14.7$ Hz, 1H), 1.10 (s, 3H), 1.12 (s, 3H), 1.55 (s, 1H), 2.12 (ddd, $J = 2.4, 9.9, 12$ Hz, 1H), 5.04 (ddd, $J = 1.8, 10.2, 18$ Hz, 2H), 5.54 (ddd, $J = 9.9, 10.2, 18$ Hz, 1H), 7.35 (m, 3H), 7.52 (m, 2H); ^{13}C NMR (CDCl_3) $\delta = -2.87, -1.90, 16.02, 26.11, 26.22, 51.75, 72.69, 117.85, 127.75, 128.88, 133.68, 136.60, 140.67$. Found: C, 72.52; H, 9.93%. Calcd for $\text{C}_{15}\text{H}_{24}\text{OSi}$: C, 72.60; H, 9.67%.

A Mixture of 3-Deuterio-4-dimethylphenylsilyl-1-butene (18j), (E)-1-Deuterio-4-dimethylphenylsilyl-2-butene (19j-E), and (Z)-1-Deuterio-4-dimethylphenylsilyl-2-butene (19j-Z) (50 : 39 : 11): Bp 83–85 °C (3 Torr); IR (neat) 2954, 2902, 1638, 1427, 1249, 1113, 836, 728, 697, 459, 442, 418 cm^{-1} ; ^1H NMR (CDCl_3) $\delta = 0.24$ (s, 0.66H), 0.26 (s, 5.34H), 0.83 (d, $J = 8.1$ Hz, 1H), 1.48 (m, 0.78H), 1.58 (m, 0.22H), 1.63 (d, $J = 8.1$ Hz, 0.22H), 1.70 (d, $J = 7.2$ Hz, 0.78H), 2.04 (m, 0.5H), 4.87 (ddd, $J = 10.2, 1.2, 1.2$ Hz, 0.5H), 4.97 (ddd, $J = 17.1, 1.2, 1.2$ Hz, 0.5H), 5.37 (m, 1H), 5.86 (ddd, $J = 6.0, 10.2, 17.1$ Hz, 0.5H), 7.34 (m, 3H), 7.49 (m, 2H); ^{13}C NMR (CDCl_3 , **18j** and **19j-E**) $\delta = -3.45, -3.18, 12.19$ (t, $J = 19.5$ Hz), 14.59, 17.02, 27.48 (t, $J = 19.5$ Hz), 112.88, 122.04, 125.87, 127.78, 127.82, 128.93, 129.01, 133.65, 133.71, 139.14, 139.35, 141.58. Found: C, 75.27; H+D, 10.02%. Calcd for $\text{C}_{12}\text{H}_{17}\text{DSi}$: C, 75.45; H+D, 9.95%.

A Mixture of (E)-1-Dimethylphenylsilyl-2-butene (21k-E), (Z)-1-Dimethylphenylsilyl-2-butene (21k-Z), and (E)-1-Dimethylphenylsilyl-1-butene (20k) (43 : 17 : 40): Bp 83–85 °C (3 Torr); IR (neat) 2952, 1248, 1113, 824, 696 cm^{-1} ; ^1H NMR (CDCl_3) $\delta = 0.26$ (s, 1.02H), 0.29 (s, 2.58H), 0.33 (s, 2.40H), 1.02 (t, $J = 7.5$ Hz, 1.2H), 1.51 (d, $J = 5.7$ Hz, 1.29H), 1.64 (d, $J = 5.7$ Hz, 0.51H), 1.66 (d, $J = 6.9$ Hz, 0.34H), 1.73 (d, $J = 7.5$ Hz, 0.86H), 2.17 (qd, $J = 7.5, 5.7$ Hz, 0.8H), 5.41 (m, 1.2H), 5.76 (d, $J = 18.6$ Hz, 0.4H), 6.19 (dt, $J = 18.6, 5.7$ Hz, 0.4H), 7.35 (m, 3H), 7.53 (m, 2H); ^{13}C NMR (CDCl_3) $\delta = -3.54, -3.45, -2.60, 12.47, 12.60, 17.01, 17.93, 21.46, 29.35, 112.85, 122.07, 124.02, 125.85, 125.95, 126.44, 127.76, 127.78, 127.83, 128.87, 128.95, 129.01, 133.65, 133.71, 133.91, 139.13, 139.47, 150.84$. Found: C, 75.84; H, 9.57%. Calcd for $\text{C}_{12}\text{H}_{18}\text{Si}$: C, 75.80; H, 9.47%.

4-Dimethylphenylsilyl-2,3,3-trimethyl-1-butene (18l): Bp 113–115 °C (3 Torr); IR (neat) 2958, 1636, 1427, 1376, 1248, 1190, 1112, 890, 816, 698 cm^{-1} ; ^1H NMR (CDCl_3) $\delta = 0.29$ (s, 6H), 1.06 (s, 6H), 1.07 (s, 2H), 1.69 (s, 3H), 4.63 (m, 1H), 4.70 (m, 1H), 7.35 (m, 3H), 7.52 (m, 2H); ^{13}C NMR (CDCl_3) $\delta = -1.27, 19.63, 29.16, 29.90, 30.44, 38.41, 108.28, 127.68, 128.61, 133.59, 140.94, 154.20$. Found: C, 77.31; H, 10.42%. Calcd for $\text{C}_{15}\text{H}_{24}\text{Si}$:

C, 77.57; H, 10.33%.

3-[(Dimethylphenylsilyl)methyl]-2,3,4-trimethyl-4-penten-2-ol (18m): Bp 139–141 °C (1 Torr); IR (neat) 3462, 2974, 1628, 1427, 1369, 1248, 1111, 1044, 947, 892, 815, 698 cm^{-1} ; ^1H NMR (CDCl_3) δ = 0.27 (s, 3H), 0.32 (s, 3H), 0.99 (d, J = 14.4 Hz, 1H), 1.07 (s, 3H), 1.12 (s, 3H), 1.17 (s, 3H), 1.54 (d, J = 14.4 Hz, 1H), 1.69 (s, 1H), 1.80 (s, 3H), 4.84 (s, 1H), 5.00 (s, 1H), 7.35 (m, 3H), 7.52 (m, 2H); ^{13}C NMR (CDCl_3) δ = -1.02, -0.82, 22.41, 23.24, 23.85, 25.45, 25.58, 48.20, 75.28, 114.54, 127.71, 128.65, 133.60, 141.10, 150.39. Found: C, 73.98; H, 10.42%. Calcd for $\text{C}_{17}\text{H}_{28}\text{OSi}$: C, 73.93; H, 10.14%.

The physical data for **19c**, **20i**, and **21h** were identical with those which have been described in the literature.^{1c)}

References

- 1) a) J. Hibino, S. Nakatsukasa, K. Fugami, S. Matsubara, K. Oshima, and H. Nozaki, *J. Am. Chem. Soc.*, **107**, 6416 (1985); b) K. Fugami, S. Nakatsukasa, K. Oshima, K. Utimoto, and H. Nozaki, *Chem. Lett.*, **1986**, 869; c) K. Fugami, J. Hibino, S. Nakatsukasa, S. Matsubara, K. Oshima, K. Utimoto, and H. Nozaki, *Tetrahedron*, **44**, 4277 (1988).
- 2) H. Hayami, M. Sato, S. Kanemoto, Y. Morizawa, K. Oshima, and H. Nozaki, *J. Am. Chem. Soc.*, **105**, 4491 (1983).
- 3) a) K. Nozaki, K. Wakamatsu, T. Nonaka, W. Tückmantel, K. Oshima, and K. Utimoto, *Tetrahedron Lett.*, **27**, 2007 (1986); b) K. Wakamatsu, T. Nonaka, Y. Okuda, W. Tückmantel, K. Oshima, K. Utimoto, and H. Nozaki, *Tetrahedron*, **42**, 4427 (1986).
- 4) I. Fleming, T. W. Newton, and F. Roessler, *J. Chem. Soc., Perkin Trans. 1*, **1981**, 2527.
- 5) K. Tamao, M. Akita, R. Kanatani, N. Ishida, and M. Kumada, *J. Organomet. Chem.*, **226**, C9 (1982).
- 6) H. Gilman and G. D. Lichtenwalter, *J. Am. Chem. Soc.*, **80**, 607 (1958).
- 7) Allylmagnesiation of acetylenes catalyzed by manganese salts has been reported. K. Okada, K. Oshima, and K. Utimoto, *J. Am. Chem. Soc.*, **118**, 6076 (1996).
- 8) Three different types of organomanganese(II) reagents (RMnX , R_2Mn , $\text{R}_3\text{Mn}^-\text{Mtl}^+$) have been reported. J. F. Normant and G. Cahiez, *Organomanganese Reagents*. In "Modern Synthetic Methods," ed by R. Scheffold, Otto Salle Verlag GmbH & Co., Frankfurt am Main (1983), Vol. 3.
- 9) We assume that the same silylmanganese reagent, $(\text{PhMe}_2\text{Si})_3\text{MnMgMe}$ is an active species in a catalytic reaction.
- 10) Phenylated adducts Ph_2CHOH and $p\text{-MeOC}_6\text{H}_4\text{CH}(\text{OH})\text{Ph}$ were detected in the reaction mixture.
- 11) H. Oda, M. Sato, Y. Morizawa, K. Oshima, and H. Nozaki, *Tetrahedron*, **41**, 3257 (1985).