

An Efficient Synthesis of Racemic Necine Bases from a Common Intermediate

Haruki Niwa,^{*,†} Akio Kuroda, Tomoyo Sakata, and Kiyoyuki Yamada

Department of Chemistry, Faculty of Science, Nagoya University, Chikusa, Nagoya 464

[†]Department of Applied Physics and Chemistry, The University of Electro-Communications, Chofu 182


(Received April 30, 1997)

Necine bases of pyrrolizidine alkaloids, turneforcidine, hastanecine, and platynecine are synthesized from ethyl 1-hydroxy-2,3,5,6-tetrahydro-1*H*-pyrrolizine-7-carboxylate in racemic form.


Pyrrolizidine alkaloids are known to possess remarkable hepatotoxicity and, in certain cases, carcinogenicity and antitumor activity.¹⁾ The characteristic structures coupled with diverse biological activities have made these alkaloids attractive synthetic targets. During the past decade we have engaged in synthetic studies on macrocyclic pyrrolizidine alkaloids.²⁾ In connection with continuous synthetic studies, we wish to describe an efficient synthesis of racemic necine bases ((±)-turneforcidine (7), (±)-hastanecine (8), and (±)-platynecine (10)) from a common intermediate, ethyl (±)-1-hydroxy-2,3,5,6-tetrahydro-1*H*-pyrrolizine-7-carboxylate (2).

Results and Discussion

Previously, we reported on the syntheses of (±)-retronecine (3)³⁾ and (±)-otonecine (4)⁴⁾ from a common intermediate, hydroxy unsaturated ester 2 prepared by the γ -hydroxylation of the β -amino- α,β -unsaturated ester system




Scheme 1.



Scheme 2.

in ethyl 2,3,5,6-tetrahydro-1*H*-pyrrolizine-7-carboxylate (1) (Scheme 1). We have found that hydroxy unsaturated ester 2 was isomerized smoothly into bicyclic keto ester 5 in 54% yield upon an AcOH treatment (Scheme 2). This isomerization of 2 into 5 may proceed through intermediates **A** and **B**, as shown in Scheme 3. Thus, the initial protonation of 2 provides iminium salt **A**, which in turn is deprotonated to give enol **B**. Tautomerization of **B** gives 5. The stereochemistry of 5 was tentatively assigned, which was confirmed by successive transformations of 5 into (±)-turneforcidine (7). Thus, the hydrogenation of keto ester 5 over PtO₂ gave hydroxy ester 6 (40%), from which (±)-turneforcidine (7) was synthe-



Scheme 3.

sized by LiAlH_4 reduction in 86% yield. On the other hand, the direct reduction of keto ester **5** with LiAlH_4 gave ($\pm$)-hastanecine (**8**) along with the ($\pm$)-turneforcidine (**7**) in 40% and 26% yield, respectively. Finally, ($\pm$)-platynecine (**10**) was synthesized by the LiAlH_4 reduction of lactone **9**, which was previously employed in our synthesis of ($\pm$)-retronecine (**3**).³⁾

In the present and previous synthesis we achieved efficient syntheses of necine bases, turneforcidine (**7**), hastanecine (**8**), platynecine (**10**), retronecine (**3**), and otonecine (**4**) from the common intermediate hydroxy unsaturated ester **2** in racemic form.

Experimental

The melting points are uncorrected. IR spectra were taken on a JASCO IR-810 spectrophotometer. ^1H NMR spectra were recorded on a JEOL JNM EX-270 (270 MHz) spectrometer. The chemical shifts (δ) are reported in ppm downfield from internal tetramethylsilane, and the coupling constants in Hz. Low-resolution (EIMS) and high-resolution mass spectra (HREIMS) were measured on a JEOL JMS-LG2000 instrument. Fuji-Davison silica gel BW-820MH and Merck aluminum oxide 90 (activity II-III, Art. 1097) (alumina) were used for column chromatography. Merck precoated silica gel 60 F₂₅₄ plates, 0.25 mm thickness and Merck precoated alumina 150 F₂₅₄ (Type T) plates, 0.25 mm thickness were used for analytical thin-layer chromatography (TLC) and for preparative TLC. Dichloromethane (CH_2Cl_2) and diisopropylamine were distilled from calcium hydride (CaH_2) under nitrogen. Benzene and toluene were distilled from sodium (Na) under nitrogen. Tetrahydrofuran (THF) was distilled from sodiumbenzophenone ketyl. Methanol (MeOH) and ethanol (EtOH) were distilled from $\text{Mg}(\text{OMe})_2$ and $\text{Mg}(\text{OEt})_2$ under nitrogen, respectively. Chloroform (CHCl_3) was distilled from phosphorus pentaoxide.

Ethyl ($\pm$)-1-Hydroxy-2,3,5,6-tetrahydro-1H-pyrrolizine-7-carboxylate (2**).** (A) **Hydroxylation with ($\pm$)-3-Phenyl-2-phenylsulfonyloxaziridine.** To a cooled (-78°C), stirred solution of ethyl 2,3,5,6-tetrahydro-1H-pyrrolizine-7-carboxylate (**1**)⁵⁾ (29.2 mg, 0.161 mmol) in THF (1.5 ml) under nitrogen was added dropwise a 0.5 M (1 M = 1 mol dm^{-3}) toluene solution of potassium bis(trimethylsilyl)amide (1.13 ml, 0.565 mmol). After the mixture was stirred at -78°C for 1 h, a solution of ($\pm$)-3-phenyl-2-phenylsulfonyloxaziridine⁶⁾ (88.5 mg, 0.339 mmol) in THF (3.0 ml) was added to the cooled solution. The mixture was stirred at -78°C for 30 min, and the reaction was quenched by the addition of saturated $\text{Na}_2\text{S}_2\text{O}_3$ solution (1 ml). The mixture was diluted with H_2O (3 ml) and extracted with EtOAc (2 ml $\times$ 5). The combined extracts were washed, dried, and concentrated under reduced pressure. The residue was purified by repeated column chromatography [(1) silica gel 3 g, CHCl_3 containing 1% Et_3N ; (2) silica gel 2.0 g, benzene containing 1% Et_3N], affording **2** (19.7 mg, 62%) as a colorless oil: IR (CHCl_3) 3340, 1595 cm^{-1} ; ^1H NMR (270 MHz, CDCl_3) δ = 1.27 (3 H, t, J = 7.3 Hz), 2.31 (1 H, m), 2.96–3.10 (3 H, m), 3.20 (1 H, dt, J = 3.6, 8.7 Hz), 3.28–3.48 (2 H, m), 4.16 (2 H, q, J = 7.3 Hz), 4.92 (1 H, t, J = 7.9 Hz); EIMS m/z (rel intensity) 197 (M^+ ; 30), 179 (56), 169 (38), 152 (16), 135 (96), 96 (100). HREIMS. Found: m/z 197.1053. Calcd for $\text{C}_{10}\text{H}_{15}\text{NO}_3$: M, 197.1052.

(B) **Hydroxylation with $\text{MoO}_5\cdot\text{HMPA}\cdot\text{Py}$.** To a cooled (-78°C), stirred solution of **1**⁵⁾ (250.8 mg, 1.836 mmol) in THF (6.3 ml) under nitrogen was added dropwise a 0.215 M THF solution of lithium diisopropylamide (19 ml, 4.08 mmol). After the mixture was stirred at -78°C for 1 h, $\text{MoO}_5\cdot\text{HMPA}\cdot\text{Py}$ ⁷⁾ (1.2 g, 2.8 mmol)

was added to the cooled solution. The mixture was stirred at -78°C for 20 min, and the reaction was quenched by the addition of saturated Na_2SO_3 solution (4.4 ml). The mixture was diluted with H_2O (1.5 ml) and extracted with EtOAc (60 ml $\times$ 2). The combined extracts were washed, dried, and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel (5.2 g, CH_2Cl_2 –EtOAc (5 : 1) $\rightarrow$ EtOAc $\rightarrow$ MeOH), affording **2** (193.3 mg, 71%).

Ethyl ($1R^*$, $7aR^*$)-($\pm$)-7-Oxo-2,3,5,6,7,7a-hexahydro-1H-pyrrolizine-1-carboxylate ($\pm$)-(5**).** A mixture of **2** (47.6 mg, 0.426 mmol) and AcOH (1 ml) was stirred at room temperature for 2 d. The reaction mixture was concentrated in vacuo. The residue was dissolved in saturated NaHCO_3 (1 ml) and the mixture was extracted with EtOAc (4 ml $\times$ 3). The combined extracts were washed, dried, and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel [3 g, CHCl_3 –MeOH (50/1) $\rightarrow$ (20/1)], affording **5** (25.5 mg, 54%) as a colorless oil: IR (CHCl_3) 1750, 1725 cm^{-1} ; ^1H NMR (270 MHz, CDCl_3) δ = 1.28 (3 H, t, J = 7.3 Hz), 2.08–2.17 (2 H, m), 2.44 (2 H, t, J = 7.3 Hz), 2.71 (1 H, tt, J = 10.6, 7.3 Hz), 2.94–3.03 (2 H, m), 3.20 (1 H, ddd, J = 12.5, 6.9, 5.6 Hz), 3.40 (1 H, dt, J = 11.5, 7.3 Hz), 3.75 (1 H, d, J = 4.9 Hz), 4.18 (2 H, q, J = 7.3 Hz); EIMS m/z (rel intensity) 197 (M^+ ; 34), 169 (47), 152 (30), 96 (100), 68 (40). HREIMS. Found: m/z 197.1051. Calcd for $\text{C}_{10}\text{H}_{15}\text{NO}_3$: M, 197.1052.

Ethyl ($1R^*$, $7R^*$, $7aR^*$)-($\pm$)-7-Hydroxy-2,3,5,6,7,7a-hexahydro-1H-pyrrolizine-1-carboxylate ($\pm$)-(6**).** A mixture of **5** (31.1 mg, 0.158 mmol) and PtO_2 (22.1 mg) in EtOH (1.2 ml) was stirred at room temperature for 2.5 h under a hydrogen atmosphere and then filtrated through a pad of Celite. The combined filtrate and washings were concentrated under reduced pressure. The residue was purified by column chromatography on alumina [1 g, EtOAc–MeOH (3/1)], providing **6** (12.7 mg, 40%) as a colorless oil: IR (CHCl_3) 3420, 1720 cm^{-1} ; ^1H NMR (270 MHz, CDCl_3) δ = 1.27 (3 H, t, J = 7.3 Hz), 1.93–2.02 (1 H, m), 2.05–2.18 (1 H, m), 2.20–2.28 (3 H, m), 2.56–2.72 (2 H, m), 3.06–3.23 (2 H, m), 3.72 (1 H, dd, J = 5.6, 4.6 Hz), 3.40 (1 H, dt, J = 11.5, 7.3 Hz), 3.75 (1 H, d, J = 4.9 Hz), 4.18 (2 H, q, J = 7.3 Hz), 4.38 (1 H, m); EIMS m/z (rel intensity) 199 (M^+ ; 8), 181 (25), 167 (8), 155 (42), 82 (52), 58 (100). Found: m/z 199.1215. Calcd for $\text{C}_{10}\text{H}_{17}\text{NO}_3$: M, 199.1208.

($1R^*$, $7R^*$, $7aR^*$)-($\pm$)-7-Hydroxy-7-hydroxymethyl-2,3,5,6,7,7a-hexahydro-1H-pyrrolizine-1-methanol ($\pm$)-(7**) [($\pm$)-Turneforcidine].** To an ice-cooled, stirred solution of **6** (12.7 mg, 0.064 mmol) in THF (2 ml) was added a 1 M THF solution of LiAlH_4 (0.19 ml, 0.191 mmol). The mixture was stirred at 0°C for 3 h, and the reaction was quenched by the addition of 5% aqueous THF (0.5 ml). The mixture was stirred for a while and filtered through a pad of Celite. The combined filtrate and washings were concentrated under reduced pressure. The residue was purified by TLC on alumina [20 cm $\times$ 20 cm, 0.25 mm thickness, CH_2Cl_2 –MeOH–aq NH_3 (5/1/0.05)], providing ($\pm$)-**7** (8.6 mg, 86%) as a colorless oil: IR (CHCl_3) 3370 cm^{-1} ; ^1H NMR (270 MHz, CDCl_3) δ = 1.55–1.71 (2 H, m), 1.88–2.11 (2 H, m), 2.43–2.74 (3 H, m), 3.06 (1 H, m), 3.21 (1 H, m), 3.32 (1 H, dd, J = 8.2, 5.6 Hz), 3.85 (1 H, dd, J = 9.9, 4.6 Hz), 4.34 (1 H, dd, J = 10.0, 5.6 Hz); EIMS m/z (rel intensity) 157 (M^+ ; 27), 133 (10), 113 (39), 89 (14), 82 (100). Found: m/z 157.1090. Calcd for $\text{C}_8\text{H}_{15}\text{NO}_2$: M, 157.1110. Spectral properties of synthetic ($\pm$)-**7** were identical with those of authentic **7**⁸⁾ in all respects.

($1R^*$, $7S^*$, $7aR^*$)-($\pm$)-7-Hydroxy-2,3,5,6,7,7a-hexahydro-1H-pyrrolizine-1-methanol ($\pm$)-(8**) [($\pm$)-Hastanecine].** To an ice-cooled, stirred solution of **5** (27.9 mg, 0.141 mmol) in THF (4 ml)

was added a 1 M THF solution of LiAlH_4 (0.43 ml, 0.43 mmol). The mixture was stirred at 0 °C for 2 h, and the reaction was quenched by the addition of 5 % aqueous THF (1 ml). The mixture was stirred for a while and filtered through a pad of Celite. The combined filtrate and washings were concentrated under reduced pressure. The residue was purified by TLC on alumina [20 cm×20 cm, 0.25 mm thickness, CH_2Cl_2 -MeOH-aq NH_3 (3/1/0.05)], providing (±)-**8** (8.7 mg, 40%) as a colorless oil along with (±)-**7** (26%). (±)-**8**: IR (CHCl_3) 3350 cm^{-1} ; $^1\text{H NMR}$ (270 MHz, CDCl_3) $\delta = 1.63$ (1 H, m), 1.86–2.03 (3 H, m), 2.10 (1 H, m), 2.52 (1 H, dt, $J = 10.6$, 5.6 Hz), 2.64 (1 H, dt, $J = 10.6$, 6.6 Hz), 3.11–3.26 (3 H, m), 3.53 (1 H, dd, $J = 10.2$, 8.3 Hz), 3.81 (1 H, dd, $J = 10.2$, 4.6 Hz), 4.07 (1 H, dt, $J = 4.3$, 5.9 Hz); EIMS m/z (rel intensity) 157 (M^+ ; 17), 113 (25), 82 (100). Found: m/z 157.1111. Calcd for $\text{C}_8\text{H}_{15}\text{NO}_2$: M, 157.1110. The spectral properties of synthetic (±)-**8** was identical with those of authentic **8**⁹⁾ in all respects.

(1R*,7S*,7aS*)-(±)-**7-Hydroxy-2,3,5,6,7,7a-hexahydro-1H-pyrrolizine-1-methanol** (±)-(**10**) [(±)-Platynecine]. To a stirred solution of **9**^{3,10)} (20.0 mg, 0.130 mmol) in THF (2 ml) was added a 1 M THF solution of LiAlH_4 (0.4 ml, 0.4 mmol). The mixture was heated under reflux with stirring for 2 h. After the reaction mixture was cooled, the reaction was quenched by the addition of 5% aqueous THF (1 ml). The mixture was stirred for a while and filtered through a pad of Celite. The combined filtrate and washings were concentrated under reduced pressure. The residue was purified by TLC on alumina [20 cm×20 cm, 0.25 mm thickness, CH_2Cl_2 -MeOH-aq NH_3 (3/1/0.05)], providing (±)-**10** (12.0 mg, 60%) as colorless oil: IR (CHCl_3) 3370 cm^{-1} ; $^1\text{H NMR}$ (270 MHz, CDCl_3) $\delta = 1.60$ –1.70 (2 H, m), 1.90–2.01 (2 H, m), 2.40–2.70 (3 H, m), 3.00 (1 H, m), 3.20 (1 H, m), 3.30 (1 H, dd, $J = 8.2$, 3.0 Hz),

3.90–4.00 (2 H, m), 4.18 (1 H, ddd, $J = 4.2$, 4.0, 3.5 Hz); EIMS m/z (rel intensity) 157 (M^+ ; 15), 113 (25), 82 (100). Found: m/z 157.1095. Calcd for $\text{C}_8\text{H}_{15}\text{NO}_2$: M, 157.1110. Spectral properties of synthetic (±)-**10** were identical with those of authentic **10**⁹⁾ in all respects.

References

- 1) a) A. R. Mattocks, "Chemistry and Toxicology of Pyrrolizidine Alkaloids," Academic Press, London (1986), Chaps. 1, 3, 7, and 11; b) J. R. Liddell, *Nat. Prod. Rep.*, **13**, 187 (1996), and references cited therein.
- 2) H. Niwa, T. Sakata, and K. Yamada, *Bull. Chem. Soc. Jpn.*, **67**, 2345 (1994), and references cited therein.
- 3) H. Niwa, A. Kuroda, and K. Yamada, *Chem. Lett.*, **1983**, 125.
- 4) H. Niwa, Y. Uosaki, and K. Yamada, *Tetrahedron Lett.*, **24**, 5731 (1983).
- 5) N. J. Leonard and T. Sato, *J. Org. Chem.*, **49**, 1066 (1984).
- 6) F. A. Davis, L. C. Vishwarkarama, J. M. Billmers, and J. Finn, *J. Org. Chem.*, **49**, 3241 (1984).
- 7) E. Vedejs, D. A. Engler, and J. E. Teleschow, *J. Org. Chem.*, **43**, 188 (1978).
- 8) H. Niwa, H. Ishiwata, A. Kuroda, and K. Yamada, *Chem. Lett.*, **1983**, 789.
- 9) A. J. Aasen, C. C. J. Culvenor, and L. W. Smith, *J. Org. Chem.*, **34**, 4137 (1969).
- 10) M. Viscontini and H. Buzek, *Helv. Chim. Acta*, **55**, 670 (1972).