

Accounts

Reduction of CO₂ Directed toward Carbon–Carbon Bond Formation

Koji Tanaka

Institute for Molecular Science, Myodaiji, Okazaki 444

(Received June 12, 1997)

This paper describes electrochemical reduction of CO₂ directed toward carbon–carbon bond formation via metal–CO₂ adducts. An electrophilic attack of CO₂ to penta-coordinated low valent polypyridyl Ru complexes affords a Ru– η^1 -CO₂ adduct, which is easily converted to Ru–CO species either by an acid–base equilibrium in protic media and oxide transfer to CO₂ under aprotic conditions. Two-electron reduction of resultant Ru–CO in protic solutions competitively causes a cleavage of the Ru–CO bond (CO evolution) and formation of a thermally labile Ru–CHO bond. Besides further reduction of the latter to Ru–CH₂OH as precursors to CH₃OH and HOOCCH₂OH, Ru–CHO reacts with CO₂ to afford HCOOH with regenerating Ru–CO as the precursor to CO. Thus, the difficulty of multi-electron reduction of CO₂ in protic solutions is ascribed to the thermal lability and strong hydride donor character of Ru–CHO. On the other hand, two-electron reduction of Ru–CO in the presence of (CH₃)₄N⁺ or CH₃I under aprotic conditions produces thermally stable R–C(O)CH₃, which works as a precursor to CH₃C(O)CH₃. Two-electron reduction of M₃(μ_3 -S)₂ clusters (M = Co, Rh, Ir) causes an M–M bond cleavage and the nucleophilicity of the μ_3 -S ligand is also enhanced. As a result, two CO₂ molecules reductively activated probably on μ_3 -S and metal sites undergo the coupling reaction to give oxalate selectively.

Much attention has been paid to the utilization of CO₂ as a C1 source for chemicals and fuels to cope with an increase in the concentration in air and the oil shortage predicted for the near future. Carbon dioxide behaves as an electrophile under normal conditions. It is well known that CO₂ smoothly inserts into metal–alkyl and –aryl (M–R) bonds to form the corresponding M–OC(O)R complexes as precursors to carboxylic acids.¹⁾ Dienes and alkynes coordinated to metals are also likely to undergo an electrophilic attack of CO₂ to give lactones and pyrones.²⁾ In contrast to such electrophilic attack of CO₂ to organic substrates activated on metals, incorporation of CO₂ activated on metals to relatively inert organic substrates still remains unusual in organic syntheses. Carbon dioxide usually binds to low valent of metals with an η^1 - or η^2 -mode, where the electron density of the CO₂ group is greatly enhanced due to back electron transfer from the metal to CO₂. The η^1 -CO₂ mode is composed of electron transfer from the filled d_{z²} orbital of d⁸ metal centers to the CO₂ π^* orbital, and the η^2 -CO₂ mode involves electron transfer from a filled CO₂ π to metal in addition to that from a filled metal πd to the empty CO₂ π^* orbital. The first metal CO₂ complex (M–CO₂) was Ni(PCy₃)₂(η^2 -CO₂) prepared by Aresta et al. in 1975.³⁾ Since then, a variety of metal complexes with η^1 - η^2 -, μ^2 -, and μ^3 -CO₂ modes have been prepared.⁴⁾ Most M– η^1 -CO₂ complexes are thermally labile and are easily oxidized by air. A C–O bond of M– η^1 -CO₂ complexes is easily cleaved either by protonation in protic

media or by oxide transfer to free CO₂ in aprotic media, while such transformation from M– η^2 -CO₂ to M–CO has not been demonstrated so far. Accordingly, CO evolution in electro- and photochemical reduction of CO₂ catalyzed by metal complexes has been simply explained by M– η^1 -CO₂ intermediates, but little direct evidence has been obtained. One of the most crucial problems in electro- and photochemical reduction of CO₂ using homogeneous catalysts is why the products are limited to only CO and/or HCOOH.⁵⁾ Elucidation of the problem would greatly contribute to constructing a catalytic system that enables multi-electron reduction of CO₂ with forming new carbon–carbon bonds near the equilibrium potentials of the reactions.

A key question for the formation of M– η^1 -CO₂ complexes how to create coordinatively unsaturated low valent metal centers under mild conditions. On the other hand, adduct formation between CO₂ and organic bases is generally much easier than the formation of M– η^1 -CO₂ ones. The OCO angle of both adducts probably depends on the number of electrons transferred to the CO₂ moieties. The linear OCO of free CO₂ is hardly influenced at all by adduct formation with a weak base such as NH₃CO₂. On the other hand, the


OCO angle of the imidazolidone–CO₂ adduct (OOC–NC(O)–NHCH₂CH₂)[–] is calculated as 132°,⁶⁾ which is quite close to those of [Co(Pr–salen)K(η^1 -CO₂)] (132°) and [Rh(diars)–

$2(\eta^1\text{-CO}_2)]$ (126°).⁷⁾ Thus, CO_2 bonded to strong organic bases is also reductively activated similar to the process on low valent metals. A fundamental difference in the reductive activation of CO_2 on metals and organic bases is that the CO_2/CO conversion takes place on the former, but practically does not on the latter. Reductive activation of CO_2 on organic bases, therefore, may be utilized in CO_2 fixation without any accompanying C–O bond cleavage. It is worthy of note that acidity and basicity of neighboring groups (ligands) of the central metals are largely influenced by redox reactions of metal complexes even if the reactions are simply explained by the change of oxidation numbers of the central metals. Accordingly, basic ligands as well as metals are possible binding sites for the reductive activation of CO_2 , and the reactivity of the activated CO_2 on both sites are regulated by the redox reaction of the metal complexes.


This article focuses on activation of CO_2 on metals and ligands directed toward multi-electron reduction of CO_2 accompanied by carbon–carbon bond formation in the electrochemical reduction of CO_2 catalyzed by metal complexes.

Metal– $\eta^1\text{-CO}_2$ Complexes

The first $\text{M-}\eta^1\text{-CO}_2$ (M = transition metal) complex was prepared by the reaction of CO_2 with $[\text{Co}^{\text{I}}(\text{salen})\text{M}]^+$ ($\text{M}=\text{Li, Na, K, and Cs}$), where alkali metals used for the reduction of $[\text{Co}^{\text{II}}(\text{salen})]$ remains in the $\text{Co-}\eta^1\text{-CO}_2$ adducts and strongly interacts with oxygen of CO_2 binding to $\text{Co}(\text{I})$ (Eq. 1).^{7a)}




Since then, a series of the $\text{Co}^{\text{I}}\text{-}\eta^1\text{-CO}_2$ complexes with Schiff bases and macrocyclic ligands have been characterized.⁸⁾ The electron donor ability of the central metal atoms gives crucial influences on the stability of the $\text{M-}\eta^1\text{-CO}_2$ bond. In fact, $\text{Co}(\text{I})$ complexes with the redox potential of the $\text{Co}(\text{I})/\text{Co}(\text{II})$ couple more positive than -1.2 V essentially lose the ability to bind CO_2 at room temperature.⁹⁾ Similar to $\text{Co}^{\text{I}}\text{-}\eta^1\text{-CO}_2$ complexes, anionic $[\text{CpM}(\text{CO})_2(\eta^1\text{-CO}_2)]^-$ ($\text{M}=\text{Fe, Ru}$)¹⁰⁾ and $[\text{W}(\text{CO})_5(\eta^1\text{-CO}_2)]^{2-}$ ¹¹⁾ were prepared by the reaction of $[\text{CpM}(\text{CO})_2]^-$ and $[\text{W}(\text{CO})_5]^{2-}$ with CO_2 at low temperatures. Neutral Rh- and $\text{Ir-}\eta^1\text{-CO}_2$ complexes are prepared by an electrophilic attack of CO_2 to penta-coordinated $[\text{Ir}^{\text{I}}\text{Cl}(\text{dmpe})_2]^{12)}$ and $[\text{Rh}^{\text{I}}\text{Cl}(\text{diars})_2]^{7b)}$. Another synthetic route for $\text{M-}\eta^1\text{-CO}_2$ complexes is deprotonation of M-C(O)OH complexes derived from a nucleophilic attack of OH^- (or H_2O) to carbonyl carbon of electron deficient M-CO complexes (Eq. 2).




As $\text{M-}\eta^1\text{-CO}_2$ complexes are generally not stable in H_2O , only a series of $[\text{Ru}(\text{bpy})_2(\text{CO})(\eta^1\text{-CO}_2)]$, $[\text{Ru}(\text{bpy})_2(\text{CO})\{\text{C(O)OH}\}]^+$, and $[\text{Ru}(\text{bpy})_2(\text{CO})_2]^{2+}$ have been shown to exist as equilibrium mixtures in H_2O (see below).¹³⁾ A general tendency for the relative stability of M-C-

$(\text{O})\text{OH}$ and $\text{M-}\eta^1\text{-CO}_2$ complexes is not clear, because $[\text{FeCp}(\text{PPh}_3)(\text{CO})\{\text{C(O)OH}\}]$ loses CO_2 faster than $[\text{FeCp}(\text{PPh}_3)(\text{CO})(\eta^1\text{-CO}_2)]^-$,¹⁴⁾ but the reverse is true for $[\text{ReCp}(\text{NO})(\text{CO})\{\text{C(O)OH}\}]^{15)}$ and $[\text{IrCl}_2(\text{PMe}_2\text{Ph})_2(\text{CO})\{\text{C(O)OH}\}]^{16)}$. The basicity of an $\eta^1\text{-CO}_2$ group, that has a fundamental importance to understand the reactivity of the $\text{M-}\eta^1\text{-CO}_2$ complexes, can be reasonably evaluated from the conjugated acids, M-C(O)OH complexes. In contrast to pK_a values of organic carboxylic acids, the values of M-C(O)OH complexes widely range from ca. 2 to over 14 (Table 1) due to a large difference in the electron donor ability of the central metals to the CO_2 group. Despite such large differences in the basicity of $\text{M-}\eta^1\text{-CO}_2$ complexes, the knowledge concerning the structures and the basicity of $\eta^1\text{-CO}_2$ group is quite limited, since only three $\text{M-}\eta^1\text{-CO}_2$ complexes have had their molecular structures determined X-ray analysis so far. The electronic states of $\text{M-}\eta^1\text{-CO}_2$ complexes would be represented by resonance forms of I, II, and III (Eq. 3).



The amounts of electrons transferred to CO_2 increased in the order $\text{I} < \text{II} < \text{III}$, and the M-CO_2 bond will be strengthened in the same order. The alkaline metal (M') as a counter ion of $[\text{Co}^{\text{I}}(\text{salen})\text{M}'(\eta^1\text{-CO}_2)]^+$ (Eq. 1) which strongly interacts with oxygen of the CO_2 group must assist the electron flow from Co^{I} to CO_2 to stabilize the $\text{Co-}\eta^1\text{-CO}_2$ bond.

An addition of 2 equiv of OH^- to an $\text{H}_2\text{O}/\text{CH}_3\text{OH}$ solution of $[\text{RuL}_2(\text{CO})_2]^{2+}$ ($\text{L}=2,2'$ -bipyridine) crystallizes red $[\text{RuL}_2(\text{CO})(\eta^1\text{-CO}_2)] \cdot 3\text{H}_2\text{O}$; this complex is exceptionally stable among those metal– $\eta^1\text{-CO}_2$ complexes reported so far. The interconversion among $\text{cis-}[\text{RuL}_2(\text{CO})_2]^{2+}$, $\text{cis-}[\text{RuL}_2(\text{CO})\{\text{C(O)OH}\}]^+$ and $\text{cis-}[\text{RuL}_2(\text{CO})(\eta^1\text{-CO}_2)]$ in H_2O is very rapid (Eq. 4) and pK_a of $[\text{RuL}_2(\text{CO})\{\text{C(O)OH}\}]^+$ is determined as 9.6 at 20°C .¹³⁾



Molecular structures of the three complexes are determined by X-ray analysis.^{17,18)} The framework of the *trans*- N-Ru-CO , *trans*- N-Ru-C(O)OH , and *trans*- N-Ru-CO_2 moieties of $[\text{RuL}_2(\text{CO})_2](\text{PF}_6)_2$, $[\text{RuL}_2(\text{CO})(\text{C(O)OH})] \cdot (\text{H}_2\text{O})\text{CF}_3\text{COO}$, and $[\text{RuL}_2(\text{CO})(\eta^1\text{-CO}_2)] \cdot 3\text{H}_2\text{O}$, respectively, are depicted in Fig. 1. Red crystals of $[\text{RuL}_2(\text{CO})(\eta^1\text{-CO}_2)] \cdot 3\text{H}_2\text{O}$ contain three-dimensional hydrogen bonding

Table 1. pK_a Values of Hydroxycarbonyl Metal Complexes

Complex	pK_a	Ref.
$\text{ReCp}(\text{NO})(\text{CO})(\text{CO}_2\text{H})$	11	15
$[\text{RuL}_2(\text{CO})(\text{CO}_2\text{H})]^+$	9.6	13
$\text{Pt}(\text{C}_6\text{H}_5)(\text{PEt}_3)(\text{CO}_2\text{H})$	14	19
$[\text{CoL}(\text{CO}_2\text{H})]^{2+}$	3.1	8d
$[\text{Co}(\text{en})_2(\text{H}_2\text{O})(\text{CO}_2\text{H})]^{2+}$	2.5	8e




Fig. 1. *trans*-N-Ru-C frameworks of $[\text{RuL}_2\text{X}(\text{CO})]^{2+}$, $[\text{RuL}_2\text{X}(\text{C}(\text{O})\text{OH})]^+$, and $[\text{RuL}_2\text{X}(\text{CO}_2)]^0$ ($\text{L}=\text{bpy}$; $\text{X}=\text{CO}$).



networks among two oxygens of the $\eta^1\text{-CO}_2$ ligand and three hydrate molecules, which must assist electron flow to the CO_2 group from Ru and stabilizes the Ru-CO₂ bond. Nonequivalent two C-O bond distances (1.25 and 1.28 Å) of the $\eta^1\text{-CO}_2$ group, therefore, are caused by the differences in the number of hydrogen bondings between the two oxygen and hydrate water molecules. The Ru-CO₂ bond distance (2.06 Å) is close to that of $[\text{RhCl}(\text{diars})_2(\eta^1\text{-CO}_2)]$ (2.05 Å)^{7b)} but longer than of $[\text{Co}(\text{Pr-salen})(\text{K})(\eta^1\text{-CO}_2)]$ (1.99 Å).^{7a)} The average of the C-O bond distance in the $\eta^1\text{-CO}_2$ group (1.27 Å) is somewhat longer than the average of the C-O ones (1.23 Å) of $[\text{RhCl}(\text{diars})_2(\eta^1\text{-CO}_2)]$ and $[\text{Co}(\text{Pr-salen})(\text{K})(\eta^1\text{-CO}_2)]$. The OCO angle (121°) of the Ru-CO₂ group is narrower than those of Rh- $\eta^1\text{-CO}_2$ (126°) and Co- $\eta^1\text{-CO}_2$ (135°). The C-O bond distance and the OCO angle of CO_2^- are calculated as 1.24 Å and 135.2°, respectively, by ab initio MO calculations.¹⁹⁾ If an OCO angle of metal- $\eta^1\text{-CO}_2$ complexes is correlated with the amount of electrons transferred to the CO₂ ligands, σ -donation from Ru to CO₂ is stronger than that from Co and Rh to CO₂. Based on the $\text{p}K_a$ values of Ru-C(O)OH (9.6) and Co- $\eta^1\text{-C}(\text{O})\text{OH}$ complexes (2–3) (Table 1), and the OCO angles of $[\text{RuL}_2(\text{CO})(\eta^1\text{-CO}_2)]$ (121°), $[\text{Co}(\text{Pr-salen})(\text{K})(\eta^1\text{-CO}_2)]$ (135°),¹⁷⁾ and CO_2^- (135°), the electronic states of the Ru- and Co- $\eta^1\text{-CO}_2$ complexes are approximated by the resonance forms III and II, respectively. Accordingly, $[\text{RhCl}(\text{diars})_2(\eta^1\text{-CO}_2)]^{7b)}$ may be expressed by an intermediate between the resonance forms II and III on the basis of the OCO angle, though the $\text{p}K_a$ value of the conjugated acid is not clear.

The relatively high electron density on the CO₂ group of $[\text{RuL}_2(\text{CO})(\eta^1\text{-CO}_2)]$ is responsible for the smooth transformation from CO₂ to CO on the Ru atom in H₂O. The C=O and C-O bond distances (1.242 and 1.345 Å) and the OCO angle of 119° of $[\text{RuL}_2(\text{CO})(\text{C}(\text{O})\text{OH})]^+$ are close to the values of $[\text{Pt}(\text{C}_6\text{H}_5)(\text{PET}_3)_2\{\text{C}(\text{O})\text{OH}\}]$,¹⁹⁾ which exists as a hydrogen-bonded dimer structure similar to the solid state of acetic acid. Protonation of the CO₂ group of $[\text{RuL}_2(\text{CO})(\eta^1\text{-CO}_2)]$ causes shortening both the Ru- $\eta^1\text{-CO}_2$ and the Ru-N (*trans* to Ru-CO₂) bonds. Dehydroxylation of $[\text{RuL}_2(\text{CO})\{\text{C}(\text{O})\text{OH}\}]^+$ further shortens the Ru-C and Ru-N (*trans* to Ru-C(O)OH) bonds (Fig. 1). The Ru- $\eta^1\text{-CO}_2$ bond is primarily composed of σ -donation from Ru to CO₂, and the Ru-CO bond consists of π -back donation from Ru to CO and weak σ -donation from CO to Ru. The gradual elongation of the Ru-C bond distances from Ru-CO, Ru-C(O)OH to Ru-CO₂, therefore, strongly reflects the decrease in the π -bonding rather than the increase in the σ -bonding in the Ru-C bonds. It is worthy of note that an increase in the σ -donation from metals


to CO₂ strengthens M- $\eta^1\text{-CO}_2$ bonds, but does not always shorten the Ru-C bond distances. The agreement of the order of the Ru-CO < Ru-C(O)OH < Ru-CO₂ bond distances with *trans*-Ru-N ones strongly indicates that bpy as a σ -donor and π -acceptor ligand plays a role as an electron reservoir in the serious changes in the electronic structures of Ru in the equilibrium of Eq. 4. It should be noticed that $[\text{RuL}_2(\text{CO})(\eta^1\text{-CO}_2)]$ does not undergo a serious change at 100 °C in H₂O at pH 11 for 3 h, while $[\text{CpRu}(\text{CO})_2(\eta^1\text{-CO}_2)]\text{Na}$ decomposes to $[\text{CpRu}(\text{CO})_2(\text{H})]$ in THF at 0 °C.²⁰⁾ Thus, bpy ligands greatly contribute to the exceptional thermal stability of $[\text{RuL}_2(\text{CO})(\eta^1\text{-CO}_2)]$ as a metal- $\eta^1\text{-CO}_2$ complex.

Conversion from M-CO₂ to M-CO under Aprotic Conditions

Anionic $[\text{CpFe}(\text{CO})_2(\eta^1\text{-CO}_2)]^-$ and $[\text{W}(\text{CO})_5(\eta^1\text{-CO}_2)]^{2-}$ smoothly react with CO₂ to produce $[\text{CpFe}^{\text{I}}(\text{CO})_3]^+$ and $[\text{W}^0(\text{CO})_6]$, respectively, with forming CO_3^{2-} (Eqs. 5 and 6).^{20a,20b)}




Thus, $[\text{M}-\eta^1\text{-CO}_2]^{n+}$ with a very strong basic CO₂ group undergoes the oxide transfer reaction by free CO₂ to generate CO_3^{2-} and $[\text{M}-\text{CO}]^{(n+2)+}$, the latter of which works as the precursor to CO in electrochemical reduction of CO₂ (reductive disproportionation of CO₂) under aprotic conditions (Eq. 7).




But no oxide transfer reaction from CO_3^{2-} to M-CO complexes (reverse reaction of Eqs. 5 and 6) has been reported so far. Comparison of the basicity of M- $\eta^1\text{-CO}_2$ and CO_3^{2-} as oxide donors would greatly help us to understand the differences in the two bonds on the basis of $\text{p}K_a$ values of the conjugated acids: $[\text{Ru}(\text{bpy})_2(\text{CO})\{\text{C}(\text{O})\text{OH}\}]^+$ (9.6)⁷⁾ and HOCO_2^- (10.3).²²⁾ Red crystals of $[\text{RuL}_2(\text{CO})(\eta^1\text{-CO}_2)] \cdot 3\text{H}_2\text{O}$ are soluble only in protic media such as H₂O, CH₃OH, C₂H₅OH, but are insoluble in aprotic solvents. The characteristic solubility of the red crystals in both media is explained by the three dimensional hydrogen bonding networks of solvated water molecules in the solid state. On the other hand, the red crystals are smoothly solubilized into a CH₃CN solution containing LiCF₃SO₃ due to destruction of 3-dimensional hydrogen bonding networks of $[\text{RuL}(\text{CO})(\text{CO}_2)] \cdot 3\text{H}_2\text{O}$ by interaction of Li⁺ with oxygens of the $\eta^1\text{-CO}_2$ group.

The IR spectrum of a CD₃CN solution containing [RuL(CO)(CO₂)]·3H₂O and 3 equiv of LiCF₃SO₃ showed strong $\nu_{\text{asym}}(\text{CO}_2)$ and $\nu_{\text{sym}}(\text{CO}_2)$ bands at 1467 and 1246 cm⁻¹, which are close those of [RuL₂(CO)(η^1 -CO₂)]·3H₂O in KBr disks (1428 and 1242 cm⁻¹).¹⁷⁾ Thus, the molecular structure of [RuL₂(CO)(η^1 -CO₂)]·3H₂O in the solid state is maintained also in CH₃CN containing LiCF₃SO₃. Solubilization of [RuL₂(CO)(η^1 -CO₂)] in aprotic media enabled the direct interconversion between [RuL₂(CO)(η^1 -CO₂)] and [RuL₂(CO)₂]²⁺ without passing through [RuL₂(CO){C(O)OH}]⁺. Similarly, we found that [RuL₂(CO)(η^1 -CO₂)] is solubilized in CH₃CN in the presence of [K(crown)]⁺.


The reaction of [K(crown)]₂CO₃ with [RuL₂(CO)₂](PF₆)₂ afforded [RuL₂(CO)(η^1 -CO₂)] and CO₂ in CH₃CN (Eq. 8).²¹⁾



Thus, the reaction of Eq. 8 is the first example of the oxide transfer from CO₃²⁻ to M-CO, and the resultant [Ru(bpy)₂(CO)(η^1 -CO₂)] remained unchanged even after CO₂ bubbling into the solution (Eq. 9).




Moreover, the reaction of [RuL₂(¹²CO)(¹³CO)]²⁺ with [K(crown)]₂¹²CO₃ evolved only ¹²CO₂ in CH₃CN. On the other hand, an addition of LiCF₃SO₃ to the CO₂-saturated CH₃CN solution containing [RuL₂(CO)(η^1 -CO₂)] resulted in gradual precipitation of Li₂CO₃ with the generation of [RuL₂(CO)₂]²⁺ (Eq. 10).



Thus, removal of CO₃²⁻ from the solution as a precipitation forced the oxide transfer from [RuL₂(CO)(η^1 -CO₂)] to CO₂ to take place in CH₃CN. The interaction of Li⁺ with the η^1 -CO₂ would induce the electron transfer from Ru to CO₂, which results in an increase of nucleophilicity of the CO₂ ligand to lead to the formation of CO₃²⁻. It is worthy of note that the oxide transfer from [K(crown)]₂CO₃ to [RuL₂(CO)₂]²⁺ in CH₃CN (Eq. 8) finished in a few minutes, while it took an almost one day to complete the reverse reaction in the presence of LiCF₃SO₃ in CO₂-saturated CH₃CN (Eq. 10).

The rate of the oxide transfer reaction from [(CH₃)₄N]₂CO₃ to [RuL₂(CO)₂]²⁺ in DMSO is slow and the reaction intermediate was detected by the ¹³C NMR spectra. The ¹³C NMR spectra of a d₆-DMSO solution right after

mixing [RuL₂(¹²CO)(¹³CO)](PF₆)₂ and [(CH₃)₄N]₂CO₃ displayed two signals at δ =201.7 and 205.2 ppm. The chemical shifts of these signals are almost identical with those of [RuL₂(CO){C(O)OH}]⁺ in DMSO, suggesting the formation of a 1:1 adduct by an attack of CO₃²⁻ to a carbonyl carbon of [RuL₂(CO)₂]²⁺ (Eq. 11).




The fission of either the RuCO₂-CO₂ or RuC(O)-CO₃ bond of the 1:1 adduct determines the direction of the oxide transfer reactions of Eqs. 8, 9, and 10 (Scheme 1). The unusual oxide transfer reaction of Eq. 8 via the RuC(O)O-CO₂ bond fission (Scheme 1) is interpreted as the conversion from a weak base (CO₃²⁻) to a weaker one ([RuL₂(CO)(η^1 -CO₂)]). Although the oxide transfer reaction of Scheme 1 lies so far to [RuL₂(CO)(η^1 -CO₂)], the shift to [RuL₂(CO)₂]²⁺ in the equilibrium by an addition of Li⁺ indicates the small energy difference in the formation of the two complexes.


Electrochemical Reduction of CO₂ Catalyzed by Metal Complexes

A large number of metal complexes have proven to be active as catalyst precursors in electro- and photochemical reduction of CO₂.⁵⁾ Among those metal complexes, three metal complexes: [ReCl(L)(CO)₃], [Ni(cyclam)]²⁺ and [RuL₂(CO)₂]²⁺, have attracted much attention because of their characteristic reactivity and high efficiency in the reduction of CO₂.

Lehn et al. reported electrochemical reduction of CO₂ catalyzed by [ReCl(L)(CO)₃] in 1984.²³⁾ A series of [ReCl(L)-(L')(CO)₂] (L'=neutral ligand) have bifunctional ability as photosensitizers and as catalysts in the photochemical reduction of CO₂ to CO. Since they proposed a reaction mechanism for CO evolution in 1986 (Scheme 2), the arguments concerning the reaction mechanisms have still continued because no reasonable reaction intermediate has been characterized.²⁴⁾ The main arguments are i) whether Cl⁻ or CO initially dissociates from one-electron reduced form, ii) whether another one-electron reduction takes place or not before the resultant five-coordinate (or solvated) Re complex undergoes an electrophilic attack of CO₂. A recent IR spectroelectrochemical study suggests the occurrence of the CO₂ attack to both [ReL(CO)₃]⁻ (two electron reduced form) and [Re(dmbpy)(CO)₃][•] (one electron reduced form) (dmbpy=4,4'-dimethyl-2,2'-bipyridine) depending on their stability under electrolysis conditions.²⁵⁾





Scheme 1. The first reversible oxide transfer through a metal-C(O)-O-CO₂ adduct.


Scheme 2. Proposed mechanisms for the electrochemical reduction of CO₂ catalyzed by [Re(bpy)(CO)₃Cl].

Sauvage et al. found [Ni(cyclam)]²⁺ adsorbed on an Hg electrode evolves CO with an almost 100% current efficiency in the electrochemical reduction of CO₂ in H₂O.²⁹⁾ The reaction is explained by Scheme 3, though none of Ni-CO₂, -C(O)OH or -CO species have been identified. Sakaki et al. reported SCF ab initio calculations for [NiF(NH₃)₄]⁺ as the model of [Ni(cyclam)]⁺ adsorbed on Hg, and proposed a Ni^I-η¹-CO₂ adduct with the Ni-C bond distance of 1.92 Å and the OCO angle of 135.3° as the active species for the CO₂ reduction.³⁰⁾ The selectivity of CO/H₂ formation in the CO₂ reduction catalyzed by Ni(macrocyclic) complexes largely depends on the substituents of the ligands, and some of the derivatives are more reactive toward the selective CO₂ reduction than [Ni(cyclam)]²⁺.³¹⁾ However, little is known about why [Ni(cyclam)]²⁺ adsorbed on the surface of Hg selectively reduces CO₂ in H₂O via a possible intermediate of the Ni-η¹-CO₂ complex (Chart 1).


Tanaka et al. reported that electrolysis of [RuL₂(CO)₂]²⁺ at -1.20 V (vs. SCE) in DMF/H₂O (1 : 9 v/v, pH 6.0) under CO₂ atmosphere produced CO and H₂, while the same electrolysis conducted at pH 9.5 gave H₂, CO, and HCOO⁻ with a mole ratio of 1 : 1 : 1 in 1985.²⁶⁾ Furthermore, the electrochemical reduction of CO₂ by [RuL₂(CO)₂]²⁺ in dry CH₃CN in the presence of Me₃NHCl as a proton source predominantly produces HCOO⁻ with a current efficiency of 85%.²⁷⁾ The alternation of the main product from CO to HCOO⁻ with decreasing the proton concentrations in the reaction media is correlated with the equilibrium among [RuL₂(CO)₂]²⁺,

[RuL₂(CO){C(O)OH}]⁺ and [RuL₂(CO)(η¹-CO₂)]⁰ (Eq. 4) in the solution. Moreover, photochemical reduction of CO₂ catalyzed by the [RuL₂(CO)₂]²⁺/[RuL₃]²⁺/N(CH₂OH)₃ system in dry DMF selectively produces HCOO⁻ in a quantum yield (η=14%), while CO becomes the main product (η=14.8%) in the similar CO₂ reduction catalyzed by the [RuL₂(CO)₂]²⁺/[RuL₃]²⁺/BNAH system (BNAH=1-benzyl-1,4-dihydronicotinamide) in DMF/H₂O mixture.²⁷⁾ These facts indicate that the electro- and photochemical reductions of CO₂ catalyzed by [RuL₂(CO)₂]²⁺ proceed in similar mechanisms. Electrochemical reduction of CO₂ catalyzed by

Scheme 3. Proposed mechanism for the electrochemical reduction of CO₂ by [Ni(cyclam)]²⁺.Chart 1. RRSS-NiHTIM²⁺, NiDMC²⁺, and Nicyclam²⁺.

Scheme 4. Reduction of CO₂ catalyzed by [Ru(bpy)₂(CO)₂]²⁺.

[RuL₂(CO)₂]²⁺ is explained by Scheme 4; [RuL₂(CO)₂]²⁺ undergoes irreversible two-electron reduction at -1.0 V (vs. SCE) (vide infra) to evolve CO with generating penta-coordinated [RuL₂(CO)]⁰, which reacts with CO₂ to form [RuL₂(CO)(η^1 -CO₂)]. Protonation of [RuL₂(CO)(η^1 -CO₂)] gives [RuL₂(CO)₂]²⁺ through [RuL₂(CO){C(O)OH}]⁺, both of which function as the precursors to CO and HCOO⁻ formation. It is worthy of note that [RuL₂(CO)(η^1 -CO₂)] is stable in CH₃OH as expected from the pK_a value of 9.6, but the η^1 -CO₂ complex is completely protonated to give [RuL₂(CO){C(O)OH}]⁺ under CO₂ in the same solvent due to the exothermic formation of CH₃OCO₂⁻ (Eq. 12).




The complete protonation of [Ru(bpy)₂(CO)(η^1 -CO₂)] affording [Ru(bpy)₂(CO){C(O)OH}]⁺ in CH₃OH under CO₂ (Eq. 12) is, therefore, responsible for the predominant formation of HCOO⁻ in the photo- and electrochemical reduction of CO₂ in the presence of weak proton donors. Besides [RuL₂(CO){C(O)OH}]⁺ derived from the equilibrium of Eq. 4, [RuL₂(CO){OC(O)H}] resulting from CO₂ insertion into the Ru-H bond of [RuL₂(CO)H]⁰ is also suggested as a precursor to HCOO⁻ in the reduction of CO₂ catalyzed by [RuL₂(CO)₂]²⁺.²⁸⁾ Although an intramolecular rearrangement from [RuL₂(CO){OC(O)H}]⁺ to [RuL₂(CO){C(O)OH}]⁺ to explain CO evolution has been also proposed, the conversion between M-OC(O)H and M-C(O)OH has not been demonstrated so far.

Multi-Electron Reduction of CO₂ under Protic Conditions


Electro- and photochemical reduction of CO₂ catalyzed by metal complexes under protic conditions generates CO and/or HCOOH together with H₂. A competitive electrophilic attack of CO₂ and proton to low valent metal centers produces either M-CO₂ or M-H bonds. CO evolution is simply ascribed to a reductive cleavage of a M-CO bond resulting from the acid-base equilibrium of M-CO₂ (Eq. 2).

Besides M-C(O)OH complexes, M-OC(O)H complexes derived from insertion of CO₂ into M-H bonds work as precursors to HCOO⁻ generation. The most crucial problem in those CO₂ reductions is why multi-electron reduction of CO₂ with C-C bond formation does not take place in homogeneous reactions. It should be noticed that multi-electron reduction of CO₂ is energetically favored compared with two-electron reduction of CO₂. The standard redox potentials *E*⁰ (vs. SCE) for CO₂ reduction shift to positive potentials (pH 7.0, 20 °C) as the number of electron involved in the reduction increases.




Although general pathways for 4-, 6-, and 8-electron reduction of CO₂ have not been demonstrated so far, the reduction of M-CO rather than M-CO₂ is considered as the key process for multi-electron reduction of CO₂ by considering the smooth transformation between CO₂ and CO on metals. Some of the cationic metal carbonyl complexes are successively reduced to methyl derivatives through formyl-, hydroxymethyl complexes with hydride donors such as NaBH₄ and metal-hydrides (Scheme 5).³³⁾ As expected from chemical reduction of M-CO to M-CH₃ (Scheme 5), reduction of M-CO to M-CHO prior to M-CO bond cleavage (CO evolution) is likely to be a key step to make the multi-electron reduction of CO₂ in the homogeneous electrochemical reduction.


There are great differences in chemical and electrochemical reductions of M-CO complexes: electrochemical reduction of [RuL₂(CO)₂]²⁺ resulted in a Ru-CO bond cleavage in both protic and aprotic solutions, while a CO group of the complex is reduced to CH₃OH in a 10% yield by treatments with 4 equiv of NaBH₄ in CH₃CN/H₂O. Both [RuL₂(CO)(CHO)]⁺ and [RuL₂(CO)(CH₂OH)]⁺ were isolated as the reaction intermediates of CH₃OH in the same reaction conducted at -20 °C (Scheme 6).³⁴⁾ Similarly, treatments of [RuL(terpy)(CO)]²⁺ (terpy = 2,2':6',2''-terpyridine) with 4 equiv of BH₄⁻ quantitatively produced CH₃OH and [RuL(terpy)(CH₃CN)]⁺ in the same solvent, where [RuL(terpy)-



Scheme 5. Stepwise reduction of metal-CO complex.

Scheme 6. Reduction of [Ru(bpy)₂(CO)₂]²⁺ with NaBH₄.

(CHO)]⁺ was initially formed in the same reaction conducted at -40°C . Despite the formation of $[\text{RuL}_2(\text{CO})(\text{CHO})]^+$ and $[\text{RuL}_2(\text{CO})(\text{CH}_2\text{OH})]^+$ as the precursor to CH_3OH in the reaction of $[\text{RuL}_2(\text{CO})_2]^{2+}$ with BH_4^- (Scheme 6), electrochemical reduction of CO_2 catalyzed by the same complex produced only CO and HCOOH even at -20°C . On the other hand, the similar CO_2 reduction catalyzed by $[\text{RuL}(\text{terpy})(\text{CO})](\text{PF}_6)_2$ under the same electrolysis conditions produced not only CO and HCOO^- but also HC(O)H , CH_3OH , H(O)-CCOOH , and HOCH_2COOH (Eq. 13) (Fig. 2).



The difference in the catalytic activity of $[\text{RuL}(\text{terpy})(\text{CO})]^{2+}$ and $[\text{RuL}_2(\text{CO})_2]^{2+}$ toward the multi-electron reduction of CO_2 is associated with the stability of the reduced forms of these complexes at that temperature. Figure 3 shows the cyclic voltammogram (CV) of both complexes in CH_3CN under N_2 and CO_2 ; $[\text{RuL}_2(\text{CO})_2]^{2+}$ undergoes irreversible two-electron reduction at around -1.30 V (vs. Ag/Ag^+) at 20 and -20°C . The CV of $[\text{RuL}(\text{terpy})(\text{CO})]^{2+}$ displays a reversible redox couples at $E_{1/2} = -1.37$ and an irreversible cathodic wave at -1.69 V at 20°C , while the second redox reaction becomes a reversible electron transfer process at -20°C . On the basis of the fact that these redox reactions takes place in polypyridyl ligands, the redox behavior of the two complexes (Fig. 3) is explained as follows: $[\text{Ru}(\text{bpy})-(\text{bpy})(\text{CO})_2]^0$ readily dissociates CO , while $[\text{Ru}(\text{bpy})-(\text{terpy})(\text{CO})]^0$ is stable at -20°C . Moreover, the reaction




Fig. 2. Multi-electron reduction of CO_2 catalyzed by $[\text{Ru}(\text{bpy})(\text{terpy})(\text{CO})]^+$ in $\text{EtOH}/\text{H}_2\text{O}$ at -20°C .





Fig. 3. Cyclic voltammograms of $[\text{RuL}_2(\text{CO})_2](\text{PF}_6)_2$ (a and b) and $[\text{RuL}(\text{terpy})(\text{CO})](\text{PF}_6)_2$ (c and d) in CH_3CN at 20°C and -20°C under N_2 (solid lines) and CO_2 (dotted lines).

of $[\text{RuL}(\text{terpy})(\text{CO})]^0$ with H_2O at -30°C produced $[\text{RuL}(\text{terpy})(\text{CHO})]^+$. Accordingly, the pathway for multi-electron reduction of CO_2 by $[\text{RuL}(\text{terpy})(\text{CO})]^{2+}$ is explained by Scheme 7. Transformation from $[\text{RuL}(\text{terpy})(\text{CO}_2)]^0$ to $[\text{RuL}(\text{terpy})(\text{CO})]^{2+}$ is ascribed to the acid-base equilibrium of Eq. 2. Two-electron reduction of $[\text{RuL}(\text{terpy})(\text{CO})]^{2+}$ under protic conditions results in the formation of $[\text{RuL}(\text{terpy})(\text{CHO})]^+$ and CO evolution competitively. Protonation or carboxylation of $[\text{RuL}(\text{terpy})(\text{CHO})]^+$ under the electrolysis conditions gives HCHO and HOOCCHO , respectively. Further reduction and protonation of $[\text{RuL}(\text{terpy})(\text{CHO})]^+$ produces $[\text{RuL}(\text{terpy})(\text{CH}_2\text{OH})]^+$, which reacts with proton and CO_2 to generate CH_3OH and HOOCCH_2OH . It is, however, worthy of note that HCOO^- is still formed as the main product in the CO_2 reduction (Fig. 2), even though the concentration of $[\text{RuL}(\text{terpy})\{\text{C}(\text{O})\text{OH}\}]^+$ must be negligibly low because of the rapid conversion from $[\text{RuL}(\text{terpy})(\eta^1\text{-CO}_2)]$ to $[\text{RuL}(\text{terpy})(\text{CO})]^{2+}$ in $\text{H}_2\text{O}/\text{C}_2\text{H}_5\text{OH}$. A trace amount of CO evolution also reflects the smooth transformation from $[\text{RuL}(\text{terpy})(\text{CO}_2)]^0$ to $[\text{RuL}(\text{terpy})(\text{CHO})]^+$ prior to CO dissociation from $[\text{RuL}(\text{terpy})(\text{CO})]^{2+}$ under the electrolysis conditions. Moreover, $[\text{RuL}(\text{terpy})(\text{CHO})]^+$, that was obtained in a stoichiometric reaction of LiBEt_3H with $[\text{RuL}(\text{terpy})(\text{CO})](\text{PF}_6)_2$ in CD_3CN , smoothly reacts with CO_2 to produce $[\text{RuL}(\text{terpy})(\text{CO})]^{2+}$ and HCOO^- (60% yield) even at -20°C (Eq. 14).





Thus, $[\text{RuL}(\text{terpy})(\text{CHO})]^+$ is the branch intermediate for two- and multi-electron reduction of CO_2 . The existence of the feedback path from thermally labile $[\text{RuL}(\text{terpy})(\text{CHO})]^+$ to stable $[\text{RuL}(\text{terpy})(\text{CO})]^{2+}$ (as the precursor for CO evo-

Scheme 7. Multi-electron reduction of CO₂ catalyzed by [Ru(bpy)₂(trpy)(CO)]⁺.


lution) with generating HCOO⁻ under CO₂ (Eq. 14) reasonably explains why only CO and/or HCOOH are produced in electro- and photochemical reduction of CO₂ catalyzed by metal complexes reported so far.⁵⁾

Multi-Electron Reduction of CO₂ under Aprotic Conditions


Although [W(CO)₅(η¹-CO₂)]²⁻ and [CpFe(CO)₂(η¹-CO₂)]⁻ smoothly react with CO₂ to produce [W(CO)₆] and [CpFe(CO)₃]⁺,^{20a,20b)} these reactions are not suitable for reductive disproportionation reduction of CO₂ because the reduction of W(CO)₆ to [W(CO)₅]²⁻ takes place at potentials more negative than that of the direct reduction of CO₂. The basicity of [RuL₂(CO)(η¹-CO₂)] is not enough to undergo the oxide transfer reaction by CO₂ so that the equilibrium between [RuL₂(CO)₂]²⁺ and [RuL₂(CO)(η¹-CO₂)] lies so far to the latter under the normal conditions (Scheme 1). Replacement of a CO ligand of [RuL₂(CO)(η¹-CO₂)] by qu (qu=quinoline) greatly enhances the basicity of the η¹-CO₂ ligand, since electrochemically prepared [RuL₂(qu)-(CH₃CN)]⁰ rapidly reacts with CO₂ to give [RuL₂(qu)-(CO)]²⁺ and CO₃²⁻ through [RuL₂(qu)(η¹-CO₂)] (Eqs. 15 and 16).³⁵⁾



Indeed, the controlled potential electrolysis of [RuL₂(qu)-(CH₃CN)]²⁺ (or [RuL₂(qu)(CO)]²⁺) in the presence of LiBF₄ in CH₃CN under CO₂ at -1.40 V effectively catalyzes the reductive disproportionation of CO₂ to produce CO and CO₃²⁻ under the same electrolysis conditions (Eq. 7). On the other hand, the similar electrochemical reduction of CO₂ using [(CH₃)₄N]BF₄ in place of LiBF₄ in CH₃CN/DMSO at -1.50 V (vs. SCE) produced not only CO₃²⁻ and CO (η=42%) but also CH₃C(O)CH₃, HCOO⁻ and CH₃C(O)CH₂COO⁻ (η=16, 7, and 6%, respectively) (Eq. 17).




Unexpected formation of CH₃C(O)CH₃ in the electrochemical CO₂ reduction results from double methylation of [RuL₂(qu)(CO)]⁰ derived from [RuL₂(qu)(η¹-CO₂)]⁰, and both HCOO⁻ and CH₃C(O)CH₂COO⁻ are the products in the electrochemical carboxylation of CH₃C(O)CH₃ catalyzed by [RuL₂(qu)(η¹-CO₂)] (Eq. 18).




The rate of the formation of CH₃C(O)CH₃ in the electrochemical reduction of CO₂ (Eq. 17) is remarkably accelerated in the presence of CH₃I, and only CH₃C(O)CH₃, HCOO⁻, and CH₃C(O)CH₂COO⁻ were produced without evolving CO (Eq. 19).



The complete depression of CO evolution in these CO₂ reductions results from smooth formation of [RuL₂(qu){C(O)CH₃}]⁺ in the reaction of [RuL₂(qu)(CO)]⁰ with CH₃I prior to the Ru-CO bond breaking (Eqs. 20 and 21).



The mechanism for the formation of CH₃C(O)CH₃ in the electrochemical reduction of CO₂ catalyzed by [RuL₂(qu)-(CO)]²⁺ is represented in Scheme 8. An electrophilic attack of CO₂ to the two-electron reduced form of [RuL₂(qu)-(solvent)]²⁺ produces [RuL₂(qu)(η¹-CO₂)]⁰. Oxide transfer from [RuL₂(qu)(η¹-CO₂)]⁰ to CO₂ generates [RuL₂(qu)-(CO)]²⁺ and CO₃²⁻. Successive alkylation of [RuL₂(qu)-(CO)]⁰ by CH₃I (or Me₄N⁺) gives CH₃C(O)CH₃. Besides the oxide transfer from [RuL₂(qu)(CO₂)]⁰ to CO₂, the complex plays the precursor to HCOO⁻ in the presence of CH₃C(O)CH₃ as a proton source, and the resultant CH₃C(O)CH₂⁻ is trapped by CO₂ to form CH₃C(O)CH₂COO⁻. The qu ligand effectively blocks the attack of bulky CH₃I to Ru, since neither C₂H₆ nor CH₃COO⁻ was formed in the reduction of CO₂ by [RuL₂(qu)(CH₃CN)]²⁺. This was because the



Scheme 8. Catalytic formation of acetone in electrochemical reduction of CO_2 by $[RuL_2(qu)(solvent)]^{2+}$.

electrochemical reduction of CO_2 catalyzed by $[RuL_2(L')-(CH_3CH)]^{2+}$ (L' = iso-quinoline) in the presence of CH_3I produced C_2H_6 , which became the main product in the similar CO_2 reduction catalyzed by $[RuL(terpy)(CO)]^{2+}$ under the similar reaction conditions.

Activation of Ru-CO Derived from Ru- η^1 - CO_2

A nucleophilic attack of organic substrates to a CO group of cationic M-CO complexes is widely utilized for carbon-carbon bond formation in organic synthesis. On the other hand, an electrophilic attack of proton and CH_3I to a CO group of the two-electron reduced form of $[RuL_2(L')-(CO)]^{2+}$ derived from $[RuL_2(L')-(CO_2)]^0$ is the key reaction in the multi-electron reduction of CO_2 under electrolysis conditions. The efficiency of the multi-electron reduction of CO_2 accompanied by carbon-carbon bond formation, therefore, is dependent on reductive activation of not only M- CO_2 bonds but also M-CO ones. Spectroelectrochemical IR spectra greatly help to understand the reactivity of an M-CO bond depending on the oxidation states of the complexes. The $\nu(C\equiv O)$ band of $[RuL_2(qu)(CO)]^{2+}$ at 2015 cm^{-1} in CD_3CN undergoes the bathochromic shift to 1980 and 1939 cm^{-1} upon electrochemical one- and two-electron reductions of the complex at -1.21 and -1.50 V , respectively (Fig. 4 and Scheme 9). Thus, the ligand (bpy and qu) localized redox reaction of $[RuL_2(qu)(CO)]^{2+}$ causes the bathochromic shift of $\nu(CO)$ band by ca. 40 cm^{-1} per one electron. It is worthy of note that effective depression of CO dissociation from $[RuL_2(qu)(CO)]^0$, in contrast to $[RuL_2(CO)_2]^0$, is explained by the participation of π^* orbital of qu in the redox reaction. The $\nu(C\equiv O)$ band at 1980 cm^{-1} of $[RuL_2(qu)(CO)]^+$ is not affected by the presence of CH_3I , while two electron reduction of $[RuL_2(qu)(CO)]^{2+}$ in the presence of an equiv amount





Fig. 4. Time dependent IR spectra of $[RuL_2(qu)(CO)](PF_6)_2$ under the electrolysis at -1.21 V (a) and subsequently at -1.50 V in CD_3CN containing $LiBF_4$ under N_2 .


of CH_3I developed a new $\nu(C\equiv O)$ band of $[RuL_2(qu)(C(O)CH_3)]^+$ at 1568 cm^{-1} as the precursor for $CH_3C(O)CH_3$ in the electrochemical reduction of CO_2 . It is, however, not clear whether $[RuL_2(qu)(C(O)CH_3)]^+$ is formed by a direct attack of CH_3I to the CO group of $[RuL_2(qu)(CO)]^0$ or by CO insertion to a Ru- CH_3 intermediate such as $[RuL_2(qu)(CH_3)(CO)]^+$.

Intramolecular cyclization by taking advantage of a ligand localized redox reaction also greatly serves for the reductive activation of M-CO bond derived from M- CO_2 . The molecular structure of $[RuL_2(napy-\kappa N)(CO)]^{2+}$ ($napy$ = 1,8-naphthyridine) is close to that of $[RuL_2(qu)(CO)]^{2+}$. One-electron reduction of the former takes place on the napy ligand at -1.03 V (vs. $Ag/AgCl$). Introduction of one-electron into a π^* orbital of the monodentate napy ligand results in a remarkable enhancement of the nucleophilicity of the non-coordinated nitrogen atom. As a result, the $\nu(C\equiv O)$ band at 2003 cm^{-1} of $[RuL_2(napy-\kappa N)(CO)]^{2+}$ shifted to 1585 cm^{-1} upon one-electron reduction at -1.10 V in CD_3CN . Similar one-electron reduction of $[RuL_2(napy-\kappa N)(^{13}CO)]-(PF_6)_2$ also caused the red shift of the $\nu(^{13}C\equiv O)$ band from 1958 to 1543 cm^{-1} . The large red shift of the $\nu(CO)$ band by 418 cm^{-1} upon one-electron reduction is associated with the formation of the five-membered carbamoyl ring by an in-



Scheme 9. $\nu(CO)$ band depending on $[Ru(bpy)_2(qu)(CO)]^{n+}$ ($n=0, 1, 2$).

tramolecular nucleophilic attack of the non-bonded nitrogen of napy- π N to the carbonyl carbon (Eq. 22).³⁶⁾




The rate of the $\text{CH}_3\text{C}(\text{O})\text{CH}_3$ formation in the electrochemical reduction of CO_2 by using $(\text{CH}_3)_4\text{N}^+$ as a methylation agent is remarkably increased by participation of the five-membered carbamoyl ring (Eq. 22).


Activation of CO_2 on Non-Transition Metals

Monomeric $\text{M}-\eta^1\text{-CO}_2$ complexes may not be suitable species for CO_2 fixation to organic molecules if one considers a smooth C–O bond cleavage in protic and aprotic media. Such a C–O bond fission is practically inhibited in CO_2 bonded on organic bases. An electrophilic attack of CO_2 to O, S, and N atoms of $\text{M}-\text{XR}$ ($\text{X}=\text{O}, \text{S}$) and $\text{M}-\text{NR}_2$ groups is usually followed by CO_2 insertion into these polarized bonds. Accordingly, $\mu_3\text{-S}$ ligands in an $\text{M}_3(\mu_3\text{-S})$ moiety may provide feasible binding sites for the activation of CO_2 without any accompanying C–O bond cleavage, because CO_2 insertion into an $\text{M}-\text{S}$ bond of $\text{M}_3(\mu_3\text{-S})$ cores can probably be neglected due to the steric hindrance.

The $\text{p}K_a$ values of $\mu_3\text{-S}$ of $[\text{Fe}_4(\mu_3\text{-S})_4(\text{SC}_6\text{H}_4\text{C}_n\text{H}_{2n+1})_4]^{3-}$ and $[\text{Mo}_2\text{Fe}_6(\mu_3\text{-S})_8(\text{SEt})_3(\text{SC}_6\text{H}_4\text{C}_{2n+1})_6]^{5-}$ ($n=4\text{--}12$) were determined as ca. 9 and ca. 11, respectively, on the basis of the pH dependent redox reactions of $[\text{Fe}_4\text{X}_4(\text{YC}_6\text{H}_4\text{C}_n\text{H}_{2n+1})_4]^{2-}$ ($\text{X}, \text{Y}=\text{S}$ and Se) in aqueous micellar solutions.³⁷⁾ In accordance with the difference in the $\text{p}K_a$ values, strong interaction between $[\text{Mo}_2\text{Fe}_6\text{S}_8(\text{SEt})_9]^{5-}$ and CO_2 was confirmed in the cyclic voltammogram (CV) in dry CH_3CN , while no interaction was detected between $[\text{Fe}_4\text{S}_4(\text{SEt})_4]^{3-}$ and CO_2 in the same solvent. Furthermore, $[\text{Mo}_2\text{Fe}_6\text{S}_8(\text{SEt})_9]^{3-}$ catalyzes CO_2 fixation to $\text{RC}(\text{O})\text{SEt}$ ($\text{R}=\text{CH}_3, \text{C}_2\text{H}_5$, and C_6H_5) producing $\text{RC}(\text{O})\text{COO}^-$ without evolving CO under the electrolysis at -1.50 V vs. SCE in CH_3CN (Eq. 23).³⁸⁾



Catalytic formation of α -keto acids by CO_2 fixation to the carbonyl carbon of $\text{RC}(\text{O})\text{SR}'$ is essentially the same reaction as the CO_2 fixation by pyruvate synthase. Based on the reactivity depending on substituents of terminal RS^- ligands, the CO_2 fixation of Eq. 23 is explained in Scheme 10: i) an electrophilic attack of CO_2 on $\mu_3\text{-S}$ of $[\text{Mo}_2\text{Fe}_6(\mu_3\text{-S})_8(\text{SEt})_9]^{5-}$, ii) substitution of EtS^- ligated on Fe by $\text{RC}(\text{O})\text{SEt}$, iii) an addition of the reductively activated CO_2 on sulfur to the carbonyl carbon of $\text{RC}(\text{O})\text{SEt}$ ligated on Fe. The




Scheme 10. Proposed path for catalytic formation of α -keto acids.


process steps i and ii are supported by the observation that an addition of free EtS^- to the solution does not block the CO_2 adduct formation with $[\text{Mo}_2\text{Fe}_6\text{S}_8(\text{SEt})_9]^{5-}$, but strongly interferes with the α -keto acid formation due to depression of dissociation of EtS^- from $\text{Fe}-\text{SEt}$ group.

Coupling Reaction of Two CO_2 Molecules Activated on Metal Sulfur Clusters

Oxalate formation attracts much attention from the viewpoints of carbon-carbon bond formation by one- or two-electron reduction of CO_2 . Uncatalyzed electrochemical reduction of CO_2 on electrodes produces oxalate through a coupling reaction of CO_2^- (Eq. 24).



Savéant et al. have reported that anion radicals of aryl-esters and -nitrils effectively accelerate the coupling reaction of CO_2^- , though the redox potentials of those aryl compounds are quite negative and close to that of $E^\circ(\text{CO}_2/\text{CO}_2^-)$ at -2.21 V (vs. SCE).³⁹⁾ A new route for oxalate generation without passing through free CO_2^- is, therefore, desired thermodynamically. Two-electron reduction of trinuclear metal-sulfur clusters. $[(\text{MCp})_3(\mu_3\text{-S})_2]^{2+}$ ($\text{M}=\text{Co}, \text{Rh}, \text{Ir}$) causes an $\text{M}-\text{M}$ bond cleavage in the $\text{M}_3(\mu_3\text{-S})$ core. In addition, the basicity of the $\mu_3\text{-S}$ ligand of $[(\text{MCp})_3(\mu_3\text{-S})_2]^{2+}$ must be increased upon two-electron reduction. Two-electron reduction of $[(\text{CpM})_3(\mu_3\text{-S})_2]^{2+}$ may, therefore, create four possible binding sites for an electrophilic attack of CO_2 . The electrolysis of $[(\text{CoCp})_3(\mu_3\text{-S})_2]^{2+}$ in CO_2 -saturated dry CH_3CN at -0.70 V selectively produced $\text{C}_2\text{O}_4^{2-}$ as a white precipitate (Eq. 25).⁴¹⁾



The oxalate formation by the reduction of CO_2 under the electrolysis at -0.7 V is particularly noteworthy since the

standard redox potential of $\text{H}_2\text{C}_2\text{O}_4$ is -0.475 V (vs. NHE) in H_2O even at pH 0 (25°C). Similarly, oxalate was selectively produced in the electrochemical reduction of CO_2 catalyzed by $[(\text{RhCp}^*)_3(\mu_3\text{-S})_2]^{2+}$ at -1.50 V (vs. SCE) in CH_3CN under CO_2 atmosphere.⁴²⁾ In accordance with this, the two electron reduced forms of $[(\text{CoCp})_3(\mu_3\text{-S})_2]^{2+}$ and $[(\text{RhCp}^*)_3(\mu_3\text{-S})_2]^{2+}$ reacted with CO_2 to produce oxalate in 80 and 60% yields, respectively, with regenerating the oxidized clusters in CH_3CN . In contrast to the reaction of $[(\text{RhCp}^*)_3(\mu_3\text{-S})_2]^0$ with CO_2 in CH_3CN , $[(\text{IrCp}^*)_3(\mu_3\text{-S})_2]^0$ reacted with the solvent molecule under CO_2 to give $[(\text{IrCp}^*)_2(\text{Ir}(\eta^4\text{-Cp}^*\text{CH}_2\text{CN})(\mu_3\text{-S})_2)]^+$ with a linear CH_2CN moiety bonded to a Cp^* ring (Fig. 5) and oxalate. The catalytic activity of $[(\text{IrCp}^*)_2(\text{Ir}(\eta^4\text{-Cp}^*\text{CH}_2\text{CN})(\mu_3\text{-S})_2)]^+$ toward oxalate generation in electrochemical reduction of CO_2 is much higher than $[(\text{CoCp})_3(\mu_3\text{-S})_2]^{2+}$ and $[(\text{RhCp}^*)_3(\mu_3\text{-S})_2]^{2+}$.

The IR spectra of $[(\text{RhCp}^*)_3(\mu_3\text{-S})_2]^{2+}$ did not show any interaction with CO_2 in CD_3CN , while the electrolysis of $[(\text{RhCp}^*)_3(\mu_3\text{-S})_2]^{2+}$ in CO_2 -saturated CD_3CN at -1.50 V resulted in appearance of new bands at 1682 and 1605 cm^{-1} together with the 1633 cm^{-1} band of oxalate in the solution IR spectra. The 1680 and 1605 cm^{-1} bands completely disappeared upon the reoxidation of the solution at 0.3 V, and the 1633 cm^{-1} band of oxalate remained unchanged after the reoxidation. The 1682 and 1605 cm^{-1} bands shifted to 1636 and 1561 cm^{-1} , respectively, when the same electrolysis was conducted under $^{13}\text{CO}_2$. On the other hand, the IR spectra of $[(\text{IrCp}^*)_2(\text{Ir}(\eta^4\text{-Cp}^*\text{CH}_2\text{CN})(\mu_3\text{-S})_2)]^+$ showed a strong peak at 1680 cm^{-1} in CO_2 -saturated CD_3CN . Moreover, another band emerged around 1600 cm^{-1} upon one-electron reduction of the cluster under the electrolysis at -1.50 V in CD_3CN under $^{12}\text{CO}_2$. Prolonged electrolysis of the same solution also resulted in the appearance of the 1633 cm^{-1} band assignable to oxalate. This peak intensified with time. Thus,




Fig. 5. Crystal structure of $[(\text{IrCp}^*)_2(\text{IrCp}^*\text{CH}_2\text{CN})(\mu_3\text{-S})_2]^+$ determined by X-ray analysis.




Fig. 6. Two possible structures of 1:2 adducts between $[(\text{Cp}^*\text{Rh})_3(\mu_3\text{-S})_2]^0$ and CO_2 as the precursor to $\text{C}_2\text{O}_4^{2-}$ generation.

the two $\nu(\text{CO}_2)$ bands at 1682 and 1605 cm^{-1} appeared at the same time in the IR spectra of $[(\text{RhCp}^*)_3(\mu_3\text{-S})_2]^0$ in CD_3CN under CO_2 , while the former was observed in $[(\text{IrCp}^*)_2(\text{Ir}(\eta^4\text{-Cp}^*\text{CH}_2\text{CN})(\mu_3\text{-S})_2)]^+$ and the latter emerged in the IR spectra of $[(\text{IrCp}^*)_2(\text{Ir}(\eta^4\text{-Cp}^*\text{CH}_2\text{CN})(\mu_3\text{-S})_2)]^0$ in CD_3CN under CO_2 . These results indicate that $[(\text{RhCp}^*)_3(\mu_3\text{-S})_2]^0$ forms a 1:2 adduct with CO_2 as a precursor to oxalate.

If CO_2 is linked to $[(\text{RhCp}^*)_3(\mu_3\text{-S})_2]^0$ with an η^1 -mode, the OCO angle (2α) of an adduct is expressed by Eq. 26,⁴³⁾

$$\left(\frac{\nu^i}{\nu}\right)^2 = \left(\frac{M_C}{M_C^i}\right) \left(\frac{M_C^i + 2M_O \sin^2 \alpha}{M_C + 2M_O \sin^2 \alpha}\right) \quad (26)$$

in which ν^i and ν represent the ν_{asym} ($^{13}\text{CO}_2$) and ν_{asym} ($^{12}\text{CO}_2$) bands (cm^{-1}), and M_C^i , M_C , and M_O are the mass number of ^{13}C , ^{12}C , and ^{16}O , respectively. On the basis of the ν_{asym} ($^{12}\text{CO}_2$) at 1682 and 1605 cm^{-1} , the OCO angles of those CO_2 molecules linked to $[(\text{RhCp}^*)_3(\mu_3\text{-S})_2]^0$ are estimated as 157 and 132° , respectively. The latter is quite close to the OCO angle of $[\text{Co}(\text{Pr-salen})(\text{CO}_2\text{Na})]^+$ (135°).

There are two possibilities for the binding sites for the attack of two CO_2 molecules to $[(\text{RhCp}^*)_3(\mu_3\text{-S})_2]^0$ (Fig. 6); one is two coordinatively unsaturated Rh atoms produced by a Rh–Rh bond fission by the two-electron reduction of the cluster and the other is Rh and $\mu_3\text{-S}$. Ligation of two CO_2 to two Rh must be sterically blocked by bulky two Cp^* ligands (Fig. 6A). On the other hand, there seems to be no serious steric hindrance for the attack of CO_2 to $\mu_3\text{-S}$. The appearance of a strong band at 1680 cm^{-1} in the IR spectra of $[(\text{IrCp}^*)_2(\text{IrCp}^*\text{CH}_2\text{CN})(\mu_3\text{-S})_2]^+$ in CO_2 -saturated CD_3CN also supports an electrophilic attack of CO_2 to $\mu_3\text{-S}$ of $[(\text{RhCp}^*)_3(\mu_3\text{-S})_2]^0$. Non-equivalence of two CO_2 molecules in the IR spectra, therefore, is explained by the attack of CO_2 molecules to Rh and S (Fig. 4B). The oxalate formation in the electrochemical reduction of CO_2 catalyzed by $[(\text{MCp})_3(\mu_3\text{-S})_2]^0$, therefore, is attributed to the coupling reaction of two CO_2 molecules bonded on the adjacent M and S atoms.

References

- 1) a) "Molecular Electrochemistry of Inorganic, Bioorganic and Organometallic Compounds," ed by C. Amatore, A. Jutand, and M. F. Nielsen, NATO ASI Ser. C (1993), 307; b) "Chemical Fixation of Carbon Dioxide," ed by M. M. Halmann, CRC Press, London (1993).
- 2) a) "Enzymatic and Model Carboxylation and Reduction Re-

- actions for Carbon Dioxide," ed by M. Arest and J. V. Schloss, NATO Ser. C (1993); b) S. Derien, J. C. Clinet, E. Dunach, and J. Perichon, *J. Org. Chem.*, **58**, 2578 (1993); c) D. Sylvie, E. Dunach, and J. Perichon, *J. Am. Chem. Soc.*, **113**, 8447 (1991).
- 3) M. Aresta, C. F. Nobile, V. G. Albano, E. Fomi, and M. Manassero, *J. Chem. Soc., Chem. Commun.*, **1975**, 636.
- 4) a) D. H. Gibson, *Chem. Rev.*, **96**, 2063 (1996); b) J. Constamagna, G. Ferraudi, J. Canales, and J. Vergas, *Coord. Chem. Rev.*, **148**, 221 (1996).
- 5) a) H. Nakajima, T. Mizukawa, H. Nagao, and K. Tanaka, *Chem. Lett.*, **1995**, 251; b) K.-M. Lam, K.-Y. Wong, S.-M. Yang, and C.-M. Che, *J. Chem. Soc., Dalton Trans.*, **1995**, 1103; c) J. Costamagna, J. Canales, J. Vargas, and G. Ferraudi, *Pure Appl. Chem.*, **67**, 1045 (1995); d) A. M. Herring, B. D. Steffey, A. Miedaner, S. A. Wander, and D. L. DuBois, *Inorg. Chem.*, **34**, 1100 (1995); f) B. D. Steffey, A. Miedaner, M. L. Maciejewski-Farmer, P. R. Bernatis, A. M. Herring, V. S. Allured, V. Carperos, and D. L. DuBois, *Organometallics*, **13**, 4844 (1994); g) A. Szymaszek and F. Pruchnik, *Rhodium Express*, **5**, 18 (1994); h) E. Fujita, J. Haff, R. Sanzenbacher, and H. Elias, *Inorg. Chem.*, **33**, 4627 (1994); i) K. Ogura, H. Sugihara, J. Yano, and M. Higasa, *J. Electrochem. Soc.*, **141**, 419 (1994); j) E. Kimura, S. Wada, M. Shionoya, and Y. Okazaki, *Inorg. Chem.*, **33**, 770 (1994); k) R. J. Haines, R. E. Wittrig, and C. P. Kubiak, *Inorg. Chem.*, **33**, 4723 (1994); l) T. Ogata, Y. Yamamoto, Y. Wada, K. Murakoshi, M. Kusaba, N. Nakashima, A. Ishida, S. Takamuku, and S. Yanagida, *J. Phys. Chem.*, **99**, 11916 (1995); m) O. Ishitani, M. W. George, T. Ibusuki, F. P. A. Johnson, K. Koike, K. Nozaki, C. Pac, J. J. Turner, and J. R. Westwell, *Inorg. Chem.*, **33**, 4712 (1994), and references therein.
- 6) Y. Musashi, T. Hamada, and S. Sakaki, *J. Am. Chem. Soc.*, **117**, 11320 (1995).
- 7) a) S. Gambarotta, F. Arena, C. Floriani, and P. F. Zanazzi, *J. Am. Chem. Soc.*, **104**, 5914 (1982); b) J. C. Calabrese, T. Herskovitz, and J. B. Kinney, *J. Am. Chem. Soc.*, **105**, 5914 (1983).
- 8) a) D. A. Gangi and R. R. Durand, Jr., *J. Chem. Soc., Chem. Commun.*, **1986**, 697; b) D. J. Pearce and D. Pletcher, *J. Electroanal. Chem.*, **197**, 317 (1986); c) M. H. Schmidt, G. M. Miskelly, and N. S. Lewis, *J. Am. Chem. Soc.*, **112**, 3420 (1990); d) C. Creutz, H. A. Schwarz, J. F. Wishart, E. Fujita, and N. Sutin, *J. Am. Chem. Soc.*, **111**, 1153 (1989); e) N. Katz, D. J. Szalda, M. H. Chou, C. Creutz, and N. Sutin, *J. Am. Chem. Soc.*, **110**, 6591 (1988).
- 9) C. Creutz, in "Electrochemical and Electrocatalytic Reactions of Carbon Dioxide," ed by B. P. Sullivan, K. Krist, and H. E. Guard, Elsevier, Amsterdam (1993), p. 55.
- 10) J. M. Maher and N. J. Cooper, *J. Am. Chem. Soc.*, **102**, 7606 (1980).
- 11) G. R. Lee and N. J. Cooper, *Organometallics*, **4**, 794 (1985).
- 12) T. Herskovitz, *J. Am. Chem. Soc.*, **99**, 2391 (1977).
- 13) K. Tanaka, M. Morimoto, and T. Tanaka, *Chem. Lett.*, **1983**, 901.
- 14) D. H. Gibson and T.-S. Ong, *J. Am. Chem. Soc.*, **109**, 7191 (1987).
- 15) a) J. R. Sweet and W. A. G. Graham, *Organometallics*, **1**, 982 (1982); b) C. P. Casey, M. A. Andrews, and J. E. Rinz, *J. Am. Chem. Soc.*, **101**, 741 (1979).
- 16) K. Bowman, A. J. Deeming, and G. P. Proud, *J. Chem. Soc., Dalton Trans.*, **1985**, 857.
- 17) H. Tanaka, B.-C. Tzeng, H. Nagao, S.-M. Peng, and K. Tanaka, *Inorg. Chem.*, **32**, 1508 (1993).
- 18) K. Toyohara, H. Nagao, T. Adachi, T. Yoshida, and K. Tanaka, *Chem. Lett.*, **1996**, 27.
- 19) C. Amatore and J.-M. Saveant, *J. Am. Chem. Soc.*, **103**, 5021 (1981).
- 20) J. R. Pinkes, C. J. Masi, R. Chiulli, B. D. Steffey, and A. R. Cutler, *Inorg. Chem.*, **36**, 70 (1997).
- 21) H. Nakajima, K. Tsuge, and K. Tanaka, *Chem. Lett.*, **1997**, 485.
- 22) D. M. Kern, *J. Chem. Educ.*, **37**, 14 (1960).
- 23) J. Hawecker, J.-M. Lehn, and R. Ziessel, *J. Chem. Soc., Chem. Commun.*, **1984**, 328.
- 24) a) i) B. P. Sullivan, C. M. Bolinger, D. Conrad, W. J. Vining, and T. J. Meyer, *J. Chem. Soc., Chem. Commun.*, **1985**, 1414; b) T. R. O'Toole, B. P. Sullivan, M. R. -M. Bruce, L. Margerum, R. W. Murray, and T. J. Meyer, *J. Electroanal. Chem.*, **217**, 259 (1989); c) P. Christensen, A. Hamnett, A. V. G. Muir, and J. A. Timney, *J. Chem. Soc., Dalton Trans.*, **1992**, 1455.
- 25) F. P. A. Johnson, M. W. George, F. Hartl, and J. J. Turner, *Organometallics*, **15**, 3374 (1996).
- 26) a) H. Ishida, K. Tanaka, and T. Yanaka, *Organometallics*, **6**, 181 (1987); b) H. Ishida, H. Tanaka, K. Tanaka, and T. Tanaka, *J. Chem. Soc., Chem. Commun.*, **1987**, 131.
- 27) H. Ishida, K. Tanaka, and T. Tanaka, *Inorg. Chem.*, **6**, 553 (1987).
- 28) J. R. Pugh, M. R. M. Bruce, B. P. Sullivan, and T. J. Meyer, *Inorg. Chem.*, **30**, 86 (1991).
- 29) M. Beley, J.-P. Collin, R. Ruppert, and J.-P. Sauvage, *J. Am. Chem. Soc.*, **106**, 7461 (1986).
- 30) a) S. Sakaki, *J. Am. Chem. Soc.*, **112**, 7813 (1990); b) **114**, 2055 (1992).
- 31) E. Fujita, J. Haff, R. Sanzenbacher, and H. Elias, *Inorg. Chem.*, **33**, 4627 (1994).
- 32) S. Chardon-Noblat, M.-N. Collomb-Dunand-Sauthier, A. Deronzier, R. Ziessel, and D. Zsoldos, *Inorg. Chem.*, **33**, 4410 (1994).
- 33) a) P. M. Treichel and R. L. Schubhin, *Inorg. Chem.*, **6**, 1328 (1967); b) C. Lapinte and D. Astruc, *J. Chem. Soc., Chem. Commun.*, **1983**, 430; c) C. Lapinte, D. Catheline, and D. Astruc, *Organometallics*, **7**, 1683 (1988).
- 34) H. Nagao, T. Mizukawa, and K. Tanaka, *Inorg. Chem.*, **33**, 3415 (1994).
- 35) H. Nakajima, Y. Kushi, H. Nagao, and K. Tanaka, *Organometallics*, **14**, 5093 (1995).
- 36) H. Nakajima and K. Tanaka, *Chem. Lett.*, **1995**, 891.
- 37) a) K. Tanaka, T. Tanaka, and I. Kawafune, *Inorg. Chem.*, **23**, 516 (1984); b) K. Tanaka, and M. Moriya, and T. Tanaka, *Inorg. Chem.*, **25**, 835 (1986); c) M. Nakamoto, K. Tanaka, and T. Tanaka, *Bull. Chem. Soc. Jpn.*, **61**, 4099 (1989).
- 38) N. Komeda, H. Nagao, T. Matsui, G. Adachi, and K. Tanaka, *J. Am. Chem. Soc.*, **114**, 3625 (1992).
- 39) A. Gennaro, A. A. Isse, J.-M. Savéant, M.-G. Severin, and E. Vianello, *J. Am. Chem. Soc.*, **118**, 7190 (1996).
- 40) a) C. R. Pulliam, J. B. Thoden, A. M. Stacy, B. Spencer, M. H. Englert, and L. F. Dahl, *J. Am. Chem. Soc.*, **113**, 7398 (1991); b) A. Venturrelli and T. B. Rauchfuss, *J. Am. Chem. Soc.*, **116**, 4824 (1994).
- 41) a) Y. Kushi, H. Nagao, T. Nishioka, K. Isobe, and K. Tanaka, *J. Chem. Soc., Chem. Commun.*, **1995**, 1223.
- 42) Y. Kushi, H. Nagao, T. Nishioka, K. Isobe, and K. Tanaka, *Chem. Lett.*, **1994**, 2175.
- 43) K. Nakamoto, "Infrared and Raman Spectra of Inorganic and Coordination Compounds," Wiley-Interscience Publication, New York (1986).



Koji Tanaka, Ph. D., Professor of Institute for Molecular Science: born in Osaka in 1946. He obtained his B. Sc. in Chemistry from Osaka University in 1969. After he finished his master's course in 1971, he was immediately promoted to an assistant professor in the Department of Chemistry, Faculty of Engineering, Osaka University. He received his Ph. D. for research in metal complexes with chalcogenide ligands with the late Professor Toshio Tanaka in 1975. From 1978 to 1979, he studied Fischer-Tropsch reactions with Prof. R. B. King in Georgia University. Since then, he has worked on redox reactions of bioinorganic compounds. In 1990 he was promoted to an associate professor of Osaka University, and then moved to the Institute for Molecular Science as a full professor in the same year. His current research is focused on transformation of stable small molecules driven by redox reactions of metal complexes.