

One-Step Synthesis of Naphthofurandione, Benzofurandione, and Phenalenofuranone Derivatives by the CAN-Mediated Cycloaddition

Kazuhiro Kobayashi,* Tomokazu Uneda, Koujiro Tanaka, Masako Mori, Hideo Tanaka, Osamu Morikawa, and Hisatoshi Konishi

Department of Materials Science, Faculty of Engineering, Tottori University, Koyama-minami, Tottori 680-0945

(Received February 19, 1998)


The 3+2-type cycloaddition reaction of 2-hydroxy-1,4-naphthoquinones with various alkenes or phenylacetylene was mediated by ammonium cerium(IV) nitrate (CAN) to give the corresponding naphtho[2,3-*b*]furan-4,9-dione and naphtho[1,2-*b*]furan-4,5-dione derivatives. The reaction of 2-hydroxy-1,4-benzoquinones with alkenes in the presence of CAN similarly proceeded to give benzofuran-4,7-dione and benzofuran-4,5-dione derivatives. 3-Hydroxy-1*H*-phenalen-1-one also underwent the CAN-mediated cycloaddition with alkenes or phenylacetylene to give the corresponding 7*H*-phenaleno[1,2-*b*]furan-7-one derivatives.

The synthesis of naphthofurandione and benzofurandione derivatives has recently attracted much attention, because a number of these compounds have been found in nature and some of them have proven to exhibit a wide range of biological activities.¹⁾ Although many strategies have been used for the construction of naphthofurandione and benzofurandione skeletons,^{2,3)} most of approaches employ several steps to secure the target skeletons from readily available precursors.²⁾ In previous papers we reported on the photoinduced regioselective 3+2-type cycloaddition of 2-hydroxy-1,4-naphthoquinones and 2-hydroxy-1,4-benzoquinones⁴⁾ with various alkenes (or alkynes) leading to a one-pot formation of naphtho[2,3-*b*]furan-4,9-diones^{2a)} and benzofuran-4,7-diones,^{2b)} respectively. In this paper we wish to report that a similar type of cycloaddition can be mediated by ammonium cerium(IV) nitrate (CAN)⁵⁾ to give naphtho[2,3-*b*]furan-4,9-dione and benzofuran-4,7-dione derivatives along with the corresponding naphtho[1,2-*b*]furan-4,5-dione and benzofuran-4,5-dione derivatives.⁶⁾ We also found that this CAN-mediated reaction using 3-hydroxy-1*H*-phenalen-1-one as a related system of 2-hydroxy-1,4-naphthoquinone afforded 7*H*-phenaleno[1,2-*b*]furan-7-one derivatives.

Synthesis of Naphthofurandione and Benzofurandione Derivatives

We began our investigation by first examining the reaction between 2-hydroxy-1,4-naphthoquinones (**1a**) and 2,3-dimethyl-2-butene (**2a**). An acetonitrile solution of **1a** and 10 molar amounts of **2a** was treated with 2 molar amounts of CAN at 0 °C. After the usual work up, purification of the crude product by preparative TLC on silica gel afforded two pure products. The major and more mobile product (a lemon yellow solid) was identified as 2,2,3,3-tetramethyl-2,3-dihydronaphtho[2,3-*b*]furan-4,9-dione (**3a**) (61%) by a direct comparison with an authentic sample previously prepared by us.^{1a)} The minor and less mobile one (an orange solid) was identified as 2,2,3,3-tetramethyl-2,3-dihydronaphtho[1,2-*b*]furan-

4,5-dione (**4a**) (10%) from its spectroscopic data, which characterize an orthoquinone structure.⁷⁾ The formation of each product was completely regioselective and no trace amounts of their corresponding regioisomers could be detected. Use of 2 molar amounts of CAN was essential for the satisfactory production of the products in the present reaction. Reactions between 2-hydroxy-1,4-naphthoquinones **1** and a range of alkenes **2** were then carried out under the same reaction conditions, and the corresponding naphthofurandiones **3** and **4** were obtained (Scheme 1). The results obtained are given in Table 1. The reactions with 2-methyl-2-butene (**2b**) and 2-ethyl-1-butene (**2c**) also gave the corresponding products in good yields (Entries 2 and 3, respectively). 1-Hexene (**2e**), on the other hand, gave the expected products only in a low yield (Entry 5). However, it is worth noting that no 3+2 cycloaddition products could be obtained by the photochemical reaction of **1a** with terminal alkenes.^{1a)} Styrenes were found to react more smoothly to give the expected adducts in higher yields and seem to favor the formation of the naphtho[1,2-*b*]furan-4,5-diones (Entries 6 and 7). The reaction of **1a** with ethyl vinyl ether (**2h**) was carried out in the presence of an equimolar amount of sodium hydrogencarbonate to afford the naphtho[2,3-*b*]furan-4,9-dione derivative **3h** as the sole



Scheme 1.

Table 1. Products from the CAN Mediated Reactions between the Naphthoquinones **1** and the Alkenes **2**

Entry	1	2	3 (Yield/%) ^{a)}	4 (Yield/%) ^{a)}
1	1a (R=R'=H)	2a (R ¹ =R ² =R ³ =R ⁴ =Me)	3a (61)	4a (10)
2	1a	2b (R ¹ =R ² =R ³ =Me, R ⁴ =H)	3b (70)	4b (12)
3	1a	2c (R ¹ =R ² =Et, R ³ =R ⁴ =H)	3c (72)	4c (17)
4	1a	2d [R ¹ =R ⁴ =H, R ² -R ³ =(CH ₂) ₃]	3d (35) ^{c)}	4d (16) ^{d)}
5	1a	2e (R ¹ = <i>n</i> -Bu, R ² =R ³ =R ⁴ =H)	3e (12)	4e (14)
6	1a	2f (R ¹ =Ph, R ² =Me, R ³ =R ⁴ =H)	3f (38)	4f (48)
7	1a	2g (R ¹ =Ph, R ² =R ³ =R ⁴ =H)	3g (39)	4g (33)
8 ^{b)}	1a	2h (R ¹ =OEt, R ² =R ³ =R ⁴ =H)	3h (34)	—
9	1a	2i (R ¹ =OAc, R ² =R ³ =R ⁴ =H)	3i (12)	4i (19)
10	1a	2j (R ¹ =CO ₂ Me, R ² =Me, R ³ =R ⁴ =H)	3j (12)	4j (3)
11	1b (R=OMe, R'=H)	2b	3k (47)	4k (24)
12	1b	2c	3l (48)	4l (20)
13	1c (R=H, R'=OMe)	2c	3m (58)	4m (19)
14	1c	2f	3n (39)	4n (44)

a) Yield of isolated product, purified by preparative TLC on SiO₂ (EtOAc-hexane). b) One molar amount of NaHCO₃ was used. c) *cis*-Fused (Ref. 1a). d) *cis*-Fused. The stereochemistry of this product was determined by a comparison of its ¹H NMR spectrum with that of **3d** (Ref. 1a).


isolated product (Entry 8). The corresponding naphtho[1,2-*b*]furan-4,5-dione derivative was not isolated, probably due to its instability under the reaction and/or work-up conditions. Vinyl acetate (**2i**) proved to be less effective; the yields of the corresponding adducts, **3i** and **4i**, were rather lower than those of the other products (Entry 9). Methyl methacrylate (**2j**) gave the corresponding products, **3j** and **4j** (Entry 10), although the yield was considerably low. Methyl acrylate and acrylonitrile, however, could not give any addition product under these conditions. When the reaction was carried out in MeOH instead of acetonitrile, the yield of the products was considerably decreased.

The foregoing results indicate that the pathway to the products of the present cycloaddition is analogous to that proposed previously for the dihydrofuran formation from 1,3-diketones and alkenes.⁵⁾ As shown in Scheme 2, the initial formation of a reactive radical intermediate **B** through **A** was followed by oxidation to give tautomeric carbenium ion intermediates, **C** and **D**, which were trapped intramolecularly with the hydroxy groups of these intermediates to give rise to naphthofurandiones **3** and **4**, respectively. The good to high yields of the products from 2-methyl-2-butene (**2b**), 2-ethyl-1-butene (**2c**), and styrenes (**2f** and **2g**), and the low yields of those from 1-hexene (**2e**) and methyl methacrylate (**2j**) give support to this pathway. It can be assumed that the low yield of the product


from vinyl acetate (**2i**) is due to loss of acetyl radical from the corresponding intermediate **B**.

The use of phenylacetylene instead of alkenes was equally effective in cycloaddition with the hydroxynaphthoquinones **1**. As shown in Scheme 3, the formation of products **6** and **7** was almost quantitative. The attempted cycloaddition of **1a** with other alkynes, such as diphenylacetylene or 1-hexyne, however, resulted in the formation of an intractable mixture in each case.





The reactions of 2-hydroxy-1,4-benzoquinones were also carried out in order to examine the scope of the present cycloaddition. When 2-hydroxy-5,6-dimethyl-1,4-benzoquinone (**8a**) was allowed to react with (1-methylvinyl)benzene (**2f**) in the presence of CAN under the above-mentioned conditions, benzofurandiones **9a** and **10** were obtained in good yield. 2-Hydroxy-5-phenyl-1,4-benzoquinone (**8b**) and **2f** gave the benzofuran-4,7-dione derivatives in moderate yield (Scheme 4). 2-Hydroxy-1,4-benzoquinone, which is too unstable to be isolated in a pure form,⁸⁾ has proven to be usable in the present reaction. When 1,2,4-benzenetriol (**11**)⁸⁾ was reacted with 2-methyl-2-butene (**2b**) or 2-ethyl-1-butene (**2c**) in the presence of 4 molar amounts of CAN at 0 °C, cycloaddition occurred via unstable 2-hydroxy-1,4-benzoquinone to afford the corresponding 2,3-dihydrobenzofuran-4,7-dione derivatives, **12a** or **12b** (Scheme 5).



Scheme 2.



Scheme 3.



Synthesis of Phenalenofuranone Derivatives

Subsequently, we examined the possibility of utilizing the present CAN-mediated cycloaddition in preparing 7*H*-phenaleno[1,2-*b*]furan-7-one derivatives. Some compounds having structures based on this phenalenofuranone skeleton have been found in nature.⁹⁾ The only report on the synthesis of this class of molecules recorded by Frost and Morrison can be found in the literature.¹⁰⁾ Among several synthetic routes described in their report, the only successful general approach, which relies on the acid-catalyzed cyclization of 3-(1-naphthoyl)-4,5-dihydrofuran-2(3*H*)-ones, suffers from a tedious preparation of the starting materials. Therefore, it is important to develop a new general method for constructing this skeleton from readily available starting materials.

The preparation of 8,9-dihydro-7*H*-phenaleno[1,2-*b*]furan-7-ones **15** from 3-hydroxy-1*H*-phenalen-1-one (**13**) and a range of alkenes is outlined in Scheme 6. The cycloaddition reactions were

Table 2. CAN-Mediated Synthesis of 8,9-Dihydro-7*H*-phenaleno[1,2-*b*]furan-7-ones **15**

Entry	14	15 (Yield/%) ^{a)}
1	14a (= 2b ; $R^1=R^2=R^3=Me$)	15a (68)
2	14b (= 2c ; $R^1=R^2=Et, R^3=H$)	15b (82)
3	14c [$R^1-R^2=(CH_2)_4, R^3=H$]	15c (55)
4	14d (= 2g ; $R^1=Ph, R^2=Me, R^3=H$)	15d (92)
5	14e ($R^1=4-MeC_6H_4, R^2=R^3=H$)	15e (77)
6	14f ($R^1=4-ClC_6H_4, R^2=R^3=H$)	15f (72)
7	14g ($R^1=R^3=Ph, R^2=H$)	15g (60) ^{b)}
8	14h (= 2i ; $R^1=OAc, R^2=R^3=H$)	15h (35)

a) Isolated yield by preparative TLC on silica gel or recrystallization. b) *trans*:*cis*=82:18 (separable by preparative TLC on SiO_2).

carried out under conditions similar to those described above for the preparation of naphtho or benzofurandiones. The reactions proceeded smoothly to afford **15** with complete regioselectivity. The results are summarized in Table 2. The yields of the products were generally high, while vinyl acetate (**14h** = **2i**) gave a rather poor result (Entry 8), as observed in the preparation of naphthofurandiones. The spiro derivative **15c** was obtained with methylenecyclopentane (**14c**) (Entry 3). The reaction with *trans*-stilbene (**14g**) gave a separable mixture of stereoisomers **15g** (*trans*:*cis*=82:18) (Entry 7). The stereochemistries of these products were determined on the basis of ¹H NMR spectral analyses. Thus, the *E* isomer exhibited absorption signals assignable to the 8- and 9-H at much higher fields than the *Z* isomer. The coupling constants between these protons were reasonable, taking the conformation of the fused dihydrofuran ring into consideration (see Experimental). The reaction of **13** with phenylacetylene (5) in the presence of CAN also proceeded to give the expected cycloaddition product, 9-phenyl-7*H*-phenaleno[1,2-*b*]furan-7-one (**16**), in a satisfactory yield, as shown in Scheme 7.

In conclusion, the present CAN-mediated 3+2-type cycloaddition reactions can provide a one-step general method for preparing naphthofurandiones, benzofurandiones, and phenalenofuranones. The present method may found some value in organic synthesis because of simple manipulations as well as the ready availability of the starting materials.

Experimental

General Methods. The melting points were measured on a Laboratory Devices MEL-TEMP II melting apparatus and are uncorrected. IR spectra were determined for KBr disks, unless stated otherwise, with a Perkin-Elmer 1600 Series FT IR spectrometer. The ¹H NMR spectra were determined in $CDCl_3$ with either a JEOL JNX-PMX 60 NMR spectrometer operating at 60 MHz or a JEOL JNM-GX270 FT NMR spectrometer operating at 270 MHz. The chemical shifts were referenced relative to tetramethylsilane as an internal standard. Mass spectra were recorded with a JEOL AUTOMASS 20 spectrometer (Center for Cooperative Research and Development, this University). TLC was carried out on Merck Kieselgel 60 PF₂₅₄.

Materials. 2-Hydroxy-6-methoxy-1,4-naphthoquinone (**1b**),¹¹⁾ 2-hydroxy-8-methoxy-1,4-naphthoquinone (**1c**),¹²⁾ 2-hydroxy-5,6-dimethyl-1,4-benzoquinone (**8a**),¹³⁾ 2-hydroxy-5-phenyl-1,4-benzoquinone (**8b**),¹⁴⁾ and 1,2,4-benzenetriol (**11**)⁸⁾ were prepared by the appropriate reported procedures. All other chemicals, including CAN (95%, Kanto Chemicals Co., Inc.), were commercially avail-

able and used without further purification, except that acetonitrile was dried over CaH₂ and distilled prior to use.

2,2,3,3-Tetramethyl-2,3-dihydronaphtho[2,3-*b*]furan-4,9-dione (3a) and 2,2,3,3-Tetramethyl-2,3-dihydronaphtho[1,2-*b*]furan-4,5-dione (4a). General Synthetic Procedure. 2-Hydroxy-1,4-naphthoquinone (**1a**) (0.17 g, 1.0 mmol) and 2,3-dimethyl-2-butene (**2a**) (0.84 g, 10 mmol) were combined in acetonitrile (40 ml) and dissolved under stirring at room temperature. This solution was cooled to 0 °C and CAN (1.1 g, 2.0 mmol) was added in several portions to the yellow solution, which immediately turned into orange. The reaction was quenched after 20 min with a saturated aqueous NH₄Cl solution. After evaporation of the solvent and excess alkene, the crude product was extracted with Et₂O three times. The combined organic layers were washed successively with aqueous NaHCO₃ and brine, dried over anhydrous MgSO₄, and concentrated in vacuo. The crude product was purified by preparative TLC (SiO₂, 2:1 hexane–EtOAc) to afford the title compounds **3a** as a lemon yellow solid (0.16 g, 61%) and **4a** as an orange solid (26 mg, 10%). **3a**: Mp 101–102 °C (hexane–Et₂O) (lit.^{1a}) mp 101–102 °C). **4a**: *R*_f 0.46 (1:2 EtOAc–hexane); an amorphous; IR (neat) 1701, 1640, and 1614 cm^{−1}; ¹H NMR (270 MHz) δ = 1.34 (6H, s), 1.46 (6H, s), 7.5–7.65 (3H, m), and 8.05 (1H, d, *J* = 7.6 Hz); MS *m/z* (%) 256 (M⁺; 84) and 241 (100). Found: C, 75.28; H, 6.01%. Calcd for C₁₆H₁₆O₃: C, 74.98; H, 6.29%.

The naphthofurandione (**3,4,6**, and **7**), benzofurandione (**9** and **10**), and phenalenofuranone derivatives (**15** and **16**) were prepared following the above-mentioned procedure, except for the reaction of **1a** with **2h**, yielding **3h**, which was carried out in the presence of NaHCO₃ (84 mg, 1.0 mmol).

2,2,3-Trimethyl-2,3-dihydronaphtho[2,3-*b*]furan-4,9-dione (3b): Mp 168–169 °C (hexane–Et₂O) (lit.^{1a}) mp 168–169 °C).

2,2,3-Trimethyl-2,3-dihydronaphtho[1,2-*b*]furan-4,5-dione (4b): *R*_f 0.40 (1:2 EtOAc–hexane); mp 122–124 °C (hexane–CH₂Cl₂); IR 1697, 1646, 1633, and 1611 cm^{−1}; ¹H NMR (60 MHz) δ = 1.28 (3H, d, *J* = 7.2 Hz), 1.43 (3H, s), 1.62 (3H, s), 3.22 (1H, q, *J* = 7.2 Hz), 7.45–7.7 (3H, m), and 7.95–8.15 (1H, m); MS *m/z* (%) 242 (M⁺; 63) and 227 (100). Found: C, 74.30; H, 5.56%. Calcd for C₁₅H₁₄O₃: C, 74.36; H, 5.82%.

2,2-Diethyl-2,3-dihydronaphtho[2,3-*b*]furan-4,9-dione (3c): *R*_f 0.69 (1:2 EtOAc–hexane); mp 114–116 °C (hexane–CH₂Cl₂); IR 1682 and 1622 cm^{−1}; ¹H NMR (60 MHz) δ = 0.98 (6H, t, *J* = 7.2 Hz), 1.86 (4H, q, *J* = 7.2 Hz), 2.96 (2H, s), 7.6–7.85 (2H, m), and 8.0–8.2 (2H, m); MS *m/z* (%) 256 (M⁺; 54), 241 (67), and 227 (100). Found: C, 74.83; H, 6.30%. Calcd for C₁₆H₁₆O₃: C, 74.98; H, 6.29%.

2,2-Diethyl-2,3-dihydronaphtho[1,2-*b*]furan-4,5-dione (4c): *R*_f 0.44 (1:2 EtOAc–hexane); mp 139–140 °C (hexane–CH₂Cl₂); IR 1698, 1645, and 1612 cm^{−1}; ¹H NMR (60 MHz) δ = 0.93 (6H, t, *J* = 7.0 Hz), 1.86 (4H, q, *J* = 7.0 Hz), 2.94 (2H, s), 7.55–7.75 (3H, m), and 8.0–8.25 (1H, m); MS *m/z* (%) 256 (M⁺; 76) and 159 (100). Found: C, 74.69; H, 6.35%. Calcd for C₁₆H₁₆O₃: C, 74.98; H, 6.29%.

cis-2,3,3a,10b-Tetrahydro-1H-cyclopenta[*b*]naphtho[2,3-*d*]furan-5,10-dione (3d): Mp 133–134 °C (hexane–Et₂O) (lit.^{1a}) mp 132–134 °C).

cis-7,8,9a-Tetrahydro-6bH-cyclopenta[*b*]naphtho[2,1-*d*]furan-5,6-dione (4d): An amorphous; *R*_f 0.29 (1:2 EtOAc–hexane); IR (neat) 1698, 1642, and 1613 cm^{−1}; ¹H NMR (270 MHz) δ = 1.5–1.9 (4H, m), 2.0–2.1 (1H, m), 2.2–2.3 (1H, m), 3.90 (1H, t, *J* = 8.0 Hz), 5.53 (1H, dd, *J* = 8.0 and 5.4 Hz), 7.55–7.65 (3H, m), and 8.07 (1H, d, *J* = 7.6 Hz); MS *m/z* (%) 240 (M⁺; 55) and 212 (100). Found: C, 74.83; H, 5.16%. Calcd for C₁₅H₁₂O₃:

C, 74.99; H, 5.03%.

2-Butyl-2,3-dihydronaphtho[2,3-*b*]furan-4,9-dione (3e): *R*_f 0.55 (1:3 EtOAc–hexane); mp 74–76 °C (hexane–CH₂Cl₂); IR 1682, 1650, and 1624 cm^{−1}; ¹H NMR (270 MHz) δ = 0.98 (3H, t, *J* = 7.3 Hz), 1.35–1.6 (4H, m), 1.7–1.95 (2H, m), 2.85 (1H, dd, *J* = 17.1 and 8.3 Hz), 3.29 (1H, dd, *J* = 17.1 and 10.2 Hz), 5.0–5.1 (1H, m), 7.6–7.75 (2H, m), and 8.05–8.1 (2H, m); MS *m/z* (%) 256 (M⁺; 47) and 213 (100). Found: C, 74.70; H, 6.58%. Calcd for C₁₆H₁₆O₃: C, 74.98; H, 6.29%.

2-Butyl-2,3-dihydronaphtho[1,2-*b*]furan-4,5-dione (4e): *R*_f 0.21 (1:3 EtOAc–hexane); mp 67–72 °C (hexane–CH₂Cl₂); IR 1698, 1648, and 1610 cm^{−1}; ¹H NMR (270 MHz) δ = 0.96 (3H, t, *J* = 7.3 Hz), 1.4–1.6 (4H, m), 1.7–1.9 (2H, m), 2.79 (1H, dd, *J* = 15.2 and 7.6 Hz), 3.23 (1H, dd, *J* = 15.2 and 9.8 Hz), 5.0–5.1 (1H, m), 7.5–7.7 (3H, m), and 8.07 (1H, d, *J* = 7.3 Hz); MS *m/z* (%) 256 (M⁺; 19), 172 (23), and 159 (100). Found: C, 75.26; H, 6.30%. Calcd for C₁₆H₁₆O₃: C, 74.98; H, 6.29%.

2-Methyl-2-phenyl-2,3-dihydronaphtho[2,3-*b*]furan-4,9-dione (3f): Mp 112–113.5 °C (hexane) (lit.⁷) mp 112–114 °C).

2-Methyl-2-phenyl-2,3-dihydronaphtho[1,2-*b*]furan-4,5-dione (4f): Mp 120–123 °C (hexane–Et₂O) (lit.⁷) mp 120–122 °C).

2-Phenyl-2,3-dihydronaphtho[2,3-*b*]furan-4,9-dione (3g): Mp 164–165 °C (hexane–Et₂O) (lit.^{1a}) mp 163–165 °C).

2-Phenyl-2,3-dihydronaphtho[1,2-*b*]furan-4,5-dione (4g): Mp 162–164 °C (hexane) (lit.⁷) mp 162–164 °C).

2-Ethoxy-2,3-dihydronaphtho[2,3-*b*]furan-4,9-dione (3h): Mp 118–119 °C (hexane–Et₂O) (lit.^{1a}) mp 119–120 °C).

2-Acetoxy-2,3-dihydronaphtho[2,3-*b*]furan-4,9-dione (3i): Mp 193–195 °C (hexane–Et₂O) (lit.^{1a}) mp 193–195 °C).

2-Acetoxy-2,3-dihydronaphtho[1,2-*b*]furan-4,5-dione (4i): *R*_f 0.38 (1:1 EtOAc–hexane); mp 182–185 °C (hexane–CH₂Cl₂); IR 1753, 1695, 1651, and 1616 cm^{−1}; ¹H NMR (270 MHz) δ = 2.16 (3H, s), 3.13 (1H, dd, *J* = 17.0 and 2.9 Hz), 3.41 (1H, dd, *J* = 17.0 and 7.6 Hz), 7.03 (1H, dd, *J* = 7.6 and 2.9 Hz), 7.55–7.7 (3H, m), and 8.11 (1H, dd, *J* = 7.9 and 1.5 Hz); MS *m/z* (%) 198 (M – AcOH; 76) and 170 (100). Found: C, 64.73; H, 3.97%. Calcd for C₁₄H₁₀O₅: C, 65.12; H, 3.90%.

Methyl 2-Methyl-4,9-dioxo-2,3,4,9-tetrahydronaphtho[2,3-*b*]furan-2-carboxylate (3j): Mp 167–169 °C (hexane–Et₂O–CH₂Cl₂) (lit.^{1a}) mp 167–169 °C).

Methyl 2-Methyl-4,5-dioxo-2,3,4,5-tetrahydronaphtho[1,2-*b*]furan-2-carboxylate (4j): *R*_f 0.21 (1:2 EtOAc–hexane); mp 167 °C (hexane–CH₂Cl₂); IR 1757, 1702, 1652, and 1625 cm^{−1}; ¹H NMR (270 MHz) δ = 1.82 (3H, s), 3.06 (1H, d, *J* = 16.0 Hz), 3.51 (1H, d, *J* = 16.0 Hz), 3.83 (3H, s), 7.55–7.75 (3H, m), and 8.10 (1H, dd, *J* = 7.6 and 1.5 Hz); MS *m/z* (%) 272 (M⁺; 13), 244 (22), 212 (90), and 184 (100). Found: C, 66.40; H, 4.17%. Calcd for C₁₅H₁₂O₅: C, 66.17; H, 4.44%.

6-Methoxy-2,2,3-trimethyl-2,3-dihydronaphtho[2,3-*b*]furan-4,9-dione (3k): Mp 133–134 °C (hexane–CH₂Cl₂) (lit.^{1a}) mp 133–134.5 °C).

8-Methoxy-2,2,3-trimethyl-2,3-dihydronaphtho[1,2-*b*]furan-4,5-dione (4k): *R*_f 0.29 (1:2 EtOAc–hexane); mp 178–179 °C (hexane–CH₂Cl₂); IR 1692, 1644, and 1618 cm^{−1}; ¹H NMR (60 MHz) δ = 1.28 (3H, d, *J* = 7.2 Hz), 1.51 (6H, s), 3.22 (1H, q, *J* = 7.2 Hz), 3.94 (3H, s), 7.00 (1H, dd, *J* = 8.4 and 2.2 Hz), 7.02 (1H, d, *J* = 2.2 Hz), and 8.03 (1H, d, *J* = 8.4 Hz); MS *m/z* (%) 272 (M⁺; 13), 244 (83), and 203 (100). Found: C, 70.35; H, 5.96%. Calcd for C₁₆H₁₆O₄: C, 70.58; H, 5.92%.

2,2-Diethyl-6-methoxy-2,3-dihydronaphtho[2,3-*b*]furan-4,9-

dione (3l): R_f 0.55 (1 : 2 EtOAc–hexane); mp 144–145 °C (hexane–CH₂Cl₂); IR 1672, 1644, and 1625 cm⁻¹; ¹H NMR (60 MHz) δ = 0.99 (6H, t, J = 7.2 Hz), 1.86 (4H, q, J = 7.2 Hz), 2.95 (2H, s), 3.97 (3H, s), 7.08 (1H, dd, J = 8.0 and 2.6 Hz), 7.53 (1H, d, J = 2.6 Hz), and 8.03 (1H, d, J = 8.0 Hz); MS m/z (%) 286 (M⁺; 44), 258 (45), and 189 (100). Found: C, 71.02; H, 6.39%. Calcd for C₁₇H₁₈O₄: C, 71.31; H, 6.34%.

2,2-Diethyl-8-methoxy-2,3-dihydronaphtho[1,2-*b*]furan-4,5-dione (4l): R_f 0.25 (1 : 2 EtOAc–hexane); mp 145–148 °C (hexane–CH₂Cl₂); IR 1684, 1649, and 1620 cm⁻¹; ¹H NMR (60 MHz) δ = 0.98 (6H, t, J = 7.2 Hz), 1.80 (4H, q, J = 7.2 Hz), 2.90 (2H, s), 3.95 (3H, s), 6.97 (1H, dd, J = 8.2 and 2.6 Hz), 7.11 (1H, d, J = 2.6 Hz), and 8.04 (1H, d, J = 8.2 Hz); MS m/z (%) 286 (M⁺; 86) and 257 (100). Found: C, 71.05; H, 6.04%. Calcd for C₁₇H₁₈O₄: C, 71.31; H, 6.34%.

2,2-Diethyl-8-methoxy-2,3-dihydronaphtho[2,3-*b*]furan-4,9-dione (3m): R_f 0.45 (1 : 2 THF–hexane); mp 89–90 °C (hexane–CH₂Cl₂); IR 1678 and 1643 cm⁻¹; ¹H NMR (270 MHz) δ = 0.97 (6H, t, J = 7.4 Hz), 1.7–1.9 (4H, m), 2.93 (2H, s), 3.99 (3H, s), 7.23 (1H, d, J = 8.5 Hz), 7.63 (1H, dd, J = 8.5 and 7.4 Hz), and 7.74 (1H, dd, J = 7.4 and 1.1 Hz); MS m/z (%) 286 (M⁺; 55), 257 (59), and 189 (100). Found: C, 71.31; H, 6.25%. Calcd for C₁₇H₁₈O₄: C, 71.31; H, 6.34%.

2,2-Diethyl-6-methoxy-2,3-dihydronaphtho[1,2-*b*]furan-4,5-dione (4m): R_f 0.15 (1 : 2 THF–hexane); mp 205–207 °C (hexane–CH₂Cl₂); IR 1683, 1646, and 1630 cm⁻¹; ¹H NMR (270 MHz) δ = 0.97 (6H, t, J = 7.4 Hz), 1.84 (4H, q, J = 7.4 Hz), 2.88 (2H, s), 3.98 (3H, s), 7.16 (1H, d, J = 8.4 Hz), 7.30 (1H, dd, J = 8.5 and 7.4 Hz), and 7.59 (1H, dd, J = 7.4 and 1.1 Hz); MS m/z (%) 286 (M⁺; 73) and 189 (100). Found: C, 71.25; H, 6.41%. Calcd for C₁₇H₁₈O₄: C, 71.31; H, 6.34%.

8-Methoxy-2-methyl-2-phenyl-2,3-dihydronaphtho[2,3-*b*]furan-4,9-dione (3n): R_f 0.58 (1 : 1 THF–hexane); mp 201–204 °C (hexane–CH₂Cl₂); IR 1673, 1644, and 1600 cm⁻¹; ¹H NMR (270 MHz) δ = 1.86 (3H, s), 3.35 (1H, d, J = 16.5 Hz), 3.40 (1H, d, J = 16.5 Hz), 4.01 (3H, s), 7.2–7.5 (6H, m), 7.64 (1H, dd, J = 8.4 and 7.4 Hz), and 7.73 (1H, dd, J = 7.4 and 1.1 Hz); MS m/z (%) 320 (M⁺; 100). Found: C, 74.72; H, 5.12%. Calcd for C₂₀H₁₆O₄: C, 74.99; H, 5.03%.

6-Methoxy-2-methyl-2-phenyl-2,3-dihydronaphtho[1,2-*b*]furan-4,5-dione (4n): R_f 0.33 (1 : 1 THF–hexane); mp 200–203 °C (hexane–CH₂Cl₂); IR 1682, 1651, and 1634 cm⁻¹; ¹H NMR (270 MHz) δ = 1.88 (3H, s), 3.29 (1H, d, J = 15.3 Hz), 3.34 (1H, d, J = 15.3 Hz), 4.00 (3H, s), 7.20 (1H, d, J = 8.4 Hz), 7.3–7.45 (6H, m), and 7.63 (1H, dd, J = 8.4 and 7.4 Hz); MS m/z (%) 320 (M⁺; 100). Found: C, 74.71; H, 5.16%. Calcd for C₂₀H₁₆O₄: C, 74.99; H, 5.03%.

2-Phenylnaphtho[2,3-*b*]furan-4,9-dione (6a): Mp 247–250 °C (hexane–CH₂Cl₂) (lit.¹⁵) mp 246.5–247.5 °C; lit.^{1a}) mp 245–249 °C).

2-Phenylnaphtho[1,2-*b*]furan-4,5-dione (7a): R_f 0.18 (1 : 5 EtOAc–hexane); mp 217–219 °C (hexane–CH₂Cl₂); IR 1693, 1669, and 1656 cm⁻¹; ¹H NMR (270 MHz) δ = 7.06 (1H, s), 7.25–7.5 (4H, m), 7.69 (1H, td, J = 7.9 and 1.5 Hz), 7.75–7.8 (3H, m), and 8.10 (1H, d, J = 6.9 Hz); MS m/z (%) 274 (M⁺; 90), 246 (99), and 189 (100). Found: C, 78.55; H, 3.42%. Calcd for C₁₈H₂₀O₃: C, 78.83; H, 3.67%.

8-Methoxy-2-phenylnaphtho[2,3-*b*]furan-4,9-dione (6b): R_f 0.40 (1 : 2 THF–hexane); mp 217–220 °C (hexane–CH₂Cl₂); IR 1660 cm⁻¹; ¹H NMR (270 MHz) δ = 4.04 (3H, s), 7.12 (1H, s), 7.34 (1H, d, J = 8.4 Hz), 7.4–7.5 (3H, m), 7.67 (1H, dd, J = 8.4 and 7.4 Hz), and 7.85–7.9 (3H, m); MS m/z (%) 304 (M⁺; 68) and 105

(100). Found: C, 74.98; H, 3.94%. Calcd for C₁₉H₁₂O₄: C, 74.99; H, 3.97%.

6-Methoxy-2-phenylnaphtho[1,2-*b*]furan-4,5-dione (7b): R_f 0.24 (1 : 2 THF–hexane); mp 187–191 °C (hexane–CH₂Cl₂); IR 1674 cm⁻¹; ¹H NMR (270 MHz) δ = 4.00 (3H, s), 7.03 (1H, s), 7.06 (1H, d, J = 7.4 Hz), 7.3–7.5 (4H, m), 7.62 (1H, dd, J = 8.4 and 7.4 Hz), and 7.77 (2H, d, J = 7.9 Hz); MS m/z (%) 304 (M⁺; 100). Found: C, 75.07; H, 3.96%. Calcd for C₁₉H₁₂O₄: C, 74.99; H, 3.97%.

2,5,6-Trimethyl-2-phenyl-2,3-dihydrobenzofuran-4,7-dione (9a): Mp 91–93 °C (Et₂O–hexane) (lit.^{1b}) mp 91–92 °C).

2,6,7-Trimethyl-2-phenyl-2,3-dihydrobenzofuran-4,5-dione (10): An oil; R_f 0.35 (1 : 2 EtOAc–hexane); IR (neat) 1679, 1651, and 1632 cm⁻¹; ¹H NMR (270 MHz) δ = 1.79 (3H, s), 1.90 (3H, s), 2.14 (3H, s), 3.13 (2H, s), and 7.1–7.4 (5H, m); MS m/z (%) 268 (M⁺; 100). Found: C, 75.84; H, 5.95%. Calcd for C₁₇H₁₆O₃: C, 76.10; H, 6.01%.

2-Methyl-2,5-diphenyl-2,3-dihydrobenzofuran-4,7-dione (9b): Mp 86–87 °C (Et₂O–hexane) (lit.^{1b}) 85–87 °C).

2,2,3-Trimethyl-2,3-dihydrobenzofuran-4,7-dione (12a). A solution of 1,2,4-benzenetriol (**11**) (0.13 g, 1.0 mmol) and 2-methyl-2-butene (**2b**) (0.70 g, 10 mmol) in acetonitrile (40 ml) was treated with CAN (2.8 g, 4 mmol) at 0 °C for 35 min and worked up in a similar manner as described in General Synthetic Procedure to give **12** (48 mg, 25%) as a yellow oil: R_f 0.42 (1 : 3 EtOAc–hexane); IR (neat) 1674, 1640, and 1635 cm⁻¹; ¹H NMR (270 MHz) δ = 1.23 (3H, d, J = 6.9 Hz), 1.43 (3H, s), 1.47 (3H, s), 3.15 (2H, q, J = 6.9 Hz), 6.54 (1H, d, J = 10.6 Hz), and 6.57 (1H, d, J = 10.6 Hz); MS m/z (%) 192 (M⁺; 98) and 177 (100). Found: C, 68.99; H, 6.01%. Calcd for C₁₁H₁₂O₃: C, 68.73; H, 6.29%.

2,2-Diethyl-2,3-dihydrobenzofuran-4,7-dione (12b): An oil; R_f 0.43 (1 : 3 EtOAc–hexane); IR (neat) 1678, 1645, and 1627 cm⁻¹; ¹H NMR (60 MHz) δ = 0.96 (6H, d, J = 6.8 Hz), 1.78 (4H, q, J = 6.8 Hz), 2.82 (2H, s), and 6.52 (2H, s); MS m/z (%) 206 (M⁺; 30), 191 (43), and 177 (100). Found: C, 69.50; H, 6.65%. Calcd for C₁₂H₁₄O₃: C, 69.89; H, 6.84%.

8,9,9-Trimethyl-8,9-dihydro-7H-phenaleno[1,2-*b*]furan-7-one (15a): R_f 0.33 (1 : 3 EtOAc–hexane); mp 125–127 °C (hexane) (lit.¹⁰) mp 130–131 °C).

9,9-Diethyl-8,9-dihydro-7H-phenaleno[1,2-*b*]furan-7-one (15b): R_f 0.45 (1 : 2 EtOAc–hexane); mp 95–98 °C (hexane–CH₂Cl₂); IR 1646 and 1627 cm⁻¹; ¹H NMR (270 MHz) δ = 0.90 (6H, t, J = 7.3 Hz), 1.77 (4H, q, J = 7.3 Hz), 2.92 (2H, s), 7.48 (1H, dd, J = 8.3 and 7.3 Hz), 7.59 (1H, t, J = 7.6 Hz), 7.9–8.2 (3H, m), and 8.51 (1H, d, J = 7.3 Hz); MS m/z (%) 278 (M⁺; 21) and 249 (100). Found: C, 81.80; H, 6.38%. Calcd for C₁₉H₁₈O₂: C, 81.99; H, 6.52%.

Spiro[cyclopentane-1,9'-(8'H)-7'H-phenaleno[1,2-*b*]furan]-7'-one (15c): R_f 0.20 (1 : 3 EtOAc–hexane); mp 155–158 °C (hexane–Et₂O); IR 1645 and 1626 cm⁻¹; ¹H NMR (270 MHz) δ = 1.6–2.0 (6H, m), 2.2–2.3 (2H, m), 3.21 (2H, s), 7.58 (1H, t, J = 7.4 Hz), 7.71 (1H, t, J = 7.4 Hz), 8.0–8.15 (3H, m), and 8.61 (1H, dd, J = 7.3 and 1.1 Hz); MS m/z (%) 276 (M⁺; 41) and 247 (100). Found: C, 82.64; H, 5.95%. Calcd for C₁₉H₁₆O₂: C, 82.58; H, 5.84%.

9-Methyl-9-phenyl-8,9-dihydro-7H-phenaleno[1,2-*b*]furan-7-one (15d): R_f 0.53 (1 : 2 THF–hexane); mp 97–100 °C (hexane–CH₂Cl₂); IR 1630 cm⁻¹; ¹H NMR (60 MHz) δ = 1.84 (3H, s), 3.39 (2H, s), 7.2–7.75 (7H, m), 7.85–8.2 (3H, m), and 8.49 (1H, d, J = 7.2 Hz); MS m/z (%) 312 (M⁺; 35) and 297 (100). Found: C, 84.78; H, 4.99%. Calcd for C₂₂H₁₆O₂: C, 84.59; H, 5.16%.

9-(4-Methylphenyl)-8,9-dihydro-7H-phenaleno[1,2-*b*]furan-7-one (15e): R_f 0.32 (1 : 3 EtOAc–hexane); mp 190–192 °C

(hexane-CH₂Cl₂); IR 1643 and 1625 cm⁻¹; ¹H NMR (270 MHz) δ = 2.37 (3H, s), 3.28 (1H, dd, J = 15.3 and 7.9 Hz), 3.68 (1H, dd, J = 15.3 and 10.6 Hz), 6.01 (1H, dd, J = 10.6 and 7.9 Hz), 7.22 (2H, d, J = 7.9 Hz), 7.34 (2H, d, J = 7.9 Hz), 7.60 (1H, dd, J = 7.9 and 7.4 Hz), 7.74 (1H, t, J = 7.9 Hz), 8.05–8.15 (3H, m), and 8.64 (1H, dd, J = 7.4 and 1.1 Hz); MS m/z (%) 312 (M⁺; 100). Found: C, 84.74; H, 5.10%. Calcd for C₂₂H₁₆O₂: C, 84.59; H, 5.16%.

9-(4-Chlorophenyl)-8,9-dihydro-7H-phenaleno[1,2-*b*]furan-7-one (15f): R_f 0.38 (1 : 3 EtOAc–hexane); mp 165–166 °C (hexane-CH₂Cl₂); IR 1644 and 1627 cm⁻¹; ¹H NMR (270 MHz) δ = 3.23 (1H, dd, J = 15.7 and 7.3 Hz), 3.70 (1H, dd, J = 15.7 and 10.5 Hz), 6.02 (1H, dd, J = 10.5 and 7.3 Hz), 7.38 (4H, s), 7.62 (1H, t, J = 7.8 Hz), 7.75 (1H, t, J = 7.8 Hz), 8.08 (2H, d, J = 7.8 Hz), 8.14 (1H, dd, J = 7.8 and 1.1 Hz), and 8.64 (1H, dd, J = 7.8 and 1.1 Hz); MS m/z (%) 332 (M⁺; 100). Found: C, 76.05; H, 4.14%. Calcd for C₂₁H₁₃ClO₂: C, 75.79; H, 3.94%.

trans-8,9-Diphenyl-8,9-dihydro-7H-phenaleno[1,2-*b*]furan-7-one (trans-15g): R_f 0.29 (1 : 3 EtOAc–hexane); mp 154–159 °C (hexane-CH₂Cl₂); IR 1630 cm⁻¹; ¹H NMR (270 MHz) δ = 4.72 (1H, d, J = 5.8 Hz), 5.83 (1H, d, J = 5.8 Hz), 7.25–7.45 (10H, m), 7.68 (1H, t, J = 7.9 Hz), 7.73 (1H, t, J = 7.9 Hz), 8.12 (1H, dd, J = 7.9 and 1.1 Hz), 8.15 (1H, dd, J = 7.9 and 1.1 Hz), 8.22 (1H, dd, J = 7.9 and 1.1 Hz), and 8.56 (1H, dd, J = 7.9 and 1.1 Hz); MS m/z (%) 374 (M⁺; 83) and 297 (100). Found: C, 86.42; H, 4.70%. Calcd for C₂₇H₁₈O₂: C, 86.60; H, 4.86%.

cis-8,9-Diphenyl-8,9-dihydro-7H-phenaleno[1,2-*b*]furan-7-one (cis-15g): R_f 0.27 (1 : 3 EtOAc–hexane); mp 108–110 °C (hexane-CH₂Cl₂); IR 1637 and 1628 cm⁻¹; ¹H NMR (270 MHz) δ = 5.00 (1H, d, J = 9.5 Hz), 6.31 (1H, d, J = 9.5 Hz), 6.85–7.3 (10H, m), 7.70 (1H, t, J = 7.9 Hz), 7.75 (1H, t, J = 7.9 Hz), 8.15 (1H, d, J = 7.9 Hz), 8.18 (1H, d, J = 7.9 Hz), 8.25 (1H, d, J = 7.9 Hz), and 8.59 (1H, d, J = 7.9 Hz); MS m/z (%) 374 (M⁺; 85) and 297 (100). Found: C, 86.70; H, 4.91%. Calcd for C₂₇H₁₈O₂: C, 86.60; H, 4.86%.

9-Acetoxy-8,9-dihydro-7H-phenaleno[1,2-*b*]furan-7-one (15h): R_f 0.42 (1 : 1 EtOAc–hexane); mp 153–156 °C (hexane-CH₂Cl₂); IR 1758 and 1634 cm⁻¹; ¹H NMR (270 MHz) δ = 2.14 (3H, s), 3.25 (1H, d, J = 16.9 Hz), 3.50 (1H, dd, J = 16.9 and 7.4 Hz), 7.07 (1H, d, J = 7.4 Hz), 7.6–7.8 (2H, m), 8.0–8.2 (3H, m), and 8.65 (1H, d, J = 7.3 Hz); MS m/z (%) 280 (M⁺; 4.4) and 210 (100). Found: C, 72.64; H, 4.48%. Calcd for C₁₇H₁₂O₄: C, 72.85; H, 4.32%.

9-Phenyl-7H-phenaleno[1,2-*b*]furan-7-one (16): R_f 0.41 (1 : 3 EtOAc–hexane); mp 186–188 °C (hexane-Et₂O); IR 1645 cm⁻¹; ¹H NMR (270 MHz) δ = 7.20 (1H, s), 7.34 (1H, t, J = 7.3 Hz), 7.44 (2H, t, J = 7.3 Hz), 7.59 (1H, dd, J = 8.0 and 7.3 Hz), 7.69 (1H, dd, J = 8.0 and 7.3 Hz), 7.79 (2H, d, J = 7.3 Hz), 7.93 (1H, d, J = 8.0 Hz), 8.09 (1H, d, J = 8.0 Hz), 8.14 (1H, d, J = 7.3 Hz), and 8.66 (1H, d, J = 7.3 Hz); MS m/z (%) 296 (M⁺; 100). Found: C, 84.83; H, 3.78%. Calcd for C₂₁H₁₂O₂: C, 85.12; H, 4.08%.

We wish to thank Mrs. Miyuki Tanmatsu of this department for determining the mass spectra.

References

- 1) R. H. Thomson, "Naturally Occurring Quinones IV, Recent Advances," Blackie Academic & Professional, London (1997), pp. 112–308; F. Diaz and J. D. Medina, *J. Nat. Prod.*, **59**, 423 (1996), and references cited therein. Y. Imanaka and E. Miyamoto, Japan Patent 286401 (1996); *Chem. Abstr.*, **126**, 96897s (1997); T. Shibata, T. Suzuki, A. Kinoshita, and H. Hayata, Japan Patent 15881 (1997); *Chem. Abstr.*, **126**, 231511d (1997).
- 2) For recent reports on the one-pot construction of naphthofurandione and benzofurandione skeletons: a) K. Kobayashi, H. Shimizu, A. Sasaki, and H. Sugimoto, *J. Org. Chem.*, **58**, 4614 (1993); b) K. Kobayashi, Y. Kanno, and H. Sugimoto, *J. Chem. Soc., Perkin Trans. I*, **1993**, 1449; c) T. Shu, D.-W. Chen, and M. Ochiai, *Tetrahedron Lett.*, **37**, 5539 (1996); d) K. Kobayashi, T. Uneda, M. Kawakita, O. Morikawa, and H. Konishi, *Tetrahedron Lett.*, **38**, 837 (1997).
- 3) J. M. Heerding and H. W. Moore, *J. Org. Chem.*, **56**, 4048 (1991); L. S. Liebeskind, K. L. Granberg, and J. Zhang, *J. Org. Chem.*, **57**, 4345 (1992); H. Takagaki, M. Sakai, S. Ogawa, K. Inasawa, K. Yamazaki, T. Ikegawa, and N. Ikegawa, Japan Patent, 139177 (1992); *Chem. Abstr.*, **117**, 191671m (1992); M. Aso, A. Ojida, G. Yang, and K. Kanematsu, *Heterocycles*, **35**, 33 (1993); K. Shishido, T. Takata, T. Omodani, and M. Shibuya, *Chem. Lett.*, **1993**, 557; J. Koyanagi, K. Yamamoto, K. Nakayama, and A. Tanaka, *J. Heterocycl. Chem.*, **32**, 1289 (1995); J. Koyanagi, K. Yamamoto, K. Nakayama, and A. Tanaka, *J. Heterocycl. Chem.*, **34**, 407 (1997); O. Cherkaoui, P. Nebios, and H. Flinn, *Chem. Pharm. Bull.*, **45**, 457 (1997); P. J. Perry, V. H. Pavlidis, and J. A. Hadfield, *Tetrahedron*, **53**, 3195 (1997), and references cited in Refs. 1a and 1b.
- 4) For recent reports on the utilization of these hydroxyquinones in the preparation of heterocycle-fused quinones: A. B. de Oliveira, D. T. Ferreira, and D. S. Raslan, *Tetrahedron Lett.*, **29**, 155 (1988); L. Kopanski, D. Karbach, G. Selbutschka, and W. Steglich, *Liebigs Ann. Chem.*, **1987**, 793; C. B. Saitz, J. A. Valderrama, and R. Tapia, *Synth. Commun.*, **20**, 3103 (1990).
- 5) For metal-mediated cycloaddition reactions of 1,3-dicarbonyls to alkenes forming 1,2-dihydrofuran derivatives, see: K. Ichikawa, S. Uemura, and T. Sugita, *Tetrahedron*, **22**, 407 (1966); E. I. Heiba and R. M. Dessau, *J. Org. Chem.*, **39**, 3456 (1974); M. G. Vinogradov, A. E. Kondorsky, and G. I. Nikishin, *Synthesis*, **1988**, 60; E. Baciocchi and R. Ruzziconi, *Synth. Commun.*, **18**, 1841 (1988), and references cited therein. For recent reports, see: J. M. Mellor and S. Mohammed, *Tetrahedron Lett.*, **32**, 7111 (1991); V. Nair and J. Matthew, *J. Chem. Soc., Perkin Trans. I*, **1995**, 187; **1995**, 1881; **1996**, 1487; Y. R. Lee and B. S. Kim, *Tetrahedron Lett.*, **38**, 2095 (1997); Y. R. Lee, N. S. Kim, and B. S. Kim, *Tetrahedron Lett.*, **38**, 5671 (1997). See also: J. Iqbal and M. Mukhopadhyay, *Synlett*, **1997**, 876. For other approaches to the formation of dihydrofuran rings, see: J. Yoshida, K. Sakaguchi, and S. Isoe, *Tetrahedron Lett.*, **27**, 6075 (1986); H. Hagiwara, K. Kato, T. Suzuki, and M. Ando, *Tetrahedron Lett.*, **38**, 2103 (1997).
- 6) Preliminary communication: K. Kobayashi, M. Mori, T. Uneda, O. Morikawa, and H. Konishi, *Chem. Lett.*, **1996**, 451.
- 7) K. Kobayashi, A. Sasaki, H. Takeuchi, and H. Sugimoto, *J. Chem. Soc., Perkin Trans. I*, **1992**, 115.
- 8) K. C. Kurien and P. A. Robins, *J. Chem. Soc. B*, **1970**, 855.
- 9) K. G. Neil and R. Raistrick, *Biochem. J.*, **65**, 166 (1957); N. Narasimhachari, K. S. Gopalkrishnan, R. H. Haskins, and L. C. Vining, *Can. J. Microbiol.*, **9**, 134 (1963); I. C. Paul and G. A. Sim, *J. Chem. Soc.*, **1965**, 1097; D. A. Frost and G. A. Morrison, *Tetrahedron Lett.*, **1970**, 963; N. Narasimhachari and L. C. Vining, *J. Antibiot.*, **25**, 155 (1973); J. S. Brooks and G. A. Morrison, *J. Chem. Soc., Perkin Trans. I*, **1972**, 421; D. A. Frost and G. A. Morrison, *Tetrahedron Lett.*, **1972**, 4729, and references cited therein.
- 10) D. A. Frost and G. A. Morrison, *J. Chem. Soc., Perkin Trans. I*, **1973**, 2159.
- 11) A. C. Baillie and R. H. Thomson, *J. Chem. Soc. C*, **1966**,

2184.

12) S. Collins, Y. Hong, G. J. Hoover, and H. Rapoport, *J. Org. Chem.*, **44**, 2153 (1979).

13) L. F. Fieser and A. I. Ardao, *J. Am. Chem. Soc.*, **78**, 774

(1956).

14) L. F. Fieser, *J. Am. Chem. Soc.*, **70**, 3165 (1948).

15) S. C. Hooker and A. Steyermark, *J. Am. Chem. Soc.*, **58**, 1202 (1936).
