

Kinetic Resolution of Racemic Ferrocenylphosphine Compounds by Enantioselective Oxidation Using Cyclic Selenoxides Having a Chiral Ligand

Yoshihiro Miyake, Akiyoshi Yamauchi, Yoshiaki Nishibayashi, and Sakae Uemura*

Department of Energy and Hydrocarbon Chemistry, Graduate School of Engineering, Kyoto University, Sakyo-ku, Kyoto 606-8501

(Received August 19, 2002)


Cyclic selenoxides having an optically active binaphthyl skeleton work as the reagents for enantioselective oxidation of phosphines to the corresponding phosphine oxides. Treatment of a racemic 2-oxazolin-2-ylferrocenylphosphine with one of the selenoxides in carbon tetrachloride in the presence of phenol affords the corresponding phosphine oxide together with the unreacted starting phosphine, both with moderate enantioselectivities (the phosphine oxide, up to 13% ee; the phosphine, up to 29% ee).

Ferrocene and its derivatives play an important role in many areas of synthetic and material chemistry, and thus, they have been incorporated into complex structures which display unusual properties.¹ From the viewpoint of stereochemistry, ferrocenes with planar chirality have been widely investigated. They are of increasing importance as chiral ligands in transition metal-catalyzed asymmetric reactions.² Ferrocenes with planar chirality have mainly been prepared by two methods: the optical resolution of racemic ferrocenes and the diastereoselective *ortho*-lithiation of ferrocenes having an optically active directing group.³ Recently, several groups have reported examples of direct asymmetric synthesis of ferrocenes bearing only planar chirality, where the enantioselective *ortho*-lithiation of prochiral ferrocenes⁴ and enantioselective intramolecular insertion of carbenoids into C–H bonds of ferrocenes⁵ have been investigated.

Organic selenoxides are known to be the reagents for oxidation of phosphines to the corresponding phosphine oxides.⁶ However, no successful report on the enantioselective oxidation of racemic phosphines using selenoxides has appeared so far.⁷ We now report the preparation of cyclic selenoxides **1** having an optically active binaphthyl skeleton⁸ and the application of **1** to the kinetic resolution of racemic ferrocenylphosphine compounds with planar chirality, namely enantioselective oxidation of the phosphine compounds.

Results and Discussion

At first, we designed and prepared C_2 -symmetric cyclic selenides having an optically active binaphthyl skeleton and converted them into the corresponding selenoxides (Scheme 1). In these selenoxides, it might not be necessary to consider the configuration around the selenium atom at the oxidation step because of the C_2 -symmetric binaphthyl skeleton.⁹ The dibromides **2** were prepared from the corresponding (*R*)-BINOLs¹⁰ in three steps according to the literature method.¹¹ The reaction of **2** with selenium anion afforded the corre-



Scheme 1.





Fig. 1. An ORTEP drawing of **1a**. Selected bond lengths, torsion angle ($\text{\AA}$, deg): Se1–O1 = 1.642 (9); C1–C2–C3–C4 = -68.0 (2).

sponding cyclic selenides **3**, which were then oxidized with *m*-chloroperbenzoic acid (mCPBA) to the selenoxides **1** having an optically active binaphthyl skeleton, in good yields. The molecular structure of the selenoxide **1a** was unambiguously clarified by X-ray structural determination, an ORTEP drawing being shown in Fig. 1.

First, we investigated kinetic resolution of racemic 2-oxazolin-2-ylferrocenylphosphine **4a** using the selenoxide **1a** (Scheme 2). Typical results are shown in Table 1. The reac-



Scheme 2.

Table 1. Kinetic Resolution of Oxazolinylferrocenylphosphine **4a** Using **1a**^{a)}

Entry	Additive	4a		5a	
		Recovered yield (%) ^{b)}	Ee (%) ^{c),d)}	Yield (%) ^{b)}	Ee (%) ^{c),e)}
1	—	35	15	53	12
2	H ₂ O	26	19	51	6
3	TfOH	41	4	58	5
4	Pyridine·HCl	44	17	56	8
5 ^{f)}	PhOH	48	29	52	13
6 ^{g)}	PhOH	45	18	35	15
7	6a	26	19	45	5
8	6b	31	16	53	7
9	6c	36	12	56	8
10	6d	36	18	57	11
11	6e	33	15	55	9
12	6f	30	13	49	8
13	6g	37	9	49	8
14	6h	34	8	50	6
15 ^{h)}	PhOH	33	1	45	1

a) Reaction conditions: **4a** (0.2 mmol), **1a** (0.1 mmol), additive (0.1 mmol), CCl₄ (2 mL), at room temperature for 24 h. b) Determined by ³¹P-NMR. c) Determined by HPLC analysis using suitable chiral columns (see experimental section). d) Absolute configuration is *S*. e) Absolute configuration is *R*. f) The efficiency of the kinetic resolution, the *k_f/k_s* value,¹² is estimated to be 2.3. g) At 0 °C. h) **1b** was used in place of **1a**.

tion of the racemic **4a** with **1a** (0.5 molar amount) in CCl₄ at room temperature for 24 h proceeded smoothly to give the cor-

responding phosphine oxide **5a** in 53% yield with 12% ee. The unreacted phosphine **4a** was recovered in 35% yield. It


Table 2. Kinetic Resolution of **4** Using **1a**^{a)}

Entry	Substrate	Additive	4		5	
			Recovered yield (%) ^{b)}	Ee (%) ^{c)}	Yield (%) ^{b)}	Ee (%) ^{c)}
1	4b	—	42	9	47	13
2	4b	PhOH	49	18	43	21
3	4c	—	49	6	39	12
4	4c	PhOH	67	9	33	22
5	4d	PhOH	59	6 ^{d)}	41	1 ^{e)}
6 ^{f)}	4e	—	28	1	46	3
7 ^{f),g)}	4e	—	48	8	41	10
8 ^{f),h)}	4e	—	48	8	41	9
9 ^{f),i)}	4e	—	14	13	46	5

a) Reaction conditions: **4** (0.2 mmol), **1a** (0.1 mmol), CCl₄ (2 mL), at room temperature for 24 h. b) Determined by ³¹P-NMR. c) Determined by HPLC analysis using suitable chiral columns. d) Determined by optical rotation. e) Determined by optical rotation after conversion to the corresponding phosphine **4d**. f) For 3 h. g) CH₂Cl₂ was used in place of CCl₄. h) ClCH₂CH₂Cl was used in place of CCl₄. i) C₆H₅Cl was used in place of CCl₄.

was revealed to be an enantiomer rich compound of 15% ee (Table 1, Entry 1). Then the effect of additives on the enantioselectivity of either or both compounds was examined; eventually it was disclosed that the addition of phenol was slightly effective for improving it (Table 1, Entries 2–14). Thus, **4a** was recovered with 29% ee and **5a** was obtained with 13% ee (Table 1, Entry 5). The efficiency of the kinetic resolution, the k_t/k_s value,¹² is estimated to be 2.3. When phenol was added to the reaction mixture before the addition of racemic **4a**, a white precipitate was produced, while the mixture changed to a clear solution when the reaction was completed. The white precipitate might be the adduct between **1a** and phenol¹³ and the existence of interaction such as hydrogen bonding^{8,14} between the oxygen atom of **1a** and the hydroxy group of phenol was expected. At present we consider that the interaction may change the chiral environment around the selenoxide. The reactions in other solvents such as tetrahydrofuran (THF), dichloromethane and 1,2-dichloroethane were slow and low enantioselectivities were obtained. When the selenoxide **1b** was used as an oxidant instead of **1a**, no asymmetric induction occurred, unfortunately (Table 1, Entry 15). In some cases, the formation of some unidentified products was observed by ³¹P-NMR.

Other racemic oxazolinylferrocenylphosphines (**4b** and **4c**) and *N,N*-dimethylaminomethylferrocenylphosphine (**4d**) were investigated for this kinetic resolution by using **1a**. Typical results are shown in Table 2. In all cases, the addition of phenol was revealed to be slightly effective for enantioselectivity, but higher enantioselectivity than that obtained using **4a** was not observed. Treatment of racemic **4d** under similar reaction conditions afforded **5d** in 41% yield with only 1% ee and the unreacted **4d** was recovered in 59% yield with 6% ee (Table 2, Entry 5). Interestingly, the reaction of racemic ferrocenylphosphine having a hydroxy group (**4e**) proceeded faster than that of **4a** (Table 2, Entry 6). The reaction of **4e** in a variety of solvents such as dichloromethane, 1,2-dichloroethane and chlorobenzene also proceeded smoothly to give **5e**, but the enantioselectivity was lower than that obtained using **4a** (Table 2, Entries 7–9). The hydrogen bonding between the oxygen atom of **1a** and the hydroxy group of **4e** probably accelerated



Scheme 3.

the reaction rate.

Next, we attempted the kinetic resolution of racemic phosphines **7**, hoping to obtain *P*-chiral phosphines, which have recently been shown to be effective ligands for the rhodium-catalyzed asymmetric hydrogenation (Scheme 3).¹⁵ Treatment of the racemic methyl(1-naphthyl)phenylphosphine (**7a**) with **1a** (0.5 molar amount) in the presence of phenol (0.5 equiv) in CCl₄ at room temperature for 3 h gave the corresponding phosphine oxide **8a** in 62% yield with 5% ee. The unreacted phosphine **7a** was recovered in 38% yield, but only with 5% ee. Typical results are shown in Table 3. Similarly, no satisfactory results were obtained in the oxidation of racemic methyl(2-naphthyl)phenylphosphine (**7b**) or (2-hydroxy-2,2-diphenylethyl)methyl(phenyl)phosphine (**7c**) (Table 3, Entries 3–5).

In summary, we have prepared cyclic selenoxides **1** having an optically active binaphthyl skeleton and attempted the kinetic resolution of racemic ferrocenylphosphine **4** using **1** as oxidants. Although the enantioselectivity of the produced phosphine is not yet high (up to 29% ee), this conceptually novel methodology may be interesting for the preparation of optically active ferrocene derivatives with only planar chirality.

Experimental

General Procedures. ¹H- and ¹³C-NMR spectra were recorded on 300 or 400 and 100 MHz FT-NMR spectrometer, respectively. ³¹P NMR spectra were recorded on a 160 MHz FT-NMR spectrometer using P(OMe)₃ as an external standard. ⁷⁷Se NMR spec-

Table 3. Kinetic Resolution of **7** Using **1a**^{a)}

Entry	Substrate	Additive	Time (h)	7		8	
				Recovered yield (%) ^{b)}	Ee (%) ^{c)}	Yield (%) ^{b)}	Ee (%) ^{d)}
1	7a	—	3	38	5	62	5
2	7a	phenol	3	37	0	63	2
3	7b	—	3	10	16	14	— ^{e)}
4	7b	phenol	3	57	0	43	9
5	7c	—	0.1	22	5	30	19

a) Reaction conditions: **7** (0.2 mmol), **1a** (0.1 mmol), additive (0.1 mmol), CCl₄ (2 mL), at room temperature. b) Determined by ³¹P-NMR. c) Determined by HPLC analysis using suitable chiral columns after conversion to the corresponding phosphine-borane. Absolute configuration is not determined. d) Determined by HPLC analysis using suitable chiral columns. Absolute configuration is not determined. e) Not determined.

tra were recorded on a 76 MHz FT-NMR spectrometer using Me₂Se as an external standard. Chemical shifts are reported in ppm relative to TMS in the solvents specified. High-resolution mass spectra (HRMS) were obtained with a JEOL JMS-SX102A spectrometer. Melting points are uncorrected. Column chromatographies on SiO₂ were performed with Merck silica gel 60. HPLC analyses were carried out on a HITACHI L-7100 and an L-7400 UV detector using Daicel Chiralcel OD, AD, and OJ columns at 30 °C. Elemental analyses were performed at the Microanalytical Center of Kyoto University. All reactions were performed in oven-dried or flame-dried glassware under an atmosphere of N₂ unless otherwise noted. Solvents and chemicals were obtained commercially and purified by standard procedures. Racemic phosphine compounds (**4a**,¹⁶ **4b**,¹⁷ **4c**,¹⁷ **4d**,¹⁸ **4e**,^{18,19} **7a**,²⁰ **7b**,²⁰ and **7c**²¹) were prepared according to literature procedures. Optically active BINOLs were commercially available compounds.

Preparation of (R)-3,5-Dihydrodinaphtho[2,1-c:1',2'-e]selenepin (3a). A solution of (R)-2,2'-bis(bromomethyl)-1,1'-binaphthalene (**2a**)^{11a} (2.640 g, 6.00 mmol) and *tert*-butyl alcohol (0.35 mL, 3.00 mmol) in THF (4 mL) was added at room temperature to Li₂Se²² (5.55 mmol) suspended in THF (8 mL), and the mixture was stirred at room temperature for 12 h. Then the reaction mixture was taken up in Et₂O/H₂O. After separation, the aqueous layer was extracted with Et₂O (2 × 40 mL). The ether layer was dried over MgSO₄ and concentrated under vacuum. Purification by column chromatography on SiO₂ with hexane/CH₂Cl₂ (20/1) as an eluent gave the title compound **3a** (1.90 g, 5.29 mmol, 95%): A white solid; mp 169.1–170.0 °C; [α]_D²⁵ –7.47 (c 0.1, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 3.46 (d, *J* = 10.8 Hz, 2H), 3.51 (d, *J* = 10.8 Hz, 2H), 7.18–7.25 (m, 4H), 7.40–7.45 (m, 2H), 7.54 (d, *J* = 8.0 and 13.6 Hz, 2H), 7.92 (dd, *J* = 7.8 and 13.6 Hz, 4H); ¹³C NMR (100 MHz, CDCl₃) δ 24.7, 125.4, 126.1, 126.3, 126.6, 128.2, 129.1, 132.0, 132.8, 133.4, 134.5; ⁷⁷Se NMR (76 MHz, CDCl₃) δ 438.2. Found: C, 73.50; H, 4.79%. Calcd for C₂₂H₁₆Se: C, 73.54; H, 4.49%.

(R)-2,6-Diphenyl-3,5-dihydrodinaphtho[2,1-c:1',2'-e]selenepin (3b). The compound was similarly prepared from **2b**.^{11b} A pale yellow solid; 43%; mp 175.5–176.1 °C; [α]_D²⁵ –66.5 (c 0.1, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 3.38 (d, *J* = 11.4 Hz, 2H), 3.61 (d, *J* = 11.4 Hz, 2H), 7.11–7.20 (m, 2H), 7.20–7.27 (m, 2H), 7.35–7.51 (m, 8H), 7.67 (brs, 4H), 7.81–7.92 (m, 4H); ¹³C NMR (100 MHz, CDCl₃) δ 22.2, 125.8, 126.0, 126.7, 127.1, 127.3, 128.1, 128.2, 129.7, 130.1, 131.4, 132.3, 132.5, 134.3, 139.0, 141.2; ⁷⁷Se NMR (76 MHz, CDCl₃) δ 440.6. HRMS *m/z* Found: 512.1049. Calcd for C₃₄H₂₄Se M⁺: 512.1046. Found: C, 80.31; H,

5.24%. Calcd for C₃₄H₂₄Se: C, 79.83; H, 4.73%.

Preparation of (R)-3,5-Dihydrodinaphtho[2,1-c:1',2'-e]selenepin 4-Oxide (1a). To a solution of **3a** (0.359 g, 1.00 mmol) in CH₂Cl₂ (30 mL) were added mCPBA (80%, 0.237 g, 1.1 mmol) and saturated aqueous K₂CO₃ solution (6 mL) and the resulting solution was stirred at room temperature for 10 min. The mixture was poured into saturated aqueous Na₂CO₃ solution and extracted with CH₂Cl₂ (2 × 30 mL). The combined organic layer was then dried over MgSO₄ and concentrated under vacuum to leave a crude product. Purification by column chromatography on SiO₂ with ethyl acetate/MeOH (1/1) as an eluent gave the title compound **1a** (0.336 g, 0.90 mmol, 90%). A white solid; mp 84.2–85.5 °C; [α]_D²⁵ –43.4 (c 0.1, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 3.18 (d, *J* = 10.8 Hz, 1H), 3.55 (d, *J* = 12.5 Hz, 1H), 3.81 (d, *J* = 12.5 Hz, 1H), 4.36 (d, *J* = 10.8 Hz, 1H), 7.21–7.34 (m, 4H), 7.48–7.55 (m, 3H), 7.63–7.65 (m, 1H), 7.93–8.04 (m, 4H); ¹³C NMR (100 MHz, CDCl₃) δ 48.9, 54.2, 126.2, 126.5, 126.9, 126.9, 128.0, 128.3, 128.5, 128.8, 129.0, 129.0, 129.1, 131.8, 132.4, 133.6, 133.7, 134.4, 134.6; ⁷⁷Se NMR (76 MHz, CDCl₃) δ 1033.3. HRMS *m/z* Found: 377.0446. Calcd for C₂₂H₁₇OSe M⁺: 377.0446. Found: C, 69.55. H, 4.73%. Calcd for C₂₂H₁₇OSe: C, 70.40; H, 4.30%.

X-ray Structural Analysis of 1a. Single crystals of **1a** (C₂₂H₁₆OSe·C₆H₆) suitable for X-ray analysis were prepared by recrystallization from benzene. Diffraction data were collected on a Rigaku RAXIS imaging plate area detector with Mo K_α (λ = 0.711 Å) radiation and a graphite monochromator at 23 °C. Details of crystal and data collection parameters are summarized in Table 4. For a structure analysis and refinement, computations were performed using the CrystalStructure²³ crystallographic software package of Molecular Structure. Neutral atom scattering factors were taken from Ref. 24. Anomalous dispersion effects were included in *F*_{calc}²⁵; the values of Δ*f*' and Δ*f*'' were those of Ref. 26. The structure was solved by direct methods (SIR92). All non-hydrogen atoms were refined anisotropically. Hydrogen atoms were included, but not refined.

Crystallographic data have been deposited at the CCDC, 12 Union Road, Cambridge CB2 1EZ, UK and copies can be obtained on request, free of charge, by quoting the publication citation and the deposition numbers CCDC-196734.

(R)-2,6-Diphenyl-3,5-dihydrodinaphtho[2,1-c:1',2'-e]selenepin 4-Oxide (1b). The compound was similarly prepared from **3b**. A pale yellow solid; 72%; mp 76.1–77.0 °C; [α]_D²⁵ –5.30 (c 0.1, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 3.30 (d, *J* = 12.8 Hz, 1H), 3.35 (d, *J* = 10.8 Hz, 1H), 4.28 (d, *J* = 12.8 Hz, 1H), 4.47 (d, *J* = 10.8 Hz, 1H), 7.00–8.15 (m, 20H); ¹³C NMR (100 MHz,

Table 4. Summary of Crystallographic Data of **1a**

Empirical formula	C ₂₂ H ₁₆ OSe·C ₆ H ₆
Fw	453.44
Crystal system	orthorhombic
Space group	<i>P</i> 2 ₁ 2 ₁ 2 ₁ (#19)
Crystal color	colorless
<i>a</i> /Å	8.0215(7)
<i>b</i> /Å	12.529(1)
<i>c</i> /Å	21.105(2)
<i>V</i> (Å ³)	2121.1(3)
<i>Z</i>	4
<i>D</i> _{calc} /g cm ⁻³	1.420
μ (Mo <i>K</i> α)/cm ⁻¹	1.95
<i>F</i> (000)	928.00
Radiation	Mo <i>K</i> α (λ = 0.71069 Å) graphite monochromated
Temp/°C	23.0
Scan type	ω
Max. 2 θ /°	55.0
No. of rflns measd	18857
No. of observns (<i>I</i> > 3.00 σ (<i>I</i>))	1349
Structure soln	direct methods (SIR92)
Refinement	full-matrix least squares
No. of variables	287
Reflection/parameter ratio	4.70
Residuals: <i>R</i> ; <i>R</i> _w	0.039; 0.044
GOF	0.70
Max shift/error in final cycle	3.58
Maximum peak in final diff map (e Å ⁻³)	0.58
Minimum peak in final diff map (e Å ⁻³)	-0.64

CDCl₃) δ 44.2, 51.3, 126.3, 126.4, 126.6, 126.7, 126.9, 127.1, 127.3, 127.9, 128.2, 128.3, 128.7, 129.3, 129.6, 130.0, 131.3, 131.6, 132.8, 132.9, 135.5, 135.7, 139.4, 140.1, 141.5. Found: C, 77.50%; H, 4.84%. Calcd for C₃₄H₂₄OSe: C, 77.41%; H, 4.59%.

Typical Procedure for Kinetic Resolution of Racemic Ferrocenylphosphine Compounds. A solution of **1a** (37.5 mg, 0.1 mmol) and phenol (9.4 mg, 0.1 mmol) in carbon tetrachloride (1 mL) was stirred under nitrogen atmosphere at room temperature. After 15 min, the racemic **4a** (93.5 mg, 0.2 mmol) in carbon tetrachloride (1 mL) was added to the solution and the resulting solution was stirred at room temperature for 24 h. Yields of phosphine and phosphine oxide were determined by integration of the ³¹P resonances against P(OMe)₃ added as an internal reference. Purification of the crude products by SiO₂ chromatography gave the corresponding ferrocenylphosphine and ferrocenylphosphine oxide in a pure form.

[2-(4,4-Dimethyl-2-oxazolin-2-yl)ferrocenyl]diphenylphosphine (4a). Enantiomeric excess was determined by Daicel Chiralcel AD, 30 °C, 0.5 mL/min, hexane/2-propanol = 95/5, *t*₁ = 9.27 (*S*) min, *t*₂ = 11.28 min (*R*).¹⁶

[2-(4,4-Dimethyl-2-oxazolin-2-yl)ferrocenyl]diphenylphosphine Oxide·H₂O (5a·H₂O). A reddish brown solid; mp 115.1–116.0 °C; enantiomeric excess was determined by Daicel Chiralcel AD, 30 °C, 0.5 mL/min, hexane/2-propanol = 90/10, *t*₁ = 45.77 (*S*) min, *t*₂ = 48.65 min (*R*); ¹H NMR (400 MHz, CDCl₃) δ 1.03 (brs, 3H), 1.19 (brs, 3H), 1.35 (br, 2H, *H*₂O), 2.93 (brs, 1H), 3.95 (brs, 2H), 4.49 (m, 6H), 5.09 (brs, 1H), 7.41–7.70 (m, 10H);

³¹P NMR (160 MHz, CDCl₃) δ 23.6. HRMS *m/z* Found: 484.1136. Calcd for C₂₇H₂₇FeNO₂P M⁺ + H: 484.1129. Found: C, 64.76%; H, 5.43%; N, 2.63%. Calcd for C₂₇H₂₆FeNO₂P·H₂O: C, 64.69%; H, 5.63%; N, 2.79%. The presence of H₂O was confirmed by ¹H NMR spectroscopy.

[2-(4-Isopropyl-2-oxazolin-2-yl)ferrocenyl]diphenylphosphine (4b).¹⁷ Enantiomeric excess was determined by Daicel Chiralcel AD, 30 °C, 0.5 mL/min, hexane/2-propanol = 95/5, *t*₁ = 8.20 min, *t*₂ = 9.82 min.

[2-(4-Isopropyl-2-oxazolin-2-yl)ferrocenyl]diphenylphosphine Oxide·H₂O (5b·H₂O). A reddish brown solid; mp 192.2–192.5 °C; enantiomeric excess was determined by Daicel Chiralcel AD, 30 °C, 1.0 mL/min, hexane/2-propanol = 90/10, *t*₁ = 31.84 min, *t*₂ = 41.72 min; ¹H NMR (400 MHz, CDCl₃) δ 0.61 (brs, 3H), 0.80 (brs, 3H), 1.47–1.53 (m, 1H), 2.05 (br, 2H, *H*₂O), 3.13 (m, 1H), 3.75 (m, 1H), 3.93 (brs, 1H), 4.21 (m, 1H), 4.49 (m, 1H), 4.50 (brs, 5H), 5.07 (brs, 1H), 7.27–7.74 (m, 10H); ³¹P NMR (160 MHz, CDCl₃) δ 23.7. HRMS *m/z* Found: 498.1293. Calcd for C₂₈H₂₉FeNO₂P M⁺ + H: 498.1286. Found: C, 65.28%; H, 5.62%; N, 2.59%. Calcd for C₂₈H₂₈FeO₂P·H₂O: C, 65.26%; H, 5.87%; N, 2.72%. The presence of H₂O was confirmed by ¹H NMR spectroscopy.

[2-(4-Benzyl-2-oxazolin-2-yl)ferrocenyl]diphenylphosphine (4c).¹⁷ Enantiomeric excess was determined by Daicel Chiralcel AD, 30 °C, 0.5 mL/min, hexane/2-propanol = 95/5, *t*₁ = 11.29 min, *t*₂ = 14.78 min.

[2-(4-Benzyl-2-oxazolin-2-yl)ferrocenyl]diphenylphosphine Oxide·H₂O (5c·H₂O). A reddish brown solid; mp 193.2–193.7 °C; enantiomeric excess was determined by Daicel Chiralcel AD, 30 °C, 1.0 mL/min, hexane/2-propanol = 90/10, *t*₁ = 41.11 min, *t*₂ = 46.72 min; ¹H NMR (400 MHz, CDCl₃) δ 1.75 (br, 2H, *H*₂O), 2.12 (dd, *J* = 12.3 and 14.7 Hz, 1H), 3.00 (dd, *J* = 12.3 and 13.6 Hz, 1H), 3.06 (dd, *J* = 8.0 and 8.4 Hz, 1H), 3.98 (brs, 1H), 4.12 (dd, *J* = 8.0 and 8.4 Hz, 1H), 4.20 (m, 1H), 4.39 (brs, 1H), 4.50 (s, 5H), 5.05 (brs, 1H), 7.03–7.77 (m, 15H); ³¹P NMR (160 MHz, CDCl₃) δ 23.7. HRMS *m/z* Found: 546.1273. Calcd for C₃₂H₂₉FeNO₂P M⁺ + H: 546.1286. Found: C, 68.79%; H, 5.56%; N, 2.40%. Calcd for C₃₂H₂₈FeNO₂P·H₂O: C, 68.22%; H, 5.37%; N, 2.49%. The presence of H₂O was confirmed by ¹H NMR spectroscopy.

[2-(*N,N*-Dimethylaminomethyl)ferrocenyl]diphenylphosphine (4d). Enantiomeric excess was determined by optical rotation.¹⁸

[2-(*N,N*-Dimethylaminomethyl)ferrocenyl]diphenylphosphine Oxide·H₂O (5d·H₂O). A reddish brown solid; mp 59.2–59.9 °C; enantiomeric excess of **5d** was determined by optical rotation after the reduction of **5d** into the corresponding phosphine¹⁸; ¹H NMR (400 MHz, CDCl₃) δ 1.96 (s, 6H), 3.00 (br, 2H, *H*₂O), 3.43 (d, *J* = 13.0, 2H), 3.94 (s, 1H), 4.20 (s, 5H), 4.36 (s, 1H), 4.67 (s, 1H), 7.21–7.92 (m, 10H); ³¹P NMR (160 MHz, CDCl₃) δ 23.3. HRMS *m/z* Found: 444.1183. Calcd for C₂₅H₂₇FeNOP M⁺ + H: 444.1180. Found: C, 65.47%; H, 5.93%; N, 2.70%. Calcd for C₂₅H₂₆FeNOP·H₂O: C, 65.09%; H, 6.12%; N, 3.04%. The presence of H₂O was confirmed by ¹H NMR spectroscopy.

[2-(Hydroxymethyl)ferrocenyl]diphenylphosphine (4e).^{18,19} Enantiomeric excess was determined by Daicel Chiralcel AD, 30 °C, 0.5 mL/min, hexane/2-propanol = 95/5, *t*₁ = 22.77 min, *t*₂ = 29.04 min.

[2-(Hydroxymethyl)ferrocenyl]diphenylphosphine Oxide·H₂O (5e·H₂O). A reddish brown solid; mp 91.5–92.3 °C; enantiomeric excess was determined by Daicel Chiralcel OD, 30 °C, 0.5 mL/min, hexane/2-propanol = 95/5, *t*₁ = 31.96 min, *t*₂ = 37.96 min; ¹H NMR (400 MHz, CDCl₃) δ 3.93 (s, 1H), 4.24–4.36

(m, 3H), 4.27 (s, 5H), 4.55 (s, 1H), 5.51 (s, 1H), 7.25–7.81 (m, 10H); ^{31}P NMR (160 MHz, CDCl_3) δ 30.9. HRMS m/z Found: 416.0627. Calcd for $\text{C}_{23}\text{H}_{21}\text{FeO}_2\text{P}$ M^+ : 416.0629. Found: C, 65.94; H, 5.25%. Calcd for $\text{C}_{23}\text{H}_{21}\text{FeO}_2\text{P}$: C, 66.37; H, 5.09%.

Methyl(1-naphthyl)phenylphosphine (7a).²⁰ Enantiomeric excess of **7a** was determined by Daicel Chiralcel OD, 30 °C, 1.0 mL/min, hexane/2-propanol = 90/10, t_1 = 6.41 min, t_2 = 6.98 min, after the transformation of **7a** into the corresponding phosphine-borane.^{20c}

Methyl(1-naphthyl)phenylphosphine Oxide (8a).²⁰ Enantiomeric excess was determined by Daicel Chiralcel OJ, 30 °C, 1.0 mL/min, hexane/2-propanol = 90/10, t_1 = 33.50 min, t_2 = 37.90 min.

Methyl(2-naphthyl)phenylphosphine (7b).²⁰ Enantiomeric excess of **7b** was determined by Daicel Chiralcel OD, 30 °C, 1.0 mL/min, hexane/2-propanol = 90/10, t_1 = 15.98 min, t_2 = 19.80 min, after the transformation of **7b** into the corresponding phosphine-borane.^{20c}

Methyl(2-naphthyl)phenylphosphine Oxide (8b).²⁰ Enantiomeric excess was determined by Daicel Chiralcel AD, 30 °C, 0.5 mL/min, hexane/2-propanol = 90/10, t_1 = 28.78 min, t_2 = 32.97 min.

(2-Hydroxy-2,2-diphenylethyl)methyl(phenyl)phosphine (7c).²¹ Enantiomeric excess of **7c** was determined by Daicel Chiralcel OD, 30 °C, 0.5 mL/min, hexane/2-propanol = 92/8, t_1 = 26.42 min, t_2 = 33.54 min, after the transformation of **7c** into the corresponding phosphine-borane.^{20c}

(2-Hydroxy-2,2-diphenylethyl)methyl(phenyl)phosphine Oxide (8c).²¹ Enantiomeric excess was determined by Daicel Chiralcel OD, 30 °C, 0.5 mL/min, hexane/2-propanol = 85/15, t_1 = 28.40 min, t_2 = 36.18 min.

This work was supported by a Grant-in-Aid for Scientific Research (Nos. 13305062 and 12750747) from the Ministry of Education, Science, Sports, and Culture. We also thank Dr. Izuru Takei at The University of Tokyo for his generous gift of (S)-**4a**.

References

- For a review see: A. Togni and T. Hayashi, "Ferrocenes," VCH: Weinheim, (1995) and Refs. cited therein.
- For a review see; M. Sawamura and Y. Ito, *Chem. Rev.*, **92**, 857 (1992).
- For representative examples of diastereoselective lithiation of chiral ferrocenes; a) D. Marquarding, H. Klusacek, G. Gokel, P. Hoffmann, and I. Ugi, *J. Am. Chem. Soc.*, **92**, 5389 (1970). b) O. Riant, O. Samuel, and H. B. Kagan, *J. Am. Chem. Soc.*, **115**, 5835 (1993). c) F. Reibière, O. Riant, L. Ricard, and H. B. Kagan, *Angew. Chem., Int. Ed. Engl.*, **32**, 568 (1993). d) C. J. Richards, T. Damalidis, D. E. Hibbs, and M. B. Hursthouse, *Synlett*, **1995**, 74. e) T. Sammakia, H. A. Latham, and D. R. Schaad, *J. Org. Chem.*, **60**, 10 (1995). f) Y. Nishibayashi and S. Uemura, *Synlett*, **1995**, 79. g) W. Zhang, T. Hirao, and I. Ikeda, *Tetrahedron Lett.*, **37**, 4545 (1996). h) K. H. Ahn, C.-W. Cho, H.-H. Baek, J. Park, and S. Lee, *J. Org. Chem.*, **61**, 4937 (1996).
- a) D. Price and N. S. Simpkins, *Tetrahedron Lett.*, **36**, 6135 (1995). b) M. Tsukazaki, M. Tinkl, A. Roglans, B. J. Chapell, N. J. Taylor, and V. Snieckus, *J. Am. Chem. Soc.*, **118**, 685 (1996). c) Y. Nishibayashi, Y. Arikawa, K. Ohe, and S. Uemura, *J. Org. Chem.*, **61**, 1172 (1996).
- S. Siegel and H.-G. Schmalz, *Angew. Chem., Int. Ed. Engl.*, **36**, 2456 (1997).
- a) M. Mikołajczyk and J. Łuczak, *J. Org. Chem.*, **43**, 2132 (1978). b) J. Drabowicz, J. Łuczak, P. Łyzwa, and M. Mikołajczyk, *Phosphorus, Sulfur, Silicon, Relat. Elem.*, **143**, 136 (1998).
- The example of the enantioselective oxidation of racemic phosphines with chiral oxoruthenium porphyrins has been reported by Simonneaux et al.; P. L. Maux, H. Bahri, G. Simonneaux, and L. Toupet, *Inorg. Chem.*, **34**, 4691 (1995).
- This work was partly presented at XVth International Conference on Phosphorus Chemistry July 29–August 3, 2001 Sendai, Japan, Abstracts p. 210. Rayner et al. independently reported the preparation of the selenoxide **1a** and its use in the oxidation of sulfides; D. J. Procter and C. M. Rayner, *Synth. Commun.*, **30**, 2975 (2000).
- a) F. A. Davis, O. D. Stringer, and R. T. Reddy, *Tetrahedron*, **41**, 4747 (1985). b) K. B. Sharpless, M. W. Young, and R. F. Lauer, *Tetrahedron Lett.*, **1973**, 1979. c) M. Oki and H. Iwamura, *Tetrahedron Lett.*, **1966**, 2917. d) W. J. Burlant and E. S. Gould, *J. Am. Chem. Soc.*, **76**, 5775 (1954). e) T. W. Campbell, H. G. Walker, and G. M. Coppinger, *Chem. Rev.*, **50**, 279 (1952). f) W. R. Gaythwaite, J. Kenyon, and H. Phillips, *J. Chem. Soc.*, **1928**, 2280.
- (R)-3,3'-Diphenyl-1,1'-bi-2-naphthol can be prepared from (R)-BINOL; J. M. Chong, L. Shen, and N. J. Taylor, *J. Am. Chem. Soc.*, **122**, 1822 (2000).
- a) I. G. Stará, P. Sary, M. Tichý, J. Závada, and P. Fiedler, *J. Org. Chem.*, **59**, 1326 (1994). b) T. Ooi, M. Kameda, and K. Maruoka, *J. Am. Chem. Soc.*, **121**, 6519 (1999).
- The efficiency of the kinetic resolution is characterized by k_f/k_s , the selectivity factor = (rate of fast-reacting enantiomer)/(rate of slow-reacting enantiomer). For a review, R. Noyori, M. Tokunaga, and M. Kitamura, *Bull. Chem. Soc. Jpn.*, **68**, 36 (1995).
- When LRMS spectra of the white precipitate were measured, the fragment peak of **1a** (m/z = 512) was obtained.
- a) The preparation of the salts of selenoxides with sulfonic acid has been reported and the salts worked as oxidants of sulfides to sulfoxides; D. J. Procter, M. Thornton-Pett, and C. M. Rayner, *Tetrahedron*, **52**, 1841 (1996). b) F. Toda and K. Mori, *J. Chem. Soc., Chem. Commun.*, **1986**, 1357.
- a) T. Imamoto, J. Watanabe, Y. Wada, H. Masuda, H. Yamada, H. Tsuruta, S. Matsukawa, and K. Yamaguchi, *J. Am. Chem. Soc.*, **120**, 1635 (1998). b) Y. Yamanori and T. Imamoto, *J. Org. Chem.*, **64**, 2988 (1999).
- a) I. Takei, Ph.D. Thesis, The University of Tokyo (2002). (S)-**4a** was prepared from (S)-2-iodoferrocenylcarboxylic acid by the modified Bolm's procedure; C. Bolm, K. M. Fernandez, A. Seger, G. Raabe, and K. Gunther, *J. Org. Chem.*, **63**, 7860 (1998). b) S.-L. You, X.-L. Hou, L.-X. Dai, Y.-H. Yu, and W. Xia, *J. Org. Chem.*, **67**, 4684 (2002).
- Y. Nishibayashi, K. Segawa, Y. Arikawa, K. Ohe, M. Hidai, and S. Uemura, *J. Organomet. Chem.*, **545-546**, 381 (1997).
- T. Hayashi, T. Mise, M. Fukushima, M. Kagotani, N. Nagashima, Y. Hamada, A. Matsumoto, S. Kawakami, M. Konishi, K. Yamamoto, and M. Kumada, *Bull. Chem. Soc. Jpn.*, **53**, 1138 (1980).
- O. Riant, O. Samuel, T. Flessner, S. Taudien, and H. B. Kagan, *J. Org. Chem.*, **62**, 6733 (1997).
- a) N. G. Andersen, P. D. Ramsden, D. Che, M. Parvez, and

- B. A. Keay, *J. Org. Chem.*, **66**, 7478 (2001). b) T. Imamoto, S.-i. Kikuchi, T. Miura, and Y. Wada, *Org. Lett.*, **3**, 87 (2001). c) T. Imamoto, T. Oshiki, T. Onozawa, T. Kusumoto, and K. Sato, *J. Am. Chem. Soc.*, **112**, 5244 (1990).
- 21 A. R. Muci, K. R. Campos, and D. A. Evans, *J. Am. Chem. Soc.*, **117**, 9075 (1995).
- 22 J. A. Gladysz, J. L. Hornby, and J. E. Garbe, *J. Org. Chem.*, **43**, 1204 (1978).
- 23 a) CrystalStructure 2.00: Crystal Structure Analysis Package, Rigaku and MSC 2001. b) CRYSTALS Issue 10: D. J. Watkin, J. R. Carruthers, and P. W. Betteridge, Chemical Crystallography Laboratory, Oxford, UK.
- 24 D. T. Cromer and J. T. Waber, "International Tables for X-ray Crystallography," The Kynoch Press: Birmingham, England, 1974, Vol. IV, Table 2.2A.
- 25 J. A. Ibers and W. C. Hamilton, *Acta. Crystallogr.*, **17**, 781 (1964).
- 26 D. C. Creagh and J. H. Hubbell, in "International Tables for Crystallography," ed by A. J. C. Wilson, Kluwer Academic Publishers, Boston (1992), Vol. C, Tables 4.2 and 4.3, p. 200.