

Metal Cation-Exchanged Montmorillonite (M^{n+} -Mont)-Catalyzed Friedel–Crafts Acylation of 1-Methyl-1-cyclohexene and 1-Trimethylsilyl-1-alkynes

Takahiro Nishimura,* Seiji Ohtaka, Keiji Hashimoto,¹ Takayoshi Yamauchi,² Takuji Hasegawa,³ Kaori Imanaka,³ Jun-ichi Tateiwa,³ Hiroshi Takeuchi,^{3,4} and Sakae Uemura*

Department of Energy and Hydrocarbon Chemistry, Graduate School of Engineering, Kyoto University, Nishikyo-ku, Kyoto 615-8510

¹Osaka Municipal Technical Research Institute, 1-6-5 Morinomiya, Joto-ku, Osaka 536-8553

²Nissei Chemical Inc., 2-18-110 Jyuhachijyo, Yodogawa-ku, Osaka 532-0001

³Department of Molecular Science, Graduate School of Science and Technology, Kobe University, Rokkodai-cho, Nada-ku, Kobe 657-8501


⁴Department of Chemical Science and Engineering, Faculty of Engineering, Kobe University, Rokkodai-cho, Nada-ku, Kobe 657-8501

Received March 29, 2004; E-mail: uemura@sci.kyoto-u.ac.jp

The acylation of 1-methyl-1-cyclohexene and 1-trimethylsilyl-1-alkynes with acyl chlorides has been investigated in the presence of a variety of metal cation-exchanged montmorillonites (abbreviated as M^{n+} -monts), where the catalysts are recyclable for several times after simple washing.

Organic transformation in the presence of a variety of solid catalysts is of recent interest and many useful synthetic methods have been developed.¹ Currently we have been much interested in the use of clay minerals as the solid catalysts, since they are known as environmentally-friendly materials and might work as the substitutes for several homogeneous catalysts.^{2,3} Here we describe the result of our attempts for Friedel–Crafts acylation of 1-methyl-1-cyclohexene and 1-trimethylsilyl-1-alkynes with various acyl chlorides in the presence of cation-exchanged montmorillonites (abbreviated as M^{n+} -monts), a kind of clay mineral. Although aromatic Friedel–Crafts reactions have so far been investigated using various M^{n+} -monts as catalysts,⁴ there are no examples of aliphatic acylation in the presence of M^{n+} -monts to the best of our knowledge.

First, we examined Friedel–Crafts acylation of 1-methyl-1-cyclohexene (**1**) (2 molar amounts) with butyryl chloride (**2b**) (1 molar amount) in the presence of several M^{n+} -monts



Scheme 1.

Table 1. Acylation of **1** with **2b** in the Presence of M^{n+} -Mont

Entry	M^{n+} -mont	(mmol) ^a	Isolated yield of 3b ^b /%	TON ^c
1	Fe^{3+}	0.105	45	13
2	In^{3+}	0.059	25	13
3	Zr^{4+}	0.220	9	1
4	La^{3+}	0.060	Trace	0
5	Al^{3+}	0.159	0	0
6	Ce^{3+}	0.042	0	0
7	Yb^{3+}	0.097	0	0
8	Dy^{3+}	0.018	0	0
9 ^d	—	—	0	0

Reaction conditions: **1** (6.0 mmol), **2b** (3.0 mmol), M^{n+} -mont (300 mg), benzene (8 cm^3), rt, 2 h. a) The amount of acid sites of M^{n+} -mont was estimated by NH_3 -TPD analysis. b) Mol of **3b**/mol of acid sites. c) Based on the mol of acid sites. d) 24 h.

($2\text{--}7 \times 10^{-2}$ molar amounts) (Scheme 1, Table 1). Fe^{3+} and In^{3+} -monts were revealed to be effective for this reaction (entries 1 and 2), the TON (turnover number) being 13 in both cases. The catalytic activity of Zr^{4+} and La^{3+} -monts was lower (entries 3 and 4). On the other hand, Al^{3+} , Ce^{3+} , Yb^{3+} , and Dy^{3+} -monts did not show any catalytic activity (entries 5–8). No reaction proceeded in the absence of M^{n+} -mont (entry 9). The amount of Fe^{3+} -mont slightly affected the product yield and we found that the use of 0.012 molar amounts of Fe^{3+} -mont (100 mg) gave the best result, a larger and a smaller amount of it resulting in a decrease of the product yield.

Thus, Fe^{3+} and In^{3+} -monts were applied to the reaction of **1** with a variety of acyl chlorides **2a–2f** (Scheme 1, Table 2). As a result, the corresponding 6-acyl-1-methyl-1-cyclohexene was obtained in up to 58% yield, but some unidentified isomeric compounds were detected by GC–MS analysis.

The recovered Fe^{3+} -mont was successfully reused in the reaction of **1** with **2b** to afford **3b** at least three times after washing it with 50% aqueous acetone and drying at 120 $^\circ\text{C}$ for 24 h (1st use, 45%; 2nd use, 40%; 3rd use, 40%).


Separately, we confirmed that the acylation of **1** with **2b** proceeded even in the presence of a catalytic amount of FeCl_3 and InCl_3 , but the catalytic activity of these Lewis acids was not as high as that of Fe^{3+} and In^{3+} -monts in terms of the isolated yields of **3b** (15–31%).

Next, we investigated the reaction of several 1-trimethylsilyl-1-alkynes **4** with **2b** (Scheme 2, Table 3). Treatment of **4** with **2b** in the presence of Fe^{3+} and In^{3+} -monts at room temperature for 2 h produced the corresponding acyl compounds in moderate yields by replacing a trimethylsilyl group with an acyl group; the yield was up to 64% (entry 2). Even when an excess amount of **2b** to **4a** was employed, only a monosubstituted product **5a** was obtained without any formation of a disubstituted product.

Table 2. Acylation of **1** with **2** in the Presence of Fe³⁺ and In³⁺-Mont

Entry	2	M ⁿ⁺ -mont	Isolated yield of 3 /%	TON ^a)
1	2a	Fe ³⁺ -mont	49	42
2	2b	Fe ³⁺ -mont	58	50
3	2c	Fe ³⁺ -mont	38	33
4	2d	Fe ³⁺ -mont	19	16
5	2e	Fe ³⁺ -mont	16	14
6	2f	Fe ³⁺ -mont	20	17
7	2a	In ³⁺ -mont	45	69
8	2b	In ³⁺ -mont	58	89
9	2c	In ³⁺ -mont	34	52
10	2d	In ³⁺ -mont	34	52
11	2e	In ³⁺ -mont	23	35
12	2f	In ³⁺ -mont	27	42

Reaction conditions: **1** (6.0 mmol), **2** (3.0 mmol), Mⁿ⁺-mont (100 mg; Fe³⁺-mont, 0.0349 mmol, 0.012 mol amt, In³⁺-mont, 0.0197 mmol, 0.0066 mol amt), benzene (8 cm³), rt, 2 h. a) Mol of **3**/mol of acid sites.



Scheme 2.

Table 3. Acylation of **4** with **2a** or **2b** in the Presence of Mⁿ⁺-Mont

Entry	R	M ⁿ⁺ -mont	Isolated yield of 5 /%	TON ^a)
1	SiMe ₃	Fe ³⁺ -mont	54	46
2	Ph	Fe ³⁺ -mont	64	55
3	Bu	Fe ³⁺ -mont	36	31
4	Me	Fe ³⁺ -mont	9	8
5	SiMe ₃	In ³⁺ -mont	31	48
6	Ph	In ³⁺ -mont	51	78
7	Bu	In ³⁺ -mont	31	48
8	Me	In ³⁺ -mont	9	14

Reaction conditions: **4** (6.0 mmol), **2b** (3.0 mmol), Mⁿ⁺-mont (100 mg; Fe³⁺-mont, 0.0349 mmol, 0.012 mol amt, In³⁺-mont, 0.020 mmol, 0.0066 mol amt), benzene (8 cm³), rt, 2 h. a) Mol of **5**/mol of acid sites.

We have applied this reaction to many other substrates, such as cyclohexene, α -pinene, norbornene, trimethyl(vinyl)silane, and allyltrimethylsilane, but all reactions resulted in a formation of many unidentified products unfortunately. The treatment of 2-phenylpropene produced only the corresponding dimer.^{2c}

Experimental

General Procedures. All commercially available organic and inorganic compounds were used without further purification.

In³⁺-mont was newly prepared from Kunipia[®] G (<100 mesh, 6 g) and In(NO₃)₃·3H₂O (21.3 g, 60 mmol).^{2a} The amount of acid sites (Brønsted and Lewis acid sites) was estimated by the temperature-programmed desorption of ammonia gas (NH₃-TPD) analysis to be 0.195 mmol g⁻¹. The basal spacing (*d*₀₀₁) of In³⁺-mont was estimated by a sharp peak obtained on XRD analysis to be 15.4 Å, showing that the compound has an interlayer structure.

The acyl compounds **3a–3f** and **5a–5d** are known and were characterized by their spectral data after their isolation.^{5–13}

General Procedure for Mⁿ⁺-Mont-Catalyzed Friedel–Crafts Acylation of 1-Methyl-1-cyclohexene (1**) with Acyl Chloride **2**.** To a mixture of acyl chloride **2** (3 mmol) and benzene (8 cm³) was added Fe³⁺-mont (100 mg, 0.0349 mmol as acid sites estimated by NH₃-TPD) in one portion at room temperature with magnetic stirring. After the mixture was stirred at the temperature for a few minutes, 1-methyl-1-cyclohexene (**1**) (0.71 mL, 6 mmol) was added to it and the mixture was stirred vigorously at room temperature for 2 h. The catalyst was filtered off and rinsed with Et₂O (50 cm³). The solvent in a mixture of the filtrate and the ethereal washings was removed under the reduced pressure to leave an oil, which was subjected to silica gel column chromatography (eluent, hexane:diethyl ether = 10:1) to give the corresponding 6-acyl-1-methyl-1-cyclohexene **3**.

References

- a) M. Balogh and P. Laszlo, "Organic Chemistry Using Clays," Springer-Verlag, New York (1993). b) R. L. Augustine, "Heterogeneous Catalysis for the Synthetic Chemist," Marcel Dekker, New York (1996). c) R. A. Sheldon and R. S. Downing, *Appl. Catal.*, **A**, **189**, 163 (1999). d) J. H. Clark, *Acc. Chem. Res.*, **35**, 791 (2002).
- a) J. Tateiwa, H. Horiuchi, K. Hashimoto, T. Yamauchi, and S. Uemura, *J. Org. Chem.*, **59**, 5901 (1994). b) J. Tateiwa, T. Nishimura, H. Horiuchi, and S. Uemura, *J. Chem. Soc., Perkin Trans. 1*, **1994**, 3367. c) J. Tateiwa, E. Hayama, T. Nishimura, and S. Uemura, *Chem. Lett.*, **1996**, 59. d) J. Tateiwa, A. Kimura, K. Hashimoto, T. Yamauchi, and S. Uemura, *Bull. Chem. Soc. Jpn.*, **69**, 2361 (1996). e) J. Tateiwa, A. Kimura, M. Takasuka, and S. Uemura, *J. Chem. Soc., Perkin Trans. 1*, **1997**, 2169.
- a) T. Nishimura, N. Kakiuchi, M. Inoue, and S. Uemura, *Chem. Commun.*, **2000**, 1245. b) N. Kakiuchi, T. Nishimura, M. Inoue, and S. Uemura, *Bull. Chem. Soc. Jpn.*, **74**, 165 (2001).
- a) J. H. Clark, A. P. Kybett, D. J. Macquarrie, S. J. Barlow, and P. Landon, *J. Chem. Soc., Chem. Commun.*, **1989**, 1353. b) A. Cornélis, C. Dony, P. Laszlo, and K. M. Nsunda, *Tetrahedron Lett.*, **32**, 1423 (1991). c) A. Cornélis, C. Dony, P. Laszlo, and K. M. Nsunda, *Tetrahedron Lett.*, **32**, 2901 (1991). d) M. Davister and P. Laszlo, *Tetrahedron Lett.*, **34**, 533 (1993). e) O. Sieskind and P. Albrecht, *Tetrahedron Lett.*, **34**, 1197 (1993). f) L.-J. Li, B. Lu, T.-S. Li, and J.-T. Li, *Synth. Commun.*, **28**, 1439 (1998).
- J. K. Groves and N. Jones, *J. Chem. Soc. C*, **1968**, 2215.
- R. Jacquier, M. Mousseron, and S. Boyer, *Bull. Soc. Chim. Fr.*, **1956**, 1653.
- P. Beak and K. R. Berger, *J. Am. Chem. Soc.*, **102**, 3848 (1980).
- T. Shono, I. Nishiguchi, M. Sasaki, H. Ikeda, and M. Kurita, *J. Org. Chem.*, **48**, 2503 (1983).
- E. A. Braude and C. J. Timmons, *J. Chem. Soc.*, **1955**, 3766.
- B. B. Snider and A. C. Jackson, *J. Org. Chem.*, **47**, 5393 (1982).
- C. H. Wong and C. W. Bradshaw, U. S. Patent 5225339 (1993); *Chem. Abstr.*, **119**, 115519u (1993).
- N. Rosas, P. Sharma, C. Alvarez, A. Cabrera, R. Ramirez, A. Delgado, and H. Arzoumanian, *J. Chem. Soc., Perkin Trans. 1*, **2001**, 2341.
- A. V. Kel'in, A. W. Sromek, and V. Gevorgyan, *J. Am. Chem. Soc.*, **123**, 2074 (2001).