

Remarkable Substituent Effects on the Oxidizing Ability of Tetraarylismuthonium Tetrafluoroborates in Alcohol Oxidation

Yoshihiro Matano,^{*1} Takeshi Suzuki,¹ Takaharu Iwata,¹
Tomonori Shinokura,¹ and Hiroshi Imahori^{1,2,3}

¹Department of Molecular Engineering, Graduate School of Engineering, Kyoto University, Nishikyo-ku, Kyoto 615-8510

²Institute for Integrated Cell-Material Sciences (iCeMS), Kyoto University, Nishikyo-ku, Kyoto 615-8510

³Fukui Institute for Fundamental Chemistry, Kyoto University, Takano-Nishihiraki-cho, Sakyo-ku, Kyoto 606-8103


Received June 16, 2008; E-mail: matano@scl.kyoto-u.ac.jp

Substituent effects on the oxidizing ability of tetraarylismuthonium tetrafluoroborates in alcohol oxidation are reported. Intermolecular and intramolecular competition experiments on geraniol oxidation by the combined use of tetraarylismuthonium tetrafluoroborates and *N,N,N',N'*-tetramethylguanidine (TMG) have revealed that the oxidizing ability of the bismuthonium salt increases by the introduction of methyl groups at the ortho position and an electron-withdrawing group at the para position of the aryl ligands. The intermolecular and intramolecular H/D kinetic isotope effects observed for the competitive oxidation of *p*-bromobenzyl alcohols have shown that the present oxidation reaction consists of fast pre-equilibrium leading to alkoxytetraarylismuth(V) intermediates (first step) and α -hydrogen abstraction by the aryl ligand attached to the bismuth (second step). The experimental results demonstrate that the electron-deficient aryl ligands enhance the electrophilicity at the bismuth center to put forward the first step and that the bulky ligands destabilize the alkoxybismuth(V) intermediates to accelerate the second step. The newly explored mesityl- and 2,6-xylyltriarylismuthonium salts have proven to convert primary and secondary alcohols to the corresponding carbonyl compounds with high efficiency under mild conditions. A remarkable steric effect of these oxidants has also been exhibited in the chemoselective oxidation between primary and secondary benzylic alcohols.

It is well known that organobismuth(V) compounds possess inherent oxidizing ability derived from the facile Bi^V/Bi^{III} redox process.¹ For instance, triarylismuth(V) compounds of the type Ar₃BiX₂ (X = anionic ligands) convert a variety of primary and secondary alcohols to aldehydes and ketones, respectively, under mild conditions.² In the 80s, Barton and co-workers extensively studied alcohol oxidation using triarylismuth(V) compounds from synthetic and mechanistic points of view, and disclosed that the oxidation of simple alcohols proceeds through two main steps; formation of alkoxytriarylismuth(V) intermediates (first step) and α -hydrogen abstraction by an aryl ligand attached to the bismuth center (second step).³ In the second step, carbonyl compounds are formed together with equimolar amounts of arene and diarylbismuth(III) species. This mechanism implies that the oxidizing ability of triarylismuth(V) compounds strongly depends on the intrinsic character of the Bi-bound aryl ligands. In fact, Barton and co-workers reported some results of competition experiments on the oxidation of allylic alcohols by the Ar₃BiX₂/K₂CO₃ (X = Cl, Br, and ONO₂) system, in which the relative reactivity of Ar₃BiX₂ was estimated to be in the order Ar = 4-methylphenyl < phenyl < 4-chlorophenyl < 3-nitrophenyl.⁴ Recently, we reported the first systematic study of the substituent effects on the oxidizing ability of triarylismuth dichlorides (Ar₃BiCl₂) in alcohol oxidation, where the relative reactivity was determined quantitatively by intermolecular and intramolecular competitive experiments.⁵ It has been disclosed that

aryl ligands bearing a methyl group at the ortho position and/or an electron-withdrawing group at the para position dramatically enhance the overall rate of oxidation in the Ar₃BiCl₂/base system. Hence, to design efficient organobismuth(V) oxidants, both the steric and electronic nature of the aryl ligands must be taken into consideration.

In contrast to triarylismuth(V) compounds, only limited information is available for the oxidizing ability of other polyarylismuth(V) compounds. In 1987, Barton and co-workers reported that tetraphenylbismuth tosylate and trifluoroacetate (Ph₄BiX; X = OTs and OCOCF₃) oxidized 3 β -cholestanol and neopentyl alcohol under basic conditions.⁶ Their pioneering work, however, has received little attention, and there has been no systematic study of the substituent effects on the oxidizing ability of tetraarylismuth(V) compounds. Quite recently, we reported a new organobismuth(V) oxidant, mesityl-triphenylbismuthonium tetrafluoroborate (mesityl = 2,4,6-trimethylphenyl), which efficiently converted a variety of primary and secondary alcohols to the corresponding carbonyl compounds in the presence of *N,N,N',N'*-tetramethylguanidine (TMG) or 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU).⁷ Most importantly, the mesitylbismuthonium salt was found to oxidize the alcohols much more rapidly than the 4-methylphenyl- and phenyl-substituted bismuthonium salts. Such a remarkable rate enhancement can be explained by considering steric effects of the aryl ligands attached to the bismuth center; the bulky mesityl ligand is likely to accelerate the α -hydrogen ab-



Scheme 1. Synthesis of tetraarylbismuthonium tetrafluoroborates **1a–1i**.

straction from sterically crowded alkoxybismuth(V) intermediates. These results suggested that more efficient tetraarylbismuth(V) oxidants could be explored by suitably tuning the steric and electronic characters of the aryl ligands. Thus, we set out to systematically investigate the substituent effects on the intrinsic oxidizing ability of tetraarylbismuthonium salts.

Here we report the first systematic study of intermolecular and intramolecular competition experiments on alcohol oxidation by using a series of tetraarylbismuthonium tetrafluoroborates. It has been revealed that both steric and electronic effects of the aryl substituents definitely affect the overall rate of the reaction. Oxidation of several alcohols and chemoselective oxidation between primary and secondary alcohols with newly developed tetraarylbismuth(V) oxidants are also reported.

Results and Discussion

To investigate the relative oxidizing abilities of tetraarylbismuthonium salts, we prepared a series of tetraarylbismuthonium tetrafluoroborates **1a–1i** according to a reported procedure:⁸ reaction of triarylbismuth difluorides with arylboronic acids in the presence of $\text{BF}_3 \cdot \text{OEt}_2$, followed by treatment with an aqueous solution of NaBF_4 afforded **1a–1i** in high yield (Scheme 1). The bismuthonium salts **1a–1i** were purified by recrystallization from $\text{CH}_2\text{Cl}_2\text{--Et}_2\text{O}$ and characterized by spectroscopic methods. In the MS spectra of **1a–1i**, intense peaks assignable to $[\text{Ar}^1_3\text{Ar}^2\text{Bi}]^+$ ions were detected, while in the IR spectra, strong absorptions due to BF_4^- ion were observed at $1150\text{--}950\text{ cm}^{-1}$. These data indicate that each bismuth center in **1a–1i** essentially possesses onium nature. The structure of **1g** was further elucidated by X-ray crystallography. As shown in Figure 1, the bismuth center adopts a distorted tetrahedral geometry with C–Bi–C bond angles of $97.35(14)\text{--}125.26(14)^\circ$ and Bi–C bond distances of $2.207(4)\text{--}2.212(4)\text{ \AA}$. The Bi–F(1) distance [$3.005(2)\text{ \AA}$] is much longer than the sum of their covalent radii (2.17 \AA), implying that the cationic bismuth center weakly interacts with the fluoroborate ion.

Scheme 2 depicts a plausible mechanism of the alcohol oxidation by tetraarylbismuthonium salt **1**. On the basis of the fact that equimolar amounts of arene and triarylbismuthane are produced together with carbonyl compound, the reaction is considered to proceed through two main steps; formation of alkoxytetraarylbismuth(V) intermediate **A** (first step) and production of carbonyl compound via α -hydrogen abstraction (second step). The first step is promoted by the addition





Figure 1. ORTEP diagram of **1g**. Selected bond lengths (\AA) and angles ($^\circ$): Bi–C(1), $2.207(4)$; Bi–C(10), $2.208(4)$; Bi–C(16), $2.202(4)$; Bi–C(22), $2.212(4)$; Bi–F(1), $3.005(2)$; C(1)–Bi–C(10), $102.13(13)$; C(1)–Bi–C(16), $125.26(14)$; C(1)–Bi–C(22), $112.88(14)$; C(10)–Bi–C(16), $97.35(14)$; C(10)–Bi–C(22), $107.33(14)$; C(16)–Bi–C(22), $108.94(14)$.



Scheme 2. Plausible mechanism for alcohol oxidation.


Table 1. Oxidation of Benzyl Alcohol with **1g** in CDCl_3 ^{a)}

$\text{Ph-CH}_2\text{OH} \xrightarrow[\text{CDCl}_3, \text{ r.t.}]{\mathbf{1g}, \text{ base}} \text{Ph-CHO}$			
Entry	Base	Time/h	Yield ^{b)} /%
1	TMG	1.5	>99
2	DBU	1.5	>99
3	Et_3N	20	31
4	K_2CO_3	1.5	66
5	Pyridine	4	0

a) Benzyl alcohol (0.050 mmol), **1g** (0.055 mmol), base (0.055 mmol), CDCl_3 (1 mL). b) Determined by $^1\text{H NMR}$ using DMF as an internal standard.

of a base. As shown in Table 1, TMG and 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) were found to promote the oxidation of benzyl alcohol with mesityltriphenylbismuthonium salt **1g** efficiently, whereas Et_3N and K_2CO_3 were less effective. No oxidation took place with pyridine even after 4 h in CDCl_3 at room temperature. Therefore, TMG was used as the base in the present competition experiments.

First, we determined the relative oxidizing ability of tetraarylbismuthonium salts by means of intermolecular competitive oxidation reactions using tetraphenylbismuthonium salt **1a** and ortho/para-substituted derivatives **1b**, **1c**, and **1f–1i** (Scheme 3 and Tables 2 and 3). TMG (1.1 equiv) was added to a CH_2Cl_2 solution containing geraniol **2** ($5 \times 10^{-3}\text{ M}$) ($1\text{ M} = 1\text{ mol dm}^{-3}$) and two kinds of bismuthonium salts **1x**



Scheme 3. Competitive intermolecular oxidation of geraniol.

Table 2. Competitive Intermolecular Oxidation of **2** with **1**

Entry	1x	Ar ¹	1y	Ar ²	Ar ¹ H/Ar ² H ^{a)}
1	1a	C ₆ H ₅	1b	4-MeC ₆ H ₄	89/11
2	1a	C ₆ H ₅	1c	4-ClC ₆ H ₄	2/98
3	1a	C ₆ H ₅	1g	2,4,6-Me ₃ C ₆ H ₂	4/96
4	1c	4-ClC ₆ H ₄	1i	2,4,6-Me ₃ C ₆ H ₂	32/68
5	1f	2-MeC ₆ H ₄	1g	2,4,6-Me ₃ C ₆ H ₂	9/91
6	1g	2,4,6-Me ₃ C ₆ H ₂	1h	2,6-Me ₂ C ₆ H ₃	33/67

a) Determined by GC.


and **1y** (each 3 equiv) at 24 ± 1 °C. After **2** had been completely oxidized to geranial (**3**), the reaction mixture was analyzed by gas chromatography (GC) and ¹H NMR spectroscopy. Table 2 summarizes the relative ratios of two kinds of arenes produced (Ar¹H/Ar²H), which are indices of the relative overall reaction rates. The quantitative formation of the corresponding triarylbismuthanes was confirmed by ¹H NMR spectroscopy.⁹ It should be noted that, in the oxidation using **1g–1i**, the mesityl and 2,6-xylyl ligands are cleaved with ≥99% selectivity (vide infra). In Entry 4, the amount of chlorobenzene produced from **1i** is negligible as compared to that produced from **1c**, and, in Entries 5 and 6, only a trace amount (<1%) of benzene could be detected. Therefore, the Ar¹H/Ar²H ratios listed in Table 2 basically reflect the relative oxidizing ability of the bismuthonium salts examined.

On the basis of the Ar¹H/Ar²H ratios observed for the competitive reactions using **1a–1c**, the electronic effects of para-substituted aryl ligands relative to phenyl were determined as follows: 4-MeC₆H₄ (0.12) < C₆H₅ (1.0) < 4-ClC₆H₄ (49). Apparently, the overall reaction rate increases with increasing electron-withdrawing ability of the para substituents. Similarly, on the basis of the Ar¹H/Ar²H ratios observed for the competitive reactions using **1a** and **1f–1h** the steric effects of ortho-substituted aryl ligands relative to phenyl were deter-

Table 3. Substituent Effects Obtained from the Competition Experiments

Aryl ligand	Inter ^{a)}	Intra ^{b)}
4-MeC ₆ H ₄	0.12	0.57
C ₆ H ₅	1	1
4-ClC ₆ H ₄	49	1.4
2-MeC ₆ H ₄	2.4	22
2,6-Me ₂ C ₆ H ₃	48	3.3 × 10 ²
2,4,6-Me ₃ C ₆ H ₂	24	3.7 × 10 ²

a) Relative reactivity of **1b**, **1c**, and **1f–1h** vs. **1a**, obtained from the results shown in Table 2. b) Relative hydrogen-abstraction aptitude of aryl group vs. phenyl group, obtained from the results shown in Table 4.



Scheme 4. Competitive intramolecular oxidation of geraniol.

Table 4. Competitive Intramolecular Oxidation of **2** with **1**








Entry	1	Ar ¹	Ar ²	Ar ¹ H/Ar ² H ^{a)}
1	1d	C ₆ H ₅	4-MeC ₆ H ₄	84/16
2	1e	C ₆ H ₅	4-ClC ₆ H ₄	68/32
3	1f	C ₆ H ₅	2-MeC ₆ H ₄	12/88
4	1g	C ₆ H ₅	2,4,6-Me ₃ C ₆ H ₂	0.8/99.2
5	1h	C ₆ H ₅	2,6-Me ₂ C ₆ H ₃	0.9/99.1
6	1i	4-ClC ₆ H ₄	2,4,6-Me ₃ C ₆ H ₂	1.4/98.6

a) Determined by GC.

mined as follows: Ph (1.0) < 2-MeC₆H₄ (2.1) < 2,4,6-Me₃C₆H₂ (24) < 2,6-Me₂C₆H₃ (48).¹⁰ This result exhibits that the overall reaction rate is enhanced considerably by introducing two methyl groups at the ortho (2,6-) positions. The fact that **1h** oxidizes **2** more rapidly than does **1g** is also attributable to the difference in the electronic effect between mesityl and 2,6-xylyl ligands. In addition, the oxidizing ability of **1i** was found to be higher than that of **1c**, suggesting that the steric effect induced by one mesityl ligand is slightly larger than the electronic effect induced by one chlorophenyl ligand.

To evaluate the substituent effects on the second step, we next examined intramolecular competitive oxidation reactions using tetraarylbismuthonium salts bearing two kinds of aryl groups ([Ar¹₃Ar²Bi]⁺[BF₄[−]]; **1d–1i**). This provides relative rate constants at which the Bi–C bond is cleaved via α-hydrogen abstraction. Treatment of **1d–1i** with **2** in the presence of TMG in CH₂Cl₂ at 24 ± 1 °C gave two kinds of arenes and triarylbismuthanes (Scheme 4 and Table 4). On the basis of the relative ratios of the arenes (Ar¹H/Ar²H) and the number of the aryl ligands originally attached to bismuth, the relative hydrogen-abstraction aptitudes were determined as follows: 4-MeC₆H₄ (0.57) < C₆H₅ (1.0) < 4-ClC₆H₄ (1.4) < 2-MeC₆H₄ (22) < 2,6-Me₂C₆H₃ (3.3 × 10²) < 2,4,6-Me₃C₆H₂ (3.7 × 10²).¹¹ Obviously, the electronic effect of the para substituents is small as compared to that observed for the intermolecular competition experiments, suggesting that the α-hydrogen is abstracted via a weakly polar transition state. In sharp contrast, the steric effect on the second step is significant, as

Table 5. Oxidation of Alcohols with **1g–1i**

$\text{R}-\text{CH}(\text{R}')-\text{OH} \xrightarrow[\text{solvent, r.t.}]{\text{1g-i (1.1 equiv), TMG (1.1 equiv)}} \text{R}-\text{C}(=\text{O})-\text{R}'$					
Entry	Alcohol	Solvent	Time/h, Yield/%		
			1g	1h	1i
1		CH ₂ Cl ₂	1.0, 99 ^{a)}	0.5, 95 ^{a)}	0.5, 92 ^{a)}
2		CH ₂ Cl ₂	0.5, 98 ^{a)}	0.5, 99 ^{a)}	0.5, 99 ^{a)}
3		CH ₂ Cl ₂	4.0, 91 ^{a)}	0.5, 96 ^{a)}	0.5, 88 ^{a)}
4		CDCl ₃	40, 88 ^{b)}	1.0, 87 ^{b)}	1.0, 99 ^{b)}
5		CDCl ₃	1.5, 94 ^{b)}	1.0, 99 ^{b)}	0.5, 99 ^{b)}
6		Toluene- <i>d</i> ₈	1.5, 99 ^{b)}	0.5, 88 ^{b)}	0.5, 97 ^{b)}
7		Toluene- <i>d</i> ₈	3.0, 98 ^{b)}	1.5, 88 ^{b)}	0.5, 97 ^{b)}

a) Isolated yield. b) NMR yield.

Table 6. Chemoselective Oxidation of Primary/Secondary Alcohols

$\text{Ph}-\text{CH}_2\text{OH} + \text{Ph}-\text{CH}(\text{R})\text{OH} \xrightarrow[\text{CDCl}_3, 24 \pm 1^\circ\text{C}]{\text{oxidant (0.9 equiv), TMG (1.0 equiv)}} \text{Ph}-\text{CHO} + \text{Ph}-\text{C}(=\text{O})\text{R}$				
Entry	Oxidant	R	Time/h	PhCHO/PhCOR ^{a)}
1	1g	Me	1	81/19
2	1g	Et	1	92/8
3	1g	<i>i</i> -Pr	1	94/6
4	1d	Et	7	74/26
5	1f	Et	7	77/23
6	1h	Et	0.5	92/8
7	1i	Et	0.5	91/9
8	(2-MeC ₆ H ₄) ₃ BiCl ₂	Et	3	63/37
9	DMP ^{b)}	Et	2	67/33

a) Determined by ¹HNMR. b) TMG was not added.

ically. Note that the 1°/2° selectivities attained by tetraaryl-bismuthonium salts **1g–1i** are considerably higher than those observed for tris(2-methylphenyl)bismuth dichloride (63/37) and DMP (67/33) (Entries 8 and 9). Thus, **1g–1i** discriminate between steric bulkiness of the α-substituents (H vs. Et) of the alcohols more efficiently than other oxidants examined (Scheme 5). Although there exist more primary-selective oxidants,²⁰ tetraaryl-bismuthonium salts bearing bulky aryl ligands constitute a useful addition to the existing efficient oxidants.


Conclusion

In summary, we have elucidated the substituent effects on the oxidizing ability of tetraaryl-bismuthonium salts in alcohol

oxidation by means of intermolecular and intramolecular competition experiments. It is now clear that highly efficient Bi^V oxidants of the type [Ar¹₃Ar²Bi⁺][BF₄[−]] can be constructed by the introduction of bulky aryl ligands and/or electron-deficient aryl ligands. The bulky aryl ligands seem to weaken the Bi–C bond and dramatically accelerate the α-hydrogen abstraction step, whereas the electron-deficient aryl ligands enhance the electrophilicity of the bismuth center and put forward the pre-equilibrium leading to alkoxybismuth(V) intermediates. The newly explored mesityl- and 2,6-xylyl-triaryl-bismuthonium tetrafluoroborates have proven to convert primary and secondary alcohols to the corresponding carbonyl compounds in the presence of TMG with high efficiency under mild conditions. A remarkable steric effect of these oxidants has also been exhibited in the chemoselective oxidation between primary and secondary benzylic alcohols.

Experimental

General. All melting points were determined on a Yanagimoto hot-stage apparatus and are uncorrected. ¹HNMR spectra were recorded on a JEOL EX-400 (400 MHz) or a JEOL AL300 (300 MHz) spectrometer using CDCl₃ as the solvent unless otherwise noted. Chemical shifts are reported as relative values vs. tetramethylsilane. IR spectra were observed as KBr pellets on a Shimadzu FTIR-8200A spectrophotometer. MALDI-TOF mass spectra were measured on a Shimadzu MALDI-I spectrometer using CHCA as a matrix. GC was performed on a Shimadzu GC-14A fitted with a silica gel capillary column. Elemental analyses were performed at the Microanalytical Laboratory of Kyoto University. Dichloromethane and toluene were distilled from calcium hydride before use. Crystallographic data of **1g** was reported previously⁷ and has been deposited with Cambridge Crystallo-



Scheme 5. Chemoselective oxidation of benzyl alcohol and 1-phenyl-1-propanol by **1g–1i**.

graphic Data Centre: Deposition number CCDC-635005. Copies of the data can be obtained free of charge via <http://www.ccdc.cam.ac.uk/conts/retrieving.html> (or from the Crystallographic Data Centre, 12, Union Road, Cambridge, CB2 1EZ, UK; Fax: +44 1223 336033; e-mail: deposit@ccdc.cam.ac.uk).

Compounds. Tetraarylbismuthonium salts **1a–1i** were prepared from triarylbismuth difluorides, $\text{BF}_3 \cdot \text{OEt}_2$, and arylboronic acids as described below.⁸ *p*-Bromobenzyl alcohol- α - d_2^{21} and *p*-bromobenzyl alcohol- α , α - d_2^{22} were prepared by reduction of *p*-bromobenzaldehyde and *p*-bromobenzoic acid ethyl ester, respectively, with LiAlD_4 . TMG was purified by distillation. Other reagents were used as commercially received.

Tetraarylbismuthonium Tetrafluoroborates 1. Typical Procedure: $\text{BF}_3 \cdot \text{OEt}_2$ (0.78 mL, 6.0 mmol) was added to a CH_2Cl_2 solution (25 mL) containing triphenylbismuth difluoride (2.39 g, 5.00 mmol) and mesitylboronic acid (0.90 g, 5.5 mmol) at 0°C , and the mixture was stirred for 2 h at room temperature. An aqueous solution of NaBF_4 (17 g, 150 mmol) was then added, and the resulting two-phase solution was vigorously stirred for 20 min. The aqueous phase was separated and extracted with CH_2Cl_2 . The combined organic extracts were dried over MgSO_4 , passed through a short silica gel column, and evaporated under reduced pressure to leave a solid residue, which was reprecipitated from CH_2Cl_2 – Et_2O to afford **1g** (3.17 g, 98%) as a colorless solid. $^1\text{H NMR}$ (CDCl_3): δ 2.33 (s, 6H), 2.36 (s, 3H), 7.17 (s, 2H), 7.52–7.70 (m, 9H), 7.80–7.90 (m, 6H). Other bismuthonium salts were similarly prepared in 88–99% isolated yields. The spectral and analytical data of the newly prepared bismuthonium salts **1c**, **1h**, and **1i** are summarized below, and those of **1a**, **1b**, and **1d–1g** were reported in Refs. 7 and 8.

Tetrakis(4-chlorophenyl)bismuthonium Tetrafluoroborate (1c): Mp 190 – 192°C ; $^1\text{H NMR}$ (CDCl_3): δ 7.64 (d, 8H, $J = 8.4$ Hz), 7.71 (d, 8H, $J = 8.4$ Hz); IR ν_{max} 1150–950 (BF_4^-); MS (MALDI-TOF) m/z 655 ($[\text{M} - \text{BF}_4]^+$). Found: C, 39.06; H, 2.21%. Calcd for $\text{C}_{24}\text{H}_{16}\text{BBiCl}_4\text{F}_4$: C, 38.85; H, 2.17%.

Triphenyl(2,6-xylyl)bismuthonium Tetrafluoroborate (1h): Mp 201 – 203°C ; $^1\text{H NMR}$ (CDCl_3): δ 2.38 (s, 6H), 7.32–7.40 (m, 3H), 7.56–7.64 (m, 9H), 7.86 (d, 6H, $J = 7.2$ Hz); IR ν_{max} 1150–950 (BF_4^-); MS (MALDI-TOF) m/z 545 ($[\text{M} - \text{BF}_4]^+$). Found: C, 49.24; H, 3.89%. Calcd for $\text{C}_{26}\text{H}_{24}\text{BBiF}_4$: C, 49.39; H, 3.83%.

Tris(4-chlorophenyl)mesitylbismuthonium Tetrafluoroborate (1i): Mp $>300^\circ\text{C}$; $^1\text{H NMR}$ (CDCl_3): δ 2.32 (s, 6H), 2.37 (s, 3H), 7.18 (s, 2H), 7.60 (d, 6H, $J = 8.6$ Hz), 7.77 (d, 6H, $J = 8.6$ Hz); IR ν_{max} 1150–950 (BF_4^-); MS (MALDI-TOF) m/z

661 ($[\text{M} - \text{BF}_4]^+$). Found: C, 43.03; H, 3.01%. Calcd for $\text{C}_{27}\text{H}_{23}\text{BBiCl}_3\text{F}_4$: C, 43.26; H, 3.09%.

Intermolecular Competition Reactions. TMG (0.020 mmol) was added to a mixture of two kinds of tetraarylbismuthonium salts **1x** and **1y** (0.060 mmol each), geraniol **2** (0.020 mmol), and CH_2Cl_2 (1.0 mL) at $24 \pm 1^\circ\text{C}$, and the resulting mixture was stirred at the same temperature. The ratio of arenes produced ($\text{Ar}^1\text{H}/\text{Ar}^2\text{H}$) was determined by GC analysis, and the reaction mixture was then concentrated under reduced pressure. The residue was taken up with CDCl_3 (ca. 2 mL) and the resulting CDCl_3 solution was measured by $^1\text{H NMR}$ spectroscopy. Each reaction was carried out three times and the average values of $\text{Ar}^1\text{H}/\text{Ar}^2\text{H}$ are listed in Table 2. In the competition reactions using **1f–1h**, a trace amount (less than 1%) of benzene was also formed.

Intramolecular Competition Reactions. TMG (0.020 mmol) was added to a mixture of **1d–1i** (0.020 mmol), **2** (0.020 mmol), and CH_2Cl_2 (1.0 mL) at $24 \pm 1^\circ\text{C}$. The reaction mixture was worked up and analyzed according to a procedure similar to that described above. Each reaction was carried out three times and the average values are listed in Table 4. The substituent effects listed in Table 3 were determined by taking the number of aryl ligands attached to the bismuth center of **1d–1i** into consideration. That is, the α -hydrogen-abstracting aptitudes listed in Table 3 were calculated based on the ratios of $[\text{Ar}^1\text{H}]/3[\text{Ar}^2\text{H}]$.

H/D Kinetic Isotope Experiments. TMG (0.010 mmol) was added to a mixture of **1g** (0.010 mmol), *p*- $\text{BrC}_6\text{H}_4\text{CH}_2\text{OH}$ (5.6 mg, 0.030 mmol), *p*- $\text{BrC}_6\text{H}_4\text{CD}_2\text{OH}$ (5.7 mg, 0.030 mmol), and CDCl_3 (1.0 mL) at $24 \pm 1^\circ\text{C}$. In the intramolecular competition experiments, *p*- $\text{BrC}_6\text{H}_4\text{CH}(\text{D})\text{OH}$ (1.9 mg, 0.010 mmol) was used as the substrate. After 30 min, the reaction mixture was analyzed by $^1\text{H NMR}$ spectroscopy, and the ratio of *p*- $\text{BrC}_6\text{H}_4\text{CHO}/p$ - $\text{BrC}_6\text{H}_4\text{CDO}$ was estimated by the comparison of the integral value of the formyl proton of *p*- $\text{BrC}_6\text{H}_4\text{CHO}$ with that of the aromatic protons of the two aldehydes. In these reactions, no other products derived from the alcohols were detected. Each reaction was carried out three times and the average values are shown in eqs 1 and 2.

Oxidation of Alcohols by 1g–1i/TMG. TMG (38 mg, 0.33 mmol) was added to a mixture of **1g–1i** (0.33 mmol), alcohol (0.30 mmol), and CH_2Cl_2 (10 mL), and the resulting mixture was stirred at room temperature. After the alcohol had been consumed (checked by TLC), the reaction mixture was concentrated under reduced pressure to leave an oily residue, which was then subjected on a short silica gel column (hexane–hexane/ EtOAc or CH_2Cl_2). The carbonyl compounds were easily separated from tri-

phenylbismuthane and isolated in a pure form. The yields of volatile carbonyl compounds were determined by ^1H NMR spectra of the reaction mixture.

Chemoselective Oxidation of Primary/Secondary Alcohols. TMG (0.055 mmol) was added to a mixture of two kinds of alcohols (each, 0.055 mmol), oxidant (0.05 mmol), and CDCl_3 (1 mL) at $24 \pm 1^\circ\text{C}$, and the resulting mixture was stirred at the same temperature. The relative ratio of two carbonyl compounds was determined by ^1H NMR spectra of the reaction mixture. Each reaction was carried out twice, and the average values are listed in Table 6.

This work was supported by a Grant-in-Aid (No. 17350018) from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

References

- 1 a) J. P. Kitchin, in *Organic Synthesis by Oxidation with Metal Compounds*, ed. by W. J. Mijs, C. R. H. I. de Jonge, Plenum, New York, **1986**, Chap. 15, pp. 817–837. b) M. Postel, E. Duñach, *Coord. Chem. Rev.* **1996**, *155*, 127. c) H. Suzuki, T. Ikegami, Y. Matano, *Synthesis* **1997**, 249. d) N. Komatsu, in *Organobismuth Chemistry*, ed. by H. Suzuki, Y. Matano, Elsevier, New York, **2001**, Chap. 5, pp. 371–440. e) N. M. Leonard, L. C. Wieland, R. S. Mohan, *Tetrahedron* **2002**, *58*, 8373. f) Y. Matano, in *Main Group Metals in Organic Synthesis*, ed. by H. Yamamoto, K. Oshima, Wiley-VCH, Weinheim, **2004**, Chap. 14, pp. 753–811. g) Y. Matano, in *Comprehensive Organometallic Chemistry III*, ed. by P. Knochel, Elsevier, Oxford, **2006**, Chap. 9.11, pp. 425–456.
- 2 a) F. Challenger, O. V. Richards, *J. Chem. Soc.* **1934**, 405. b) D. H. R. Barton, J. P. Kitchin, W. B. Motherwell, *J. Chem. Soc., Chem. Commun.* **1978**, 1099. c) D. H. R. Barton, D. J. Lester, W. B. Motherwell, M. T. B. Papoula, *J. Chem. Soc., Chem. Commun.* **1979**, 705. d) D. H. R. Barton, J. P. Kitchin, D. J. Lester, W. B. Motherwell, M. T. B. Papoula, *Tetrahedron* **1981**, *37*, Suppl. 1, 73. e) V. A. Dodonov, T. I. Zinov'eva, N. N. Osadchaya, *Zh. Obshch. Khim.* **1988**, *58*, 712. f) T. I. Zinov'eva, N. V. Dolganova, V. A. Dodonov, I. G. Prezhbog, *Izv. Akad. Nauk. Ser. Khim.* **1998**, 681. g) H. Suzuki, T. Ikegami, Y. Matano, *Tetrahedron Lett.* **1994**, *35*, 8197. h) Y. Matano, H. Nomura, *J. Am. Chem. Soc.* **2001**, *123*, 6443. i) Y. Matano, H. Nomura, H. Suzuki, M. Shiro, H. Nakano, *J. Am. Chem. Soc.* **2001**, *123*, 10954. j) Y. Mitsumoto, M. Nitta, *Bull. Chem. Soc. Jpn.* **2003**, *76*, 1029.
- 3 In the reaction with α -diols, the oxidative C–C bond cleavage takes place to give two equivalents of carbonyl compounds together with triarylbi-muthane.
- 4 The relative reactivity of Ar_3BiX_2 was determined from the relative ratios of triarylbi-muthanes (Ar_3Bi) recovered from the reaction mixture, but significant variation was observed for the yield of Ar_3Bi . Therefore, the relative rate values reported in Ref. 2d are not used here for quantitative discussion.
- 5 a) Y. Matano, H. Nomura, *Angew. Chem., Int. Ed.* **2002**, *41*, 3028. b) Y. Matano, T. Hisanaga, H. Yamada, S. Kusakabe, H. Nomura, H. Imahori, *J. Org. Chem.* **2004**, *69*, 8676.
- 6 In the presence of *N*-*t*-butyl-*N'*,*N'*,*N''*,*N''*-tetramethylguanidine, these alcohols were oxidized to the corresponding carbonyl compounds in 78–92% yields after 7.5–12 h. D. H. R. Barton, J.-P. Finet, W. B. Motherwell, C. Pichon, *J. Chem. Soc., Perkin Trans. I* **1987**, 251.
- 7 Y. Matano, T. Suzuki, T. Shinokura, H. Imahori, *Tetrahedron Lett.* **2007**, *48*, 2885.
- 8 a) Y. Matano, S. A. Begum, T. Miyamatsu, H. Suzuki, *Organometallics* **1998**, *17*, 4332. b) Y. Matano, S. A. Begum, T. Miyamatsu, H. Suzuki, *Organometallics* **1999**, *18*, 5668.
- 9 In most cases, the relative ratio of $\text{Ar}^1_3\text{Bi}/\text{Ar}^1_2\text{Ar}^2\text{Bi}$ could not be determined accurately due to overlapping of their aromatic protons.
- 10 Care must be taken to compare the results from the intermolecular competitive experiments of **1a–1c** ($[\text{Ar}_4\text{Bi}][\text{BF}_4]$) with those of **1f–1h** ($[\text{Ph}_3\text{ArBi}][\text{BF}_4]$), because the number of ortho/para-substituted aryl ligands attached to the bismuth center is different between these two types of oxidants.
- 11 The electronic effect observed for the present $[\text{Ar}^1_3\text{Ar}^2\text{Bi}][\text{BF}_4]/\text{TMG}$ system is somewhat larger than that reported for the $\text{Ar}^1_2\text{Ar}^2\text{BiCl}_2/\text{DBU}$ system, where the hydrogen-abstracting aptitudes are determined as follows: $4\text{-MeC}_6\text{H}_4$ (0.92) < C_6H_5 (1.0) < $4\text{-ClC}_6\text{H}_4$ (1.1) < $2\text{-MeC}_6\text{H}_4$ (1.8) (see, Ref. 5b). The observed electronic effect implies that the α -hydrogen might be abstracted as a hydride in a formal sense.
- 12 If the first step consists of a slow equilibrium compared to the second step, it would become a rate-determining step. In such a case, a small secondary isotope effect should be observed for the intermolecular competition reactions. See: H. Simon, D. Palm, *Angew. Chem., Int. Ed. Engl.* **1966**, *5*, 920.
- 13 For example, see: a) W.-B. Chiao, W. H. Saunders, Jr., *J. Am. Chem. Soc.* **1978**, *100*, 2802. b) H. Kwart, M. Brechbiel, *J. Am. Chem. Soc.* **1981**, *103*, 4650.
- 14 The intramolecular kinetic isotope effect ($k_{\text{H/D}}$) reported for the Swern oxidation of PhCH(D)OH was 2.6 ± 0.3 . See: M. Marx, T. T. Tidwell, *J. Org. Chem.* **1984**, *49*, 788.
- 15 Finet and co-workers reported the influence of ortho substituents of triarylbi-muth derivatives in base-catalyzed C-arylation reactions with phenols and enols. In their ligand coupling systems, the steric hindrance plays an inhibitory role, which is more pronounced in the mesityl ligand. See: A. Fedorov, S. Combes, J.-P. Finet, *Tetrahedron* **1999**, *55*, 1341.
- 16 We reported the electronic effect of aryl ligands of unsymmetrical tetraarylbi-muthonium salts in base-catalyzed C-arylation reaction of 2-naphthol, where the migratory aptitude decreases with increasing the electron-donating ability of the para substituents. See: Y. Matano, T. Miyamatsu, H. Suzuki, *Chem. Lett.* **1998**, 127.
- 17 It should be noted that the equatorial–apical equilibrium of the aryl ligands might affect the hydrogen-abstracting ability. Presumably, the bulky aryl ligands are apt to occupy the equatorial positions relative to the Bi-bound alkoxy group. We appreciate the reviewer's valuable comments.
- 18 Trifluoromethyl carbinols are known to resist oxidation and only a few methods are available for their conversion to the corresponding trifluoromethyl ketones. For example, see: a) R. J. Linderman, D. M. Graves, *J. Org. Chem.* **1989**, *54*, 661. b) I. E. Markó, P. R. Giles, M. Tsukazaki, I. Chellé-Regnaut, A. Gautier, S. M. Brown, C. J. Urch, *J. Org. Chem.* **1999**, *64*, 2433. c) V. Kesavan, D. Bonnet-Delpon, J.-P. Bégue, A. Srikanth, S. Chandrasekaran, *Tetrahedron Lett.* **2000**, *41*, 3327.
- 19 4-Methoxybenzyl alcohol, 4-bromobenzyl alcohol, 4-(methylthio)benzyl alcohol, 2-methylbenzyl alcohol, cinnamyl alcohol, benzhydrol, and 1-dodecanol were oxidized by **1g**/TMG in CH_2Cl_2 or toluene to give the corresponding carbonyl compounds in 94–99% yield. See Ref. 7.
- 20 For example, see: a) $\text{RuCl}_2(\text{PPh}_3)_3$: H. Tomioka, K. Takai,

- K. Oshima, H. Nozaki, *Tetrahedron Lett.* **1981**, 22, 1605. b) Electrooxidation mediated by 2,2,6,6-tetramethylpiperidine-1-oxyl (TEMPO): M. F. Semmelhack, C. S. Chou, D. A. Cortes, *J. Am. Chem. Soc.* **1983**, 105, 4492. c) 4-(Benzoyloxy)-TEMPO/NaBrO₂: T. Inokuchi, S. Matsumoto, T. Nishiyama, S. Torii, *J. Org. Chem.* **1990**, 55, 462. d) TEMPO/NCS: J. Einhorn, C. Einhorn, F. Ratajczak, J.-L. Pierre, *J. Org. Chem.* **1996**, 61, 7452. e) TEMPO/trichloroisocyanuric acid: L. De Luca, G. Giacomelli, A. Porcheddu, *Org. Lett.* **2001**, 3, 3041. f) TEMPO/RuCl₂(PPh₃)₃: A. Dijkman, A. Marino-González, A. Mairata i Payeras, I. W. C. E. Arends, R. A. Sheldon, *J. Am. Chem. Soc.* **2001**, 123, 6826. g) 4-(Acetamido)-TEMPO/Cu(ClO₄)₂/DMAP/O₂/[bmim]PF₆: N. Jiang, A. J. Ragauskas, *Org. Lett.* **2005**, 7, 3689. h) TEMPO/Cu(NO₃)₂/Cu(NO₃)₂/O₂: M. Gilhespy, M. Lok, X. Baucherel, *Chem. Commun.* **2005**, 1085, and references therein.
- 21 E. J. Corey, J. O. Link, *Tetrahedron Lett.* **1989**, 30, 6275.
22 S. K. Pollack, B. C. Raine, W. J. Hehre, *J. Am. Chem. Soc.* **1981**, 103, 6308.