

The oncoprotein gankyrin binds to MDM2/HDM2, enhancing ubiquitylation and degradation of p53

Hiroaki Higashitsuji,¹ Hisako Higashitsuji,¹ Katsuhiko Itoh,¹ Toshiharu Sakurai,¹ Toshikazu Nagao,¹ Yasuhiko Sumitomo,¹ Tomoko Masuda,¹ Simon Dawson,² Yutaka Shimada,³ R. John Mayer,² and Jun Fujita^{1,*}

¹Department of Clinical Molecular Biology, Graduate School of Medicine, Kyoto University, 54 Shogoin Kawaharacho, Sakyo-ku, Kyoto 606-8507, Japan

²Laboratory of Intracellular Proteolysis, School of Biomedical Sciences, University of Nottingham Medical School, Queens Medical Centre, Nottingham NG7 2UH, United Kingdom

³Department of Surgery and Surgical Basic Science, Graduate School of Medicine, Kyoto University, 54 Shogoin Kawaharacho, Sakyo-ku, Kyoto 606-8507, Japan

*Correspondence: jfujita@virus.kyoto-u.ac.jp

In this article (Cancer Cell 8, pp. 75–86), the name of one of the authors was listed incorrectly. The author incorrectly listed as “Haruhiko Sumitomo” is actually named Yasuhiko Sumitomo.