Cancer epigenetics is no Mickey Mouse

Andrew P. Feinberg^{1,*}

¹Departments of Medicine, Molecular Biology & Genetics, and Oncology, Johns Hopkins University School of Medicine 1064 Ross, 720 Rutland Avenue, Baltimore, Maryland 21205

*Correspondence: afeinberg@jhu.edu

(Cancer Cell 8, 267-268; October 18, 2005)

Due to a production error, the first sentence of the fifth paragraph of the main text reads, "Second, both models of the associated cancers involve *hypomethylation* rather than *hypermethylation*." It should have read "Second, both models or the associated cancers involve *hypomethylation* rather than *hypermethylation*."

DOI: 10.1016/j.ccr.2005.10.011

Global loss of imprinting leads to widespread tumorigenesis in adult mice

Teresa M. Holm, ^{1,2} Laurie Jackson-Grusby, ^{1,3} Tobias Brambrink, ¹ Yasuhiro Yamada, ^{1,4} William M. Rideout III, ^{1,5} and Rudolf Jaenisch ^{1,2,*}

(Cancer Cell 8, 275-285; October 18, 2005)

The accession number for the microarray data in this article was omitted. The data can be found at http://www.ebi.ac.uk/arrayexpress/; the accession number is E-MEXP-450.

DOI: 10.1016/j.ccr.2005.10.006

¹ Whitehead Institute for Biomedical Research, Cambridge, Massachusetts 02142

²Massachusetts Institute of Technology, Cambridge, Massachusetts 02142

³ Present address: Pathology Department, Children's Hospital Boston and Harvard Medical School, 320 Longwood Avenue, Boston, Massachusetts 02115

⁴Present address: Department of Tumor Pathology, Gifu University, Gifu 501-1194, Japan

⁵Present address: AVEO Pharmaceuticals Inc., Cambridge, Massachusetts 02139

^{*}Correspondence: jaenisch@wi.mit.edu