ERβ Impedes Prostate Cancer EMT by Destabilizing HIF-1a and Inhibiting VEGF-Mediated Snail Nuclear **Localization: Implications for Gleason Grading**

Paul Mak, Irwin Leav, Bryan Pursell, Donggoo Bae, Xiaofang Yang, Cherie A. Taglienti, Lindsey M. Gouvin, Vishva M. Sharma, and Arthur M. Mercurio*

*Correspondence: arthur.mercurio@umassmed.edu DOI 10.1016/j.ccr.2010.05.005

(Cancer Cell 17, 319-332; April 13, 2010)

It has come to our attention that there was an error in the assembly of Figure 4C, resulting in the inclusion of an incorrect photomicrograph (ER\beta 1 Knockdown: +MG132). We have now inserted the correct photomicrograph in the revised Figure 4C. This error has no bearing on the conclusion of the study and we apologize for any confusion caused by the incorrect photomicrograph.




Figure 4. ER β 1 Destabilizes HIF-1 α Protein and Represses HIF-1-Mediated Transcription of VEGF-A

(C) PC3 cells (scrambled control cells) or ER_β1 knockdown cells were treated in the absence or presence of MG132 (1 uM) for 6 hr and photographed. Scale bars represent 50 um.