



Synthesis, characterization, and antifungal activity of novel inulin derivatives with chlorinated benzene

Zhanyong Guo^{a,*}, Qing Li^a, Gang Wang^a, Fang Dong^a, Haoyuan Zhou^{a,b}, Jing Zhang^c

^a Key Laboratory of Coastal Biology and Bioresource Utilization, Yantai Institute of Coastal Zone Research, Chinese Academy of Sciences, Yantai 264003, China

^b Graduate School of Chinese Academy of Sciences, Beijing 100039, China

^c Laishan Forestry Bureau, Yantai, China

ARTICLE INFO

Article history:

Received 9 June 2013

Received in revised form 15 July 2013

Accepted 18 August 2013

Available online 25 August 2013

Keywords:

Inulin derivatives

Antifungal activity

Chlorine

ABSTRACT

A group of novel inulin derivatives containing benzene or chlorinated benzene were synthesized by reaction of chloroacetyl inulin (**CAIL**) with the Schiff bases of 4-amino-pyridine, including (2-pyridyl)acetyl inulin chloride (**PAIL**), 2-[4-(2-chlorobenzylideneamino)-pyridyl]acetyl inulin chloride (**2CPAIL**), 2-[4-(4-chlorobenzylideneamino)-pyridyl]acetyl inulin chloride (**4CPAIL**), and 2-[4-(2,4-dichlorobenzylideneamino)-pyridyl]acetyl inulin chloride (**2,4DCPAIL**). Their antifungal activity against three kinds of phytopathogens was estimated by hypha measurement *in vitro*. Of all the synthesized chitosan derivatives, **2,4DCPAIL** inhibited the growth of the tested phytopathogens with inhibitory indices of 67%, 47%, and 43% against *Colletotrichum lagenarium* (Pass) Ell. et alst., *Phomopsis asparagi* (Sacc.) Bubak and *Fusarium oxysporum* (schl.) F.sp. *niveum* (*F. oxysporum*) respectively at 1.0 mg/mL. The results indicate that all the inulin derivatives have better antifungal activity than inulin, and the inhibitory index is affected by the chlorine atom grafted to the inulin derivatives.

© 2013 Elsevier Ltd. All rights reserved.

1. Introduction

As a kind of abundant natural polysaccharide, inulin is mainly extracted from such low-requirement crops as *Helianthus tuberosus*, chicory, and yacon. Inulin consists primarily of β -fructosyl fructose units – always presented in furanose form – usually with a glucopyranose unit reducing end (GFn) (Rogge & Stevens, 2004). Inulin has unique range of molecular weight with the degree of polymerization (DP) varies from 2 to 70, which is rare as a kind of polysaccharide. This polysaccharide has exhibited some interesting properties such as beneficial nutritional attributes for human health, moderate average degree of polymerization and readiness of being obtained (Beylot, 2006; Causey, Feirtag, Gallaher, Tungland, & Slavin, 2000). As a kind of source of renewable, biodegradable and environmentally benign polysaccharide, inulin is a promising candidate to meet the continuously increasing demand for biodegradable and environmentally benign polymeric material. This structural singularity raises its interesting properties like beneficial nutritional attributes, which in turn excites the chemical modification of inulin in recent years (Beylot, 2006; Habibi & Dufresne, 2008). We have reported the antioxidant activity


and antifungal activity of inulin derivatives (Ren, Liu, Dong, & Guo, 2011; Ren et al., 2012), which proved the chemical modification could improve the bioactivity of inulin.

Many reports have demonstrated that aromatic nucleus Schiff bases had good bioactivities (Karthikeyan et al., 2006). It is also reported that salicylaldehyde derivatives, with one or more halogens in the aromatic ring, showed a variety of biological activities (Li, Guo, & Jiang, 2010). Meanwhile, as antifungal group, chlorinated benzenes are used in many fungicides such as pentachloronitrobenzene and chlorothalonil which have five or four chlorine atoms respectively (Sha, 1994). Chemical fungicides with chlorine atoms have good fungicidal activity and used widely years ago, but their toxicities and residues give rise to serious problems for environment and human health. When these groups are grafted onto inulin, they should be released slowly and may meet the requirements of environmental safety.

In this paper, we reported the synthesis and antifungal properties of a group of inulin derivatives with benzene or chlorinated benzene as substituent including **PAIL**, **2CPAIL**, **4CPAIL**, and **2,4DCPAIL**. The chloroacetyl inulin (**CAIL**) was first synthesized by reaction between the C-6 hydroxyl of inulin and chloroacetyl chloride. **CAIL** was an excellent intermediate of the project as the chlorine of **CAIL** can easily attack pyridine to give *N*-alkylpyridinium salts (Li et al., 2013; Sajomsang, 2010). Subsequently the benzene or chlorinated benzene groups were introduced into inulin through the reaction between Schiff bases of 4-amino-pyridine and **CAIL**.

* Corresponding author. Tel.: +86 535 2109171; fax: +86 535 2109000.

E-mail addresses: zhanyongguo@gmail.com, zhanyongguo@hotmail.com (Z. Guo).



Scheme 1. Synthetic routes for inulin derivatives.

The aimed inulin derivatives designed in this way were expected to have advantageous features such as high antifungal activity and good water solubility. Apart from these advantages, the reaction generates Schiff bases functional groups, which is also helpful for the constructions of bioactive inulin derivatives (Guo, Ren, Dong, Wang, & Li, 2013). The chemical structures of the derivatives were characterized by FT-IR and ^1H NMR.

Plant pathogenic fungi negatively affect a large number of important vegetables and fruits. The losses caused by plant pathogenic fungi have attracted wide attention throughout the world. For example, *Phomopsis asparagi* can cause stem blight of asparagus and subsequent yield loss; the premature plant death caused by *Colletotrichum Lagerarium* can lead to a yield loss of 6–48% (Chen & Dai, 2012; Elena, 2006). Therefore, three common plants-threatening fungi, *C. Lagerarium*, *P. asparagi*, and *Fusarium oxysporum*, were selected to evaluate the antifungal properties of the derivatives by hypha measurement *in vitro*. Control of these plants-threatening fungi could benefit the production of related vegetables and fruits.

2. Experimental

2.1. Materials and analytical methods

Inulin was purchased from E. Merck (Darmstadt, Germany). 2-Chlorobenzaldehyde, 4-chlorobenzaldehyde, and 2,4-dichlorobenzaldehyde were purchased from the Sigma–Aldrich Chemical Corp. The other reagents are all of analytical grade and used without further purification. FT-IR spectra were recorded on a Jasco-4100 Fourier Transform Infrared Spectrometer (Japan, provided by JASCO Co., Ltd Shanghai, China) with KBr disks. ^1H nuclear magnetic resonance (^1H NMR) spectra were measured with a Bruker AVIII-500 Spectrometer (Switzerland, provided by Bruker Tech. and Serv. Co., Ltd. Beijing, China.). Three plants-threatening fungi *C. lagenarium* (Pass) Ell. et Halst., *P. asparagi* (Sacc.) Bubak and *F. oxysporum* (Schl.) F. sp. niveum (*F. oxysporum*) were provided by Qingdao Academy of Agricultural Sciences.

2.2. The synthesis of the Schiff bases of 4-amino-pyridine

As shown in Scheme 1, a solution of 0.01 mol aldehydes (2-chlorobenzaldehyde, 4-chlorobenzaldehyde or 2,4-dichlorobenzaldehyde) and 0.01 mol 4-amino-pyridine were refluxed in 30 mL ethanol for 4 h. Then the solvent was concentrated under reduced pressure. The crude products were purified by crystallization from ethanol. (Kaboudin, Afsharinezhad, & Mohammad, 2010).

2.3. The synthesis of chloroacetyl inulin

Chloroacetyl inulin (**CAIL**) was synthesized as follows (Li, Guo, & Jiang, 2010). 1.61 g inulin was dissolved in 100 mL H_2O at room temperature (r.t.), and 0.02 mol chloroacetyl chloride was added drop wise. After stirring for 12 h at r.t., the solution was concentrated under reduced pressure. The concentrated solution was precipitated by the addition of excess acetone and the precipitate was filtered. The products were washed with ether dried at 60°C for 6 h.

2.4. The synthesis of inulin derivatives

A solution of 0.3 g **CAIL** and 0.5 mL pyridine or the Schiff bases of 4-amino-pyridine in 20 mL *N,N*-dimethylformamide (DMF) was stirred for 24 h at 60°C . The solution was precipitated with excess acetone and the precipitate was filtered. The unreacted pyridine, amine, and other outgrowth were extracted in a Soxhlet apparatus with ethanol for two days. The inulin derivatives were obtained after drying at 60°C for 6 h (Li, Guo & Jiang, 2010).

2.5. Antifungal assays

Antifungal assays were performed by following the plate growth rate method (Jasso de Rodríguez, Hernández-Castillo, Rodríguez-García, & Angulo-Sánchez, 2005; Li, Guo & Jiang, 2010). Briefly, the compounds were dissolved in distilled water at a concentration of 5.0 mg/mL. Then, each sample (inulin, **CAIL** and inulin derivatives) solution was added to sterilized potato dextrose agar to give a final concentration of 0.1, 0.5, and 1.0 mg/mL. After the mixture was cooled in the plate (6.0 cm diameter), the mycelium of fungi of a 5.0 mm diameter was transferred to the test plate and incubated at 27°C . When the mycelium of fungi reached the edges of the control plate (without the presence of samples), the antifungal index was calculated as follows:

$$\text{Antifungal index (\%)} = 1 - \frac{D_a}{D_b} \times 100$$

where D_a is the diameter of the growth zone in the test plates and D_b is the diameter of the growth zone in the control plate. Each experiment was performed three times, and the data were shown with mean $\pm$ SD. The Scheffe method was used to evaluate the differences in antifungal index in the antifungal tests. Results with $P < 0.05$ were considered statistically significant (Jasso de Rodríguez et al., 2005).

3. Results and discussion

3.1. Structure of the inulin derivatives

Each step of the synthesis was followed by FT-IR and ^1H NMR spectroscopy. The FT-IR spectra of inulin, **CAIL**, **PAIL**, **2CPAIL**, **4CPAIL**, and **2,4DCPAIL** are shown in Fig. 1. The FT-IR spectrum of inulin shows peaks of saccharide at 848 cm^{-1} and 1029 cm^{-1} (Ren et al., 2012). As far as the FT-IR spectra of **CAIL** and inulin are concerned, the characteristic peak of carbonyl ($\text{C}=\text{O}$) vibration deformation appears at 1747 cm^{-1} . Another new peak at 771 cm^{-1} is assigned to the $\text{C}-\text{Cl}$ group which indicates that **CAIL** is synthesized successfully. After pyridine grafted onto **CAIL**, the absorbance of $\text{C}-\text{Cl}$ at 771 cm^{-1} is weakened and new peaks appear at 1488 and 1538 cm^{-1} which are assigned to the absorbance of pyridine in the spectrum of **PAIL** (Panicker et al., 2006; Li, Guo & Jiang, 2010). In the spectra of **2CPAIL**, **4CPAIL** and **2,4DCPAIL**, new peaks appear at about 1535 , 1465 , 830 cm^{-1} , which are attributed to the characteristic absorption of benzene and pyridine. Another prominent peak at about 1654 cm^{-1} was assigned to the $\text{C}=\text{N}$ of Schiff bases group.

Fig. 2 presented the ^1H NMR spectra of inulin, **CAIL**, **PAIL**, **2CPAIL**, **4CPAIL** and **2,4DCPAIL**. It is known that all the signals at $3.0\text{--}5.4\text{ ppm}$ are assigned to the protons of inulin (Ren et al., 2012). After reacted with chloroacetyl chloride, new signals appear at $1.1\text{--}1.3\text{ ppm}$ in **CAIL**, which are related to the proton of $-\text{CH}_2\text{Cl}$. New signals appear at $7.9\text{--}8.9\text{ ppm}$ (proton of pyridine) in **PAIL**, and corresponding signals including proton of benzene and pyridine move to high field at about $6.7\text{--}8.0\text{ ppm}$ in **2CPAIL**, **4CPAIL**, and **2,4DCPAIL**, which should be assigned to the conjugate structure. Above results further confirmed the success of the preparation of inulin derivatives.




Fig. 1. FT-IR spectra of inulin, **CAIL**, **PAIL**, **2CPAIL**, **4CPAIL**, and **2,4DCPAIL**.




Fig. 2. ^1H NMR spectra of inulin, **CAIL**, **PAIL**, **2CPAIL**, **4CPAIL**, and **2,4DCPAIL**.




Fig. 3. The antifungal activity of inulin, CAIL, PAIL, 2CPAIL, 4CPAIL, and 2,4DCPAIL against *C. lagenarium*.




Fig. 4. The antifungal activity of inulin, CAIL, PAIL, 2CPAIL, 4CPAIL, and 2,4DCPAIL against *P. asparagi*.




Fig. 5. The antifungal activity of inulin, CAIL, PAIL, 2CPAIL, 4CPAIL, and 2,4DCPAIL against *F. oxysporum*.

3.2. Antifungal activity

The antifungal activities of inulin, CAIL, PAIL, 2CPAIL, 4CPAIL, and 2,4DCPAIL against *C. lagenarium*, *P. asparagi*, and *F. oxysporum* were tested and the results are shown in Figs. 3–5. As shown in those figures, inulin did not inhibit the growth of all the measured fungi under the tested concentration (1.0–0.1 mg/mL), which is in accord with the earlier report (Ren et al., 2012). All the inulin derivatives show inhibitory activity against three fungi and the antifungal indices of them enhanced with increasing concentration. After the reaction of inulin and chloroacetyl chloride, the antifungal ability of CAIL increases and the antifungal indices at 1.0 mg/mL are 22%, 17%, and 12% against *C. lagenarium*, *P. asparagi*, and *F. oxysporum*, respectively. This is possible due to the introducing of chloroacetyl group in CAIL. As shown in Figs. 3–5, PAIL has lower antifungal ability than CAIL on all the measured fungi. It would be reasonable to presume that chlorine atom should be the important factor which influences

the antifungal activity of inulin derivatives. Meanwhile, as shown in three figures, the antifungal indices had the same order as follow: 2,4DCPAIL > 4CPAIL > 2CPAIL > CAIL > PAIL > inulin. It is easily to find that the order of antifungal indices is in accord with the number of chlorine atom in those compounds. 2,4DCPAIL, which has two chlorine atoms, had the best inhibitory activity among all of the measured samples. And the inhibitory indices of 2,4DCPAIL against *C. lagenarium*, *P. asparagi*, and *F. oxysporum* are 67%, 47%, and 43%, respectively at 1.0 mg/mL. The results further confirm that chlorine grafted in the synthesized inulin derivatives contribute a lot to the antifungal action and consequently increase the antifungal activity of them.

4. Conclusions

The development of fungicides has attracted people's attention along with the food safety. And environmental problems become serious caused by the chemical pesticides. More and more effective fungicides have been prohibited in recent years, and the requirement for "green pesticide" is urgent. In this paper, four inulin derivatives containing benzene or chlorinated benzene with the Schiff bases of 4-amino-pyridine were designed and synthesized, and their antifungal activity against three kinds of phytopathogens was estimated by hypha measurement *in vitro*. All the inulin derivatives have good solubility in water, and exhibit higher inhibition indices than inulin. These data demonstrate that the chemical modification of inulin with chlorinated benzene would lead to an enhancement of the biological activity against some plant pathogenic fungi. In addition, the antifungal activity of these inulin derivatives is influenced by the number of chlorine grafted on to the periphery of molecular chains. Further study will be carried out to ascertain this hypothesis.

Acknowledgments

We thank the project of the 12th Five-Year Science and Technology Plan for Agriculture (2012BAD32B09), the Science and Technology Project of Yantai (20121321) and the Taishan Scholar Program of Shandong Province for financial support of this work.

References

- Beylot, M. (2006). Effects of inulin-type fructans on lipid metabolism in man and in animal models. *British Journal of Nutrition*, 93(Suppl. 1), S163–S168.
- Causey, J. L., Feirtag, J. M., Gallaher, D. D., Tungland, B. C., & Slavin, J. L. (2000). Effects of dietary inulin on serum lipids, blood glucose and the gastrointestinal environment in hypercholesterolemic men. *Nutrition Research*, 20(2), 191–201.
- Chen, Y. J., & Dai, G. H. (2012). Antifungal activity of plant extracts against *Colletotrichum lagenarium*, the causal agent of anthracnose in cucumber. *Journal of the Science of Food and Agriculture*, 92(9), 1937–1943.
- Elena, K. (2006). First report of *Phomopsis asparagi* causing stem blight of asparagus in Greece. *Plant Pathology*, 55, 300.
- Guo, Z. Y., Ren, J. M., Dong, F., Wang, G., & Li, P. C. (2013). Comparative study of the influence of active groups of chitosan derivatives on antifungal activity. *Journal of Applied Polymer Science*, 127(4), 2553–2556.
- Habibi, Y., & Dufresne, A. (2008). Highly filled bionanocomposites from functionalized polysaccharide nanocrystals. *Biomacromolecules*, 9(7), 1974–1980.
- Jasso de Rodríguez, D., Hernández-Castillo, D., Rodríguez-García, R., & Angulo-Sánchez, J. L. (2005). Antifungal activity *in vitro* of Aloe vera pulp and liquid fraction against plant pathogenic fungi. *Industrial Crops and Products*, 21(1), 81–87.
- Kaboudin, B., Afsharinezhad, & Mohammad, B. (2010). Synthesis of a novel class of β -lactam derivatives of 1-aminophosphonates by staudinger ketene-imine [2+2]-cycloaddition reaction. *Synthesis*, 20, 3504–3508.
- Karthikeyan, M. S., Prasad, D. J., Poojary, B., Bhat, K. S., Holla, B. S., & Kumari, N. S. (2006). Synthesis and biological activity of Schiff and Mannich bases bearing 2,4-dichloro-5-fluorophenyl moiety. *Bioorganic & Medicinal Chemistry*, 14(22), 7482–7489.
- Li, Q., Ren, J. M., Dong, F., Feng, Y., Gu, G. D., & Guo, Z. Y. (2013). Synthesis and antifungal activity of thiazole-functionalized chitosan derivatives. *Carbohydrate Research*, 373(0), 103–107.

- Li, R. C., Guo, Z. Y., & Jiang, P. A. (2010). Synthesis, characterization, and antifungal activity of novel quaternary chitosan derivatives. *Carbohydrate Research*, 345(13), 1896–1900.
- Panicker, C. Y., Varghese, H. T., Philip, D., & Nogueira, H. I. S. (2006). FT-IR, FT-Raman and SERS spectra of pyridine-3-sulfonic acid. *Spectrochimica Acta, Part A: Molecular and Biomolecular Spectroscopy*, 64(3), 744–747.
- Ren, J. M., Liu, J. L., Dong, F., & Guo, Z. Y. (2011). Synthesis and hydroxyl radicals scavenging activity of *N*-(aminoethyl)inulin. *Carbohydrate Polymers*, 85(1), 268–271.
- Ren, J. M., Wang, P. B., Dong, F., Feng, Y., Peng, D. J., & Guo, Z. Y. (2012). Synthesis and antifungal properties of 6-amino-6-deoxyinulin, a kind of precursors for facile chemical modifications of inulin. *Carbohydrate Polymers*, 87(2), 1744–1748.
- Rogge, T. M., & Stevens, C. V. (2004). Facilitated synthesis of inulin esters by transesterification. *Biomacromolecules*, 5(5), 1799–1803.
- Sha, J. J. (1994). *Handbook of chemical product* (3rd ed., pp. 190–192). Beijing: Chemical Industry.
- Sajomsang, W. (2010). Synthetic methods and applications of chitosan containing pyridylmethyl moiety and its quaternized derivatives. *Carbohydrate Polymers*, 80(3), 631–647.