

## CARBOHYDRATE RESEARCH, VOL. 113 (1983)

## SUBJECT INDEX

- Acetal-exchange reactions, preparation of mixed-acetal derivatives of carbohydrates by, 225
- 2-Acetamido-1-*N*-acyl-2-deoxy- $\beta$ -D-glucopyranosylamines, degradation during the hydrazinolysis of, C13
- Acid-catalyzed interconversion of diastereoisomeric methyl 2,3:4,6-di-*O*-benzylidene- $\alpha$ -D-mannopyranosides, 326
- Activity of fructanase in batch cultures of oral streptococci, 101
- Addition of 1-nitroalkanes to methyl 2,3-*O*-isopropylidene- $\beta$ -D-ribo-pentodialdo-1,4-furanoside and *N*<sup>6</sup>-benzoyl-2',3'-*O*-isopropylidene-adenosine-5'-aldehyde, 233
- Adenosine-5'-aldehyde, *N*<sup>6</sup>-benzoyl-2',3'-*O*-isopropylidene-, addition of 1-nitroalkanes to, 233
- Alditol acetates for monosaccharide analysis, a simple and rapid preparation of, 291
- Alditols, 2-haloethylidene acetals of some, 336
- $\omega$ -Aminoalkyl  $\beta$ -glycosides of *N*-acetylmuramyl-L-alanyl-D-isoglutamine, and their conjugates with meningococcal group C polysaccharide, 57
- Analysis, a simple and rapid preparation of alditol acetates for monosaccharide, 291
- 3,4-Anhydro-2,6-dideoxy- $\beta$ -L-*lyxo*- and *ribo*-hexopyranosides, methyl, ring opening of, 189
- Anthracyclines fragments, synthesis of, 21
- T-Antigene, synthesis of, 203
- Antitumor activity, purification of a new polysaccharide from *Cyttaria johowii* (Esp.) and studies on its, 331
- Antitumoral anthracycline antibiotics, synthesis of fragments of, 21
- L-Arabinan from apple-juice concentrates, 339
- $\alpha$ -L-Arabinofuranosyl- $\beta$ -D-xylotriose and -tetraose from hemicellulose by hemicellulose degradation, 125
- Base transformation of 2,4-*O*-benzylidene-1,6-di-*O*-*p*-tolylsulfonyl-D-glucitol into 1,3-anhydro-2,4-*O*-benzylidene-D-glucitol, reaction pathway in the, 321
- Biotechnology, Volume 1 (book review) C29
- Branched-chain amino sugars, a stereochemically defined route to, *via* spiro-oxiranes, C6
- Byproducts in the preparation of penta-*O*-acetyl-D-hexononitriles from D-galactose and D-glucose, 241
- Carbohydrate, food, developments in (book review), C27
- Carbohydrates, glycosylation of 1,2-*O*-cyanoethylidene derivatives of, C1
- Carbohydrate, side chain of native ovalbumin, a <sup>13</sup>C-n.m.r.-spectral study of galactosyltransferase specificity with regard to the, 141
- Concanavalin A, the positively cooperative binding of, to corn starch, 133
- Convenient, laboratory procedure for producing solid D-*arabino*-hexos-2-ulose (D-glucosone), 151
- Convenient, laboratory procedure for reducing D-glucosone to D-fructose, 159
- 1,2-*O*-Cyanoethylidene derivatives of carbohydrates, glycosylation of, C1
- Degradation during the hydrazinolysis of 2-acetamido-1-*N*-acyl-2-deoxy- $\beta$ -D-glucopyranosylamines, C13
- Developments in food carbohydrate—3 (book review), C27
- Diastereotopic substrates of  $\beta$ -D-galactosidase from *Escherichia coli* as probes for a catalytically active, protonating group, 71
- 2,6-Dideoxy- $\beta$ -L-*lyxo*- and *ribo*-hexopyranosides, methyl 3,4-anhydro-, ring opening of, 189
- Disaccharides (book review), C27
- Drosera capensis*, structure of acidic polysaccharide from mucin secreted by, 113
- Enzymic oxidation of D-*arabino*-hexos-2-ulose (D-glucosone) to D-*arabino*-2-hexulosonic acid ("2-keto-D-gluconic acid"), 163
- Facile synthesis of benzyl 2-acetamido-6-*O*-acetyl-4-*O*-(6-*O*-acetyl-3,4-*O*-isopropylidene- $\beta$ -D-galactopyranosyl)-2-deoxy- $\beta$ -D-glucopyranoside, a key intermediate for the synthesis of *O*- $\alpha$ -L-fucopyranosyl-(1 $\rightarrow$ 2)-*O*- $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 4)-*O*-[ $\alpha$ -L-fucopyranosyl-(1 $\rightarrow$ 3)]-2-acetamido-2-deoxy-D-glucopyranose, C18

- Formation and X-ray structural analysis of 2,3,4-tri-*O*-acetyl-1,5-anhydro-5,6-dideoxy-5-*C*-[(*S*)-phenylphosphinyl]-L-iditol, a further mechanistic study of the ring enlargement of (5*RS*)-5-deoxy-5-*C*-[(*RS*)-phosphinyl]- $\alpha$ -D-xyl/o-hexofuranoses to give 5-deoxy-5-*C*-[(*RS*)-phosphinyl]- $\alpha$ , $\beta$ -L-idopyranoses, 31
- Fructanase in batch cultures of oral streptococci, activity of, 101
- D-Fructose, convenient laboratory procedure for reducing D-glucosone to, 159
- D-Galactopyranose, *O*- $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 3)-*O*-(2-acetamido-2-deoxy- $\alpha$ -D-galactopyranosyl)-(1 $\rightarrow$ 6)-, synthesis of, 203
- $\gamma$ -D-Galactopyranoside, benzyl 2-acetamido-3,6-di-*O*-(2-acetamido-2-deoxy- $\beta$ -D-glucopyranosyl)-2-deoxy-, synthesis of, 63
- 4-*O*- $\gamma$ -D-Galactopyranosyl-D-galactose: efficient synthetic routes from "polygalacturonic acid", 219
- $\beta$ -D-Galactosidase, from *Escherichia coli*, diastereotopic substrates of, as probes for a catalytically active protonating group, 71
- $\beta$ -D-Galactosidase-bound diastereotopic substrate, location of a proton-donating group at the *re*-face of a, 93
- Gangliosides,  $^{13}\text{C}$ -n.m.r. spectra of, 173
- D-Glucal, 5-thio-, and inhibitor of glycosidases, synthesis and conformation of, 166
- D-Glucan, alkali-soluble, from the fruit body of *Laetiporus sulphureus*, 281
- D-Glucans, mechanism of synthesis of, by D-glucosyltransferases from *Streptococcus mutans* 6715, 301
- Glucitol into 1,3-anhydro-2,4-*O*-benzylidene-D-glucitol, reaction pathway in the base transformation of 2,4-*O*-benzylidene-1,6-di-*O*-p-tolylsulfonyl-D-, 321
- D-Glucopyranose, *O*- $\gamma$ -L-fucopyranosyl-(1 $\rightarrow$ 2)-*O*- $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 4)-*O*-[ $\gamma$ -L-fucopyranosyl-(1 $\rightarrow$ 3)]-2-acetamido-2-deoxy-, a key intermediate for the synthesis of, C18
- D-Glucopyranose, *O*- $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 3)-*O*-(2-acetamido-2-deoxy- $\gamma$ -D-galactopyranosyl)-(1 $\rightarrow$ 6)-, synthesis of, 203
- $\beta$ -D-Glucopyranoside, benzyl 2-acetamido-6-*O*-acetyl-4-*O*-(6-*O*-acetyl-3,4-*O*-isopropylidene- $\beta$ -D-galactopyranosyl)-2-deoxy-, a facile synthesis of, C18
- $\beta$ -D-Glucopyranoside, *p*-nitrophenyl 2-acetamido-2-deoxy-4-*O*- $\beta$ -D-galactopyranosyl-, synthesis of, 257
- D-Glucosone, convenient laboratory procedure for reducing, to D-fructose, 159
- D-Glucosone, hexos-2-ulose, convenient laboratory procedure for producing solid, 151
- D-Glucosyltransferases, from *Streptococcus mutans* 6715, mechanism of synthesis of D-glucans by, 301
- Glycosylation of 1,2-*O*-cyanoethylidene derivatives of carbohydrates, C1
- G<sub>M1</sub>- and asialo G<sub>M1</sub> ganglioside,  $^{13}\text{C}$ -n.m.r. spectra of, 173
- G<sub>M1</sub>-oligosaccharide,  $^{13}\text{C}$ -n.m.r. spectrum of, 173
- 2-Haloethylidene acetals of some alditols, 336
- Hemicellulose, degradation by hemicellulase into  $\gamma$ -L-arabinofuranosyl- $\beta$ -D-xyl/o-triose and -tetraose of, 125
- Heterocyclization, under acetylation conditions, the mechanism of, 273
- $\gamma$ -D-xyl/o-Hexofuranoses, (5*RS*)-5-deoxy-5-*C*-[(*RS*)-phosphinyl]-, a further mechanistic study of the ring enlargement of, to give 5-deoxy-5-*C*-[(*RS*)-phosphinyl]- $\alpha$ , $\beta$ -L-idopyranoses, 31
- D-Hexonitriles, penta-*O*-acetyl-, from D-galactose and D-glucose, byproducts in the preparation of, 241
- $\beta$ -L-Ido-Hexopyranoside, methyl 2,6-dideoxy-4-*O*-(2,3,6-trideoxy- $\gamma$ -L-glycerol- and  $\gamma$ -L-erythro-hexopyranosyl)-, synthesis of, 21
- Hexopyranosides, methyl 3,4-anhydro-2,6-dideoxy- $\beta$ -L-lyxo- and -thio-, ring opening of, 189
- D-arabino-Hexos-2-ulose, convenient laboratory procedure for producing solid, 151
- D-arabino-Hexos-2-ulose (D-glucosone), enzymic oxidation of, to D-arabino-2-hexulosonic acid ("2-keto-D-gluconic acid"), 163
- D-arabino-2-Hexulosonic acid ("2-keto-D-gluconic acid"), by enzymic oxidation of D-arabino-hexos-2-ulose (D-glucosone), 163
- Hydrazinolysis of 2-acetamido-1-*N*-acetyl-2-deoxy- $\beta$ -D-glucopyranosylamines, degradation during the, C13
- Hydrazones, (2,4-dichlorophenoxy)acetyl-, of saccharides, 273
- L-Iditol, 2,3,4-tri-*O*-acetyl-1,5-anhydro-5,6-dideoxy-5-*C*-[(*S*)-phenylphosphinyl]-, formation and X-ray structural analysis of, 31
- $\beta$ -L-Idofuranose, 5-deoxy-5-*C*-(diphenylphosphinyl)-1,2-*O*-isopropylidene-3-*O*-methyl-, synthesis of, C10
- D-Isoglutamine, *N*-acetylmuramyl-L-alanyl-, *m*-aminoalkyl  $\beta$ -glycosides of, and their conjugates with meningococcal group C polysaccharide, 57
- D-Isoglutamines, *N*-acetyl-6-*O*-(*m*-aminoacyl)-muramyl-L-alanyl-, methyl  $\beta$ -glycosides of, and their conjugates with meningococcal group C polysaccharide 45

- Location of a proton-donating group at the re-face of a  $\beta$ -D-galactosidase-bound, diastereotopic substrate, 93
- Maltotriitol, 2,3,4,6,2',3',6',2'',3'',6''-deca-*O*-methyl-, the selective cleavage of, into 2,3,4,6,2',3',6'-hepta-*O*-methylmaltose and 2,3,6-tri-*O*-methyl-D-glucitol, C22
- $\alpha$ -D-Mannopyranoside, *p*-nitrophenyl 6-*O*-(2-acetamido-2-deoxy-3-*O*- and -4-*O*- $\beta$ -D-galactopyranosyl- $\beta$ -D-glucopyranosyl)-, synthesis of, 257
- $\alpha$ -D-Mannopyranosides, methyl 2,3:4,6-di-*O*-benzylidene-, the acid-catalyzed interconversion of diastereoisomeric, 326
- Mechanism of synthesis of D-glucans by D-glucosyltransferases from *Streptococcus mutans* 6715, 301
- Methyl  $\beta$ -glycosides of *N*-acetyl-6-*O*-( $\omega$ -aminoacyl)muramyl-L-alanyl-D-isoglutamines, and their conjugates with meningococcal group C polysaccharide, 45
- Microbial fundamentals (book review), C29
- Mixed-acetal derivatives of carbohydrates, preparation of, by acetal-exchange reactions, 225
- Mucin secreted by *Drosera capensis*, structure of acidic polysaccharide from, 113
- Nigerose,  $\alpha$ - and  $\beta$ -, octaacetate, a simple preparation of, 281
- Nigerotriose,  $\beta$ -, a simple preparation of, 281
- 1-Nitroalkanes, addition of, to methyl 2,3-*O*-isopropylidene- $\beta$ -D-ribo-pentodialdo-1,4-furanoside and *N*<sup>6</sup>-benzoyl-2',3'-*O*-isopropylidene-adenosine-5'-aldehyde, 233
- <sup>13</sup>C-N.m.r.-spectral study of galactosyltransferase specificity with regard to the carbohydrate side-chain of native ovalbumin, 141
- <sup>13</sup>C-N.m.r. spectra of gangliosides, 173
- N.m.r. studies of D-ribosylamines derived from primary amines, 1
- C-Nucleosides, the synthesis of precursors of, C16
- Oligosaccharides, a possibility for sequential analysis of, by stepwise degradation, C22
- Oral streptococci, activity of fructanase in batch cultures of, 101
- Pentodialdo-1,4-furanoside, methyl 2,3-*O*-isopropylidene- $\beta$ -D-ribo-, addition of 1-nitroalkanes to, 233
- "Polygalacturonic acid", efficient synthetic routes to 4-*O*- $\alpha$ -D-galactopyranosyl-D-galactose from, 219
- Polysaccharide, acidic, from mucin secreted by *Drosera capensis*, structure of, 113
- Polysaccharide from *Cyttaria johowii* (Esp.) and studies on its antitumor activity, purification of a new, 331
- Positively cooperative binding of concanavalin A to corn starch: the isotherm of binding and a measure of the cooperativity, 133
- Possibility for sequential analysis of oligosaccharides by stepwise degradation. The selective cleavage of 2,3,4,6,2',3',6',2'',3'',6''-deca-*O*-methylmaltotriitol into 2,3,4,6,2',3',6'-hepta-*O*-methylmaltose and 2,3,6-tri-*O*-methyl-D-glucitol, C22
- Preparation of mixed-acetal derivatives of carbohydrates by acetal-exchange reactions, 225
- Primary amines, n.m.r. studies of D-ribosylamines derived from, 1
- Purification of a new polysaccharide from *Cyttaria johowii* (Esp.) and studies on its antitumor activity, 331
- Reaction pathway in the base transformation of 2,4-*O*-benzylidene-1,6-di-*O*-*p*-tolylsulfonyl-D-glucitol into 1,3-anhydro-2,4-*O*-benzylidene-D-glucitol, 321
- D-Ribosylamines derived from primary amines, n.m.r. studies of, 1
- Saccharide (2,4-dichlorophenoxy)acetylhydrazones, the mechanism of heterocyclization under acetylation conditions, 273
- Simple and rapid preparation of alditol acetates for monosaccharide analysis, 291
- Simple preparation of  $\alpha$ - and  $\beta$ -nigerose octaacetate and  $\beta$ -nigerotriose hendecaacetate by the acetolysis of an alkali-soluble D-glucan from the fruit body of *Laetiporus sulphureus*, 281
- Spiro-oxiranes, a stereochemically defined route to branched-chain amino sugars *via*, C6
- Starch, corn, the positively cooperative binding of, to concanavalin A, 133
- Stereochemically defined route to branched-chain amino sugars *via* spiro-oxiranes, and related studies, C6
- Synthesis and conformation of 5-thio-D-glucal, an inhibitor of glycosidases, 166
- Synthesis of 5-deoxy-5-*C*-(diphenylphosphinyl)-1,2-*O*-isopropylidene-3-*O*-methyl- $\beta$ -L-idofuranose, C10
- Synthesis of *p*-nitrophenyl 2-acetamido-2-deoxy-4-*O*- $\beta$ -D-galactopyranosyl- $\beta$ -D-glucopyranoside, and *p*-nitrophenyl 6-*O*-(2-acetamido-2-deoxy-3-*O*- and -4-*O*- $\beta$ -D-galactopyranosyl- $\beta$ -D-glucopyranosyl)- $\alpha$ -D-mannopyranoside, 257

- Synthesis of *C*-nucleoside precursors: 3-(alditol-1-yl)-5-phenyl-1,2,4-triazolo[3,4-*a*]phthalazines, C16
- Synthesis of *S*- and *Se*-(2,3,4,6-tetra-*O*-acetyl- $\beta$ -D-glucosyl) thiophosphates and selenophosphates, efficient, 317
- Synthesis of benzyl 2-acetamido-3,6-di-*O*-(2-acetamido-2-deoxy- $\beta$ -D-glucopyranosyl)-2-deoxy- $\alpha$ -D-galactopyranoside, 63
- Synthetic routes to 4-*O*- $\alpha$ -D-galactopyranosyl-D-galactose from "polygalacturonic acid", efficient, 219
- S*- and *Se*-(2,3,4,6-Tetra-*O*-acetyl- $\beta$ -D-glucosyl) thiophosphates and selenophosphates, efficient synthesis of, 317
- Thiophosphates and selenophosphates, efficient synthesis of *S*- and *Se*-(2,3,4,6-tetra-*O*-acetyl- $\beta$ -D-glucosyl), 317
- 1,2,4-Triazolo[3,4-*a*]phthalazines, 3-(alditol-1-yl)-5-phenyl-, C16
- $\beta$ -D-Xylo-triose and -tetraose from hemicellulose by hemicellulose degradation, 125

---

### INFORMATION FOR AUTHORS

Papers should deal with original work not published elsewhere in an accessible language. Manuscripts (3 copies) should be prepared in double-spaced typing.

Flexibility in presentation of results will be allowed and the general style of preparation should be similar to that adopted in this issue. Careful attention must be paid to nomenclature, which should conform to the British-American Rules of Carbohydrate Nomenclature [*J. Org. Chem.*, 28 (1963) 281; *J. Chem. Soc.*, (1962) 5307] until there is full international agreement by the International Union of Pure and Applied Chemistry.

References should be given at the end of the paper; they should be numbered and the numbers should appear in the text at the appropriate places as superscripts without parentheses. The abbreviation of journal titles should conform to those adopted by the *Bibliographic Guide for Editors and Authors*, Chemical Abstracts Service, Columbus, OH 43210, and should be followed by volume number, year (in parentheses) and inclusive pagination. For example: *Carbohydr. Res.*, 6 (1968) 63-74. Each paper should have an abstract of 80-200 words. For papers in French or German, an English abstract should be supplied.

Line drawings should be in a form suitable for reproduction, drawn in Indian ink on drawing paper or tracing paper. Each drawing should be on a separate sheet and not glued to the manuscript or drawn on the manuscript page.

Photographs should be submitted as clear black-and-white prints on glossy paper.

Tables, and a list of legends to the illustrations, should be typed on separate pages of the manuscript. Proofs of Preliminary Communications will be checked in the editorial office, and will not be sent to the author unless specifically requested. If such a request is made, the publishers cannot be held responsible for possible delay in publication.