SUBJECT INDEX

- Alpha-amylase, photolabile derivatives of maltose and maltotriose as ligands for the affinity labelling of the maltodextrin-binding site in porcine pancreatic, 129
- Alternative synthesis and enzyme-inhibitory activity of methyl l'-epiacarviosin and its 6-hydroxy analog, 279
- Arabinoxylan oligosaccharides containing two or three branched xylose residues, ¹H-n.m.r. study of enzymically generated wheat-endosperm, 19
- Arylation of 4,6-O-benzylidene-3-O-tert-butyldimethylsilyl-1-tributylstannyl-p-glucal, a stereoselective route to the basic skeleton of papulacandins and chaetiacandin by palladiummediated, 157
- 5-Azido-3-*O*-benzoyl-6-*O*-tert-butyldiphenylsilyl-5-deoxy-1,2-*O*-isopropylidene-β-L-talofuranose, crystal and molecular structure of, 271
- 2-Azidoethyl glycosides of potential use for the preparation of neoglycoconjugates, synthesis of, 303
- Azo dyes containing naphthalene nuclei, induced circular dichroism spectra of complexes of cyclomalto-oligosaccharides and, 71
- Bioglycan-immunomodulator (mytilan) isolated from the mussel *Crenomytilus grayanus*, structure of a. 221
- Boswellia serrata, isolation and structure of 4-Omethyl-glucuronoarabinogalactan from, 321
- Cell-surface receptors for C-glycosyl compounds, 243
- Cryptococcus neoformans serotype D, glucuronoxylomannan of: structural analysis by gasliquid chromatography-mass spectrometry and by ¹³C-n.m.r. spectroscopy, 263
- Crystal and molecular structure of 5-azido-3-*O*-benzoyl-6-*O*-tert-butyldiphenylsilyl-5-deoxy-1,2-*O*-isopropylidene-β-L-talofuranose, 271
- Crystal structure of D-threitol at 119 K and 298 K,
- Crystal structure of lactitol (4-O- β -D-galactopyranosyl-D-glucitol), 45
- Crystal structure of lactitol (4-*O*-β-D-galactopyranosyl-D-glucitol) dihydrate, 53
- Cyclodextrin-glucanosyltransferase, enzymatic synthesis of linear and branched gluco-oligosaccharides using, 61
- Cyclomaltohexaose, a convenient route to polysubstituted derivatives by multiple tritylation, 287

- Cyclomaltohexaoses and cyclomaltoheptaoses, di-O-triphenylmethyl (trityl-), preparation of and characterisation of three positional isomers by the "hex-5-enose degradation", 1
- Cyclomalto-oligosaccharides and azo dyes containing naphthalene nuclei, induced circular dichroism spectra of complexes of, 71
- Cyclomalto-oligosaccharides (cyclodextrins), distribution of substituents in *O*-(2-hydroxypropyl) derivatives of, 99
- Deacetylation of derivatives of 1,2-O-isopropylidene-α-D-hexofuranoses, enzymic, 137
- N_ε-(1-Deoxy-D-fructos-1-yl)hippuryl-lysine, preparation and characterisation of, 293
- Disaccharide-glycosides related to carageenans, stereoselective synthesis of, 147
- Distribution of substituents in O-(2-hydroxypropyl) derivatives of cyclomalto-oligosaccharides (cyclodextrins), 99
- Ehlers-Danlos syndrome, proteoglycans from human burn hypertrophic scar in, 209
- Enzymatic synthesis of linear and branched glucooligosaccharides, using cyclodextrin-glucanosyltransferase, 61
- Enzyme expressed by the *Pseudomonas saccha*rophila maltotetraohydrolase gene (mta) in *Es*cherichia coli, properties of, 255
- Enzyme-inhibitory activity of methyl 1'-epiacarviosin and its 6-hydroxy analog, alternative synthesis and, 279
- Enzymic deacetylation of derivatives of 1,2-Oisopropylidene-α-D-hexofuranoses, 137
- Epiacarviosin and its 6-hydroxy analog, alternative synthesis and enzyme-inhibitory activity of methyl 1'-, 279
- Escherichia coli, properties of the enzyme expressed by the *Pseudomonas saccharophila* maltotetraohydrolase gene (mta) in, 255
- Exopolysaccharides from *Rhizobium meliloti* YE-2 grown under different osmolarity conditions, viscoelastic properties, 227
- 4-O-β-D-Galactopyranosyl-D-glucitol (lactitol), crystal structure of, 45
- 4-O-β-D-Galactopyranosyl-D-glucitol (lactitol) dihydrate, crystal structure of, 53
- Gluco-oligosaccharides, enzymatic synthesis of linear and branched, using cyclodextrin-glucanosyltransferase, 61

- Glucuronoxylomannan of *Cryptococcus neoformans* serotype D: structural analysis by gasliquid chromatography-mass spectrometry and by ¹³C-n.m.r. spectroscopy, 263
- Glycoproteins, structural analysis of monosulfated side-chain oligosaccharides isolated from human tracheobronchial mucous, 187
- Glycosides of potential use for the preparation of neoglycoconjugates, synthesis of 2-azidoethyl, 303
- C-Glycosyl compounds bind to receptors on the surface of *Escherichia coli* cells and can target proteins to the organism, 243
- Glycosyl derivatives of methyl α-L-rhamnopyranoside, synthesis of di-O-, 109
- Glycosyl phosphates and azides, synthesis of, 169
- Heparin, two series of oligosaccharides derived from porcine intestinal mucosal, by degradation with heparinse, one- and two-dimensional ¹³C-n.m.r. characterization of, 81
- Hippuryl-lysine, preparation and characterisation of N_c -(1-deoxy-D-fructos-1-yl)-, 293
- Human burn hypertrophic scar, proteoglycans from, 209
- Hyaluronic acid and a $(1 \rightarrow 4)-\beta$ -D-xylan, structural elucidation of two extracellular polysaccharides of *Pasteurella multocida* as, 329
- O-(2-Hydroxypropyl) derivatives of cyclomaltooligosaccharides (cyclodextrins), distribution of substituents in, 99
- Induced circular dichroism spectra of complexes of cyclomalto-oligosaccharides and azo dyes containing naphtalene nuclei. 71
- Isolation and structure of a 4-O-methyl-glucuronoarabinogalactan from *Boswellia serrata*, 321 1,2-O-Isopropylidene-α-p-hexofuranoses.
- zymic deacetylation of derivatives of, 137
- Lactitol (4-*O*-β-D-galactopyranosyl-D-glucitol), crystal structure of, 45
- Lactitol (4-*O*-β-D-galactopyranosyl-D-glucitol) dihydrate, crystal structure of, 45
- Maltose and maltotriose, photolabile derivatives as ligands for the affinity labelling of the maltodextrin-binding site in porcine pancreatic alpha-amylase. 129
- Maltotetrahydrolase gene (mta), properties of the enzyme expressed in Escherichia coli by the Pseudomonas saccharophila, 255
- Methyl 3-*O*-(3,6-anhydro-β-D-galactopyranosyl)α-D-galactopyranoside and methyl 3,6-anhydro-4-*O*-β-D-galactopyranoside, efficient and stereoselective synthesis of, 147

- 4-O-Methyl-glucuronoarabinogalactan from *Boswellia serrata*, isolation and structure of a, 321
- Methyl α-L-rhamnopyranoside, synthesis of di-Oglycosyl derivatives of, 109
- Multiple tritylation, a convenient route to polysubstituted derivatives of cyclomaltohexaose, 287
- Mytilan isolated from the mussel *Crenomytilus* grayanus, structure of a bioglycan-immuno-modulator, 221
- Neoglycoconjugates, synthesis of 2-azidoethyl glycosides of potential use for the preparation of, 303
- ¹³C-N.m.r. characterization, one- and two-dimensional, of two series of oligosaccharides derived from porcine intestinal mucosal heparin by degradation with heparinase, 81
- ¹H-N.m.r. study of enzymically generated wheatendosperm arabinoxylan oligosaccharides containing two to three branched xylose residues, 19
- Oligosaccharides containing two or three branched xylose residues, ¹H-n.m.r. study of enzymically generated wheat-endosperm arabinoxylan, 19
- Oligosaccharides isolated from human tracheobronchial mucous glycoproteins, structural analysis of monosulfated side-chain, 187
- Papulacandins and chaetiacandin, a stereoselective route to their basic skeleton by a palladium-mediated arylation of 4,6-O-benzylidene-3-O-tert-butyldimethylsilyl-1-tributylstannyl-pglucal, 157
- Pasteurella multocida extracellular polysaccharides, structural elucidation as hyaluronic acid and a $(1 \rightarrow 4)$ - β -D-xylan, 329
- Phosphates and azides, synthesis of glycosyl, 169 Photolabile derivatives of maltose and maltotriose as ligands for the affinitiy labelling of the maltodextrin-binding site in porcine pancreatic alpha-amylase, 129
- Polysaccharides of *Pasteurella multocida*, structural elucidation of hyaluronic acid and a $(1 \rightarrow 4)$ - β -D-xylan as extracellular, 329
- Preparation and characterisation of N_e -(1-deoxy-D-fructos-1-yl)hippuryl-lysine, 293
- Proteoglycans from a patient with Ehlers-Danlos syndrome, 209
- Proteoglycans, from human burn hypertrophic scar, isolation and characterization of, 209
- 3-Pyrrolecarbohydrazide, synthesis and reactions of 2-methyl-5-(D-arabino-tetrahydroxybutyl), 311

- Receptors of the surface of *Escherichia coli* cells and can target proteins to the organism, *C*-glycosyl compounds bind to, 243
- Rhizobium melitoli YE-2 grown under different osmolarity conditions, viscoelastic properties of exopolysaccharides from, 227
- Starch from waxy rice (IR29), structure and properties during development of the grain, 235
- Stereoselective route to the basic skeleton of the antibiotics papulacandin and chaetiacandin, using palladium-mediated arylation of a D-glucal derivative, 157
- Structure and properties of waxy-rice (IR29) starch during development of the grain, 235
- Structure of mytilan, a bioglycan-immunomodulator isolated from the mussel *Crenomytilus* grayanus, 221
- Synthesis of di-O-glycosyl derivatives of methyl α-L-rhamnopyranoside, 109
- Synthesis of glycosyl phosphates and azides, 169

- Synthesis of methyl 3-*O*-(3,6-anhydro-β-D-galactopyranosyl)-α-D-galactopyranoside and methyl 3,6-anhydro-4-*O*-β-D-galactopyranosyl-α-D-galactopyranoside, efficient and stereoselective, 147
- Synthesis of (+)-validamycin H, total, 299
- β-L-Talofuranose, crystal and molecular structure of 5-azido-3-O-benzoyl-6-O-tert-butyldiphenylsilyl-5-deoxy-1,2-O-isopropylidene-, 271
- D-Threitol, crystal structure at 119 K and 298 K,
- Total synthesis of (+)-validamycin H, 299
 Tritylation, a convenient route to polysubstituted
 derivatives of cyclomaltohexaose by multiple,
 287
- Validamycin H, total synthesis of (+)-, 299 Viscoelastic properties of exopolysaccharides from *Rhizobium meliloti* YE-2 grown under different osmolarity conditions, 227
- Waxy-rice (IR29) starch, structure and properties during development of the grain, 235