

CONTENTS

<i>Publisher's Note</i>	v
The crystal structure of the inclusion complex of cyclomaltoheptaose (β -cyclodextrin) with 4- <i>tert</i> -butyltoluene I.M. Mavridis and E. Hadjoudis (Athens, Greece)	1
The crystal and molecular structures of the hexitol hexa-acetates J. Kopf, M. Morf, B. Zimmer (Hamburg, Germany), M. Bischoff and P. Köll (Oldenburg, Germany)	17
The crystal structure of α -D-glucopyranosyl potassium hydrogenphosphate T. Lis (Wrocław, Poland)	33
Molecular and crystal structure of konjac glucomannan in the mannan II polymorphic form T. Yui, K. Ogawa and A. Sarko (Syracuse, NY, USA)	41
Molecular and crystal structure of the regenerated form of (1 $\rightarrow$ 3)- α -D-mannan T. Yui, K. Ogawa and A. Sarko (Syracuse, NY, USA)	57
Synthesis and conformational analysis of substituted 2,6-dioxabicyclo[3.1.1]heptanes: 1,3-anhydro-2,4-di- <i>O</i> -benzyl-6-deoxy- β -D-glucopyranose by ^1H -NMR spectroscopy and molecular mechanics calculation X. Wu, F. Kong, D. Lu and P. Zhang (Beijing, China)	75
Addition and cyclisation reactions of sugar phenylhydrazone derivatives via a 1,4-elimination process: preparation of 3-(2-acetamido-1-phenylhydrazonobutyl)-2(1 <i>H</i>)-quinoxalinone and 3-(1-acetamidopropyl)-pyrazolo[3,4- <i>b</i>]quinoxaline (flavazole) derivatives L. Somogyi (Debrecen, Hungary)	89
Stereoselective syntheses of a di-, tri-, and tetra-saccharide fragment of <i>Shigella dysenteriae</i> type 1 O-antigen using 3,4,6-tri- <i>O</i> -acetyl-2-azido-2-deoxy- α -D-glucopyranosyl chloride as a glycosyl donor V. Pavliak, P. Kováč and C.P.J. Glaudemans (Bethesda, MD, USA)	103
Synthesis of 2- <i>O</i> -(2-acetamido-2-deoxy- β -D-glucopyranosyl)-5a-carba- α -D-mannopyranose, and 5a-carba-2- <i>O</i> - and 3,6-di- <i>O</i> -(α -D-mannopyranosyl)- α -D-mannopyranoses S. Ogawa and K. Nishi (Yokohama, Japan)	117
Synthesis and characterization of highly substituted deoxyfluorocellulose acetate N. Kasuya, K. Iiyama and A. Ishizu (Tokyo, Japan)	131
Transacetalation: a convenient, nonaqueous method for effecting the deprotection of isopropylidene and benzylidene derivatives of sugars M.A. Andrews and G.L. Gould (Upton, NY, USA)	141
<i>Notes</i>	
Phase transfer catalyzed synthesis of 4-nitrophenyl 1-thio- β -D-glycobiosides F.D. Tropper, F.O. Andersson, C. Grand-Maitre and R. Roy (Ottawa, ON, Canada)	149

Synthesis of a repeating pentasaccharide fragment of the capsular polysaccharide of <i>Streptococcus pneumoniae</i> type 18C A.M.P. Van Steijn, J.G.M. Van der Ven, P. Van Seeventer, J.P. Kamerling and J.F.G. Vliegthart (Utrecht, Netherlands)	155
Synthesis of methyl 2-C-acetamidomethyl-2-deoxy- α -D-glucopyranoside and its <i>manno</i> isomer P. Sarda, A. Olesker and G. Lukacs (Gif-sur-Yvette, France)	161
Synthesis of a chiral polysubstituted <i>cis</i> -fused decalin containing an angular C-methyl group by an intramolecular Diels–Alder reaction S. Rémy, F. Jaekel, A. Olesker and G. Lukacs (Gif-sur-Yvette, France)	167
Preparation of building blocks for carba-oligosaccharides related to cell-surface glycans S. Ogawa, T. Tonegawa, K. Nishi and J. Yokoyama (Yokohama, Japan)	173
Synthesis of feruloylated and <i>p</i> -coumaroylated methyl glycosides R.F. Helm, J. Ralph and R.D. Hatfield (Madison, WI, USA)	183
<i>Preliminary communications</i>	
Synthesis of 1-deoxynojirimycin-containing glycans related to the Lewis X and sialyl-Lewis X epitopes recognized by LEC-CAMs H. Furui, M. Kiso and A. Hasegawa (Gifu, Japan)	C1
Methyl-3- <i>O</i> -(2-acetamido-2-deoxy-6-thio- β -D-glucopyranosyl)- β -D-galactopyranoside: a slow reacting acceptor-analogue which inhibits glycosylation by UDP-D-galactose- <i>N</i> -acetyl-D-glucosamine-(1 $\rightarrow$ 4)- β -D-galactosyltransferase Y. Kajihara, H. Hashimoto and H. Kodama (Yokohama, Japan)	C5
<i>Papers</i>	
A conformational study of α -1-Rha <i>p</i> -(1 $\rightarrow$ 2)- α -L-Rhap-(1 $\rightarrow$ OMe) by NMR nuclear Overhauser effect spectroscopy (NOESY) and molecular dynamics calculations G. Widmalm, R.A. Byrd and W. Egan (Bethesda, MD, USA)	195
A nuclear magnetic resonance spectroscopic investigation of Kdo-containing oligosaccharides related to the genus-specific epitope of <i>Chlamydia</i> lipopolysaccharides K. Bock, J.U. Thomsen (Valby, Denmark), P. Kosma, R. Christian (Vienna, Austria), O. Holst and H. Brade (Borstel, Germany)	213
Karplus-type equation for vicinal carbon–proton coupling constants for the C–S–C–H pathway in 1-thioglycosides I. Tvaroska (Bratislava, Czechoslovakia), K. Mazeau, M. Blanc-Muesser, S. Lavaitte, H. Driguez and F.R. Taravel (Grenoble, France)	225
The use of neocarribose oligosaccharides with different length and sulphate substitution as model compounds for ^1H -NMR spectroscopy S.H. Knutsen and H. Grasdalen (Trondheim, Norway)	233
NMR spectroscopy and conformational analysis of substituted 1,2:5,6-di- <i>O</i> -isopropylidene- α -D-allofuranose derivatives D.C. Lankin, S.T. Nugent and S.N. Rao (Skokie, IL, USA)	245
Binding studies on internal immunodeterminants: synthesis of β -(1 $\rightarrow$ 6)-linked oligosaccharide methyl glycosides having one to four internal D-galactopyranosyl residues flanked by gentiobiose residues T. Ziegler (Stuttgart, Germany), H. Sutoris and C.P.J. Glaudemans (Bethesda, MD, USA)	271

<i>Carbohydr. Res.</i> 229 (1992) C19–C21	C21
Synthesis of derivatives of methyl 3-deoxy-3- <i>C</i> -formyl- α -D- <i>arabino</i> -pentofuranoside S. Benefice-Malouet, P.L. Coe and R.T. Walker (Birmingham, United Kingdom)	293
Influence of malondialdehyde on the Maillard degradation of Amadori compounds A. Gómez-Sánchez, I. Hermosín and I. Maya (Seville, Spain)	307
Synthesis of D-glucose 3- and 6-[2-(perfluoroalkyl)ethyl phosphates]: a new type of anionic surfactant for biomedical use A. Milius, J. Greiner and J.G. Riess (Nice, France)	323
Interaction of linear manno-oligosaccharides with three mannose-specific bulb lectins. Compari- son with mannose/glucose-binding lectins H. Kaku and I.J. Goldstein (Ann Arbor, MI, USA)	337
New mannose-specific lectins from garlic (<i>Allium sativum</i>) and ramsons (<i>Allium ursinum</i>) bulbs H. Kaku, I.J. Goldstein (Ann Arbor, MI), E.J.M. Van Damme and W.J. Peumans (Leuven, Belgium)	347
Anticoagulant and antithrombin activities of oversulfated fucans T. Nishino and T. Nagumo (Kanagawa, Japan)	355
<i>Notes</i>	
Reaction of 2-deoxy-6- <i>O</i> -[2,3-dideoxy-4,6- <i>O</i> -isopropylidene-2,3-(<i>N</i> -tosylepimino)- α -D-man- nopyranosyl]- 4,5- <i>O</i> -isopropylidene-1,3-di- <i>N</i> -tosylstreptamine with potassium hydrogenflu- oride Y. Kobayashi, N. Taneichi, T. Tsuchiya and K. Koga (Kawasaki, Japan)	363
Epimerization of reducing terminal groups of (1 $\rightarrow$ 2)-linked D-gluco- and D-manno-disacchar- ides in aqueous sodium hydroxide H. Kobayashi, N. Shibata, S. Konno, K. Hisamichi and S. Suzuki (Miyagi, Japan)	369
<i>Announcement</i>	C11
<i>Corrigendum</i>	C12
<i>Author index</i>	C13
<i>Subject index</i>	C15
<i>Contents (vol. 229)</i>	C19