

SUBJECT INDEX

- 2-Acetamido-2-deoxy monosaccharides, determination of proton–proton distances from ^1H NMR relaxation measurements in solution, 129
- N*-Acetyllactosamine derivatives, a convenient synthesis from lactal, 159
- Agarose-gel electrophoresis, qualitative and quantitative analysis of “fast moving” and “slow moving” heparins, dermatan sulfate, and chondroitin sulfate by, 263
- 2-Amino-2-deoxyaldoses, a route to higher, by lengthening of the carbon chain of sugars by the $\text{CH}(\text{NO}_2) \cdot \text{CH}(\text{OEt})_2$ fragment, 239
- 1,2-Anhydro-3,4,6-tri-*O*-benzyl- β -D-talopyranose, the crystal structure and conformational analysis of substituted 2,7-dioxabicyclo[4,1,0]heptanes, 63
- Anion-exchange chromatography, high-performance, analysis of pectate lyase-generated oligogalacturonic acids by, with pulsed amperometric detection, 1
- Antigen from *Xanthomonas maltophilia* O19, structure of the, 249
- O-Antigen fragment, controlled acid hydrolysis of, yields univalent heptasaccharide haptens containing one 3,6-dideoxyhexose epitope, 347
- O-Antigen polysaccharide, structural studies of the *Escherichia coli* O127, 255
- Branched-chain sugars, use of methinium salts in syntheses of, 315
- Carba-disaccharide α -D-mannosidase inhibitor, an imino-linked, 341
- 5a-Carba-glucopyranose, a cellobiose phosphorylase from *Cellvibrio gilvus* recognizes only the β -D-form of, 355
- Carbon chain of sugars, a route to higher 2-amino-2-deoxyaldoses by lengthening by the $\text{CH}(\text{NO}_2) \cdot \text{CH}(\text{OEt})_2$ fragment, 239
- Cellobiose phosphorylase, from *Cellvibrio gilvus* recognizes only the β -D-form of 5a-carba-glucopyranose, 355
- Chlorodeoxy trisaccharides related to the *Shigella flexneri* Y polysaccharide, synthesis of, 195
- CMP-D-Neu5Ac-D-galactoside-(2 \rightarrow 6)- α -D-sialyltransferase, rat liver, characterization of inhibitory activities and binding mode of synthetic 6'-modified methyl *N*-acetyl- β -D-lactosaminide toward, 179
- Conformational analysis of methyl β -xylobioside, the: effect of choice of potential functions, 71
- Corn syrup solids by means of cyclododecanone as selective complexant, enhanced production of γ -cyclodextrin from, 223
- Crystal and molecular structure of threitol, 119
- Crystal and molecular structure of 2,3,4-tri-*O*-acetyl- β -D-arabinopyranosyl azide, 305
- Crystal and molecular structures of four heptitol heptaacetates, 111
- Crystal structure of 6-*O*-[(*R*)-2-hydroxypropyl]- and 6-*O*-[(*S*)-2-hydroxypropyl]-cyclomaltoheptaose, 83
- γ -Cyclodextrin from corn syrup solids by means of cyclododecanone as selective complexant, enhanced production of, 223
- 3-Deoxy-2-octulosonic acid derivatives and characterization of their 3-deoxyoctitols, synthesis of, 145
- 3,6-Dideoxyhexose epitope, controlled acid hydrolysis of an O-antigen fragment yields univalent heptasaccharide haptens containing one, 347
- Diglycosylamines in the arabinose, mannose, and rhamnose series, structure of glycosylamines and, 329
- 2,3-Dihydroxyacetophenone, formation from pentoses or hexuronic acids, 211
- 2,7-Dioxabicyclo[4,1,0]heptanes, the crystal structure and conformational analysis of substituted, 1,2-anhydro-3,4,6-tri-*O*-benzyl- β -D-talopyranose, 63
- Endopolygalacturonase, purification and characterization of biologically active (1 \rightarrow 4)-

- linked α -D-oligogalacturonides after partial digestion of polygalacturonic acid with, 9
- Enhanced production of γ -cyclodextrin from corn syrup solids by means of cyclododecanone as selective complexant, 223
- Escherichia coli* O127 O-antigen polysaccharide, structural studies of the, 255
- β -L-Fucopyranosyl and β -L-rhamnopyranosyl isothiocyanate, synthesis of 2,3,4-tri-*O*-benzoyl-, 165
- Gas-liquid chromatography with flame-ionization detection, molar-response factors for the quantitative analysis of fully methylated methyl 2-acetamido-2-deoxyhexopyranosides by, 299
- Gelatinization, internal structure of the potato starch granule revealed by chemical, 279
- Glycosaminoglycans, qualitative and quantitative analysis by agarose-gel electrophoresis, 263
- Glycosides of 3,6-di-*O*- and 3,4-di-*O*- α -D-mannopyranosyl- α -D-mannopyranose, syntheses of octyl and tetradecyl, 323
- Glycosylamines and diglycosylamines in the arabinose, mannose, and rhamnose series, structure of, 329
- Haptens, heptasaccharide, controlled acid hydrolysis of an O-antigen fragment yields univalent, containing one 2,6-dideoxyhexose epitope, 347
- Heparins (fast moving and slow moving), dermatan sulfate, and chondroitin sulfate, qualitative and quantitative analysis by agarose-gel electrophoresis, 263
- Heptakis (2,6-di-*O*-methyl)cyclomaltoheptaose complex in water, revised structure of the pyrene-1-carboxylic acid-, 337
- Heptitol heptaacetates, the crystal and molecular structures of four, 111
- Hexuronic acids or pentoses, formation of 2,3-dihydroxyacetophenone from, 211
- Hexuronic acids or pentoses, formation of reductive acid from, 217
- High-performance anion-exchange chromatography, analysis of pectate lyase-generated oligogalacturonic acids by, with pulsed amperometric detection, 1
- 6-*O*-[(*R*)-2-Hydroxypropyl]- and 6-*O*-[(*S*)-2-hydroxypropyl]-cyclomaltoheptaose, crystal structure, 83
- Imino-linked carba-disaccharide α -D-mannosidase inhibitor, an, 341
- Lactal, a convenient synthesis of *N*-acetylglucosamine derivatives from, 159
- β -D-Lactosaminide, synthetic 6'-modified methyl *N*-acetyl-, characterization of inhibitory activities toward rat liver CMP-D-Neu5Ac-D-galactoside-(2 \rightarrow 6)- α -D-sialyltransferase, 179
- Lengthening of the carbon chain of sugars by the $\text{CH}(\text{NO}_2) \cdot \text{CH}(\text{OEt})_2$ fragment, a route to higher 2-amino-2-deoxyaldoses by, 239
- (1 \rightarrow 4)-Linked α -D-oligogalacturonides, purification and characterization of biologically active, after partial digestion of polygalacturonic acid with endopolygalacturonase, 9
- Lipo-oligosaccharide from *Mycobacterium linda*, synthesis of the terminal trisaccharide unit of the, 309
- Mannopyranosides, a new route for 2,4-di-*O*-protection of, 323
- Mannopyranosyl- α -D-mannopyranose, syntheses of octyl and tetradecyl glycosides of 3,6-di-*O*- and 3,4-di-*O*- α -D-, 323
- α -D-Mannosidase inhibitor, an imino-linked carba-disaccharide, 341
- Methiniminium salts, syntheses of branched-chain sugars with, 315
- Methyl β -xylobioside, the conformational analysis of: effect of choice of potential functions, 71
- Microscopic protonation constants in tobramycin, an NMR and pH study with the aid of partially *N*-acetylated derivatives, 99
- Molecular and crystal structures of four heptitol heptaacetates, 111
- Molecular and crystal structure of threitol, 119
- Molecular and crystal structure of 2,3,4-tri-*O*-acetyl- β -D-arabinopyranosyl azide, 305
- Mycobacterium linda*, synthesis of the terminal trisaccharide unit of the lipo-oligosaccharide from, 309
- NMR and pH study of microscopic protonation constants in tobramycin, with the aid of partially *N*-acetylated derivatives, 99
- NMR relaxation measurements in solution, determination of proton-proton distances in 2-acetamido-2-deoxy monosaccharides, 129
- Nystose, computer modeling, 51
- Nystose trihydrate: crystal structure analysis and hydrogen bonding, 37
- 2-Octulosonic acid derivatives, synthesis of 3-deoxy-, and characterization of their 3-deoxyoctitols, 145
- Oligogalacturonic acids, analysis of pectate lyase-generated, by high-performance anion-exchange chromatography with pulsed amperometric detection, 1

- Oligosaccharide structures on bee venom phospholipase A₂, characterization of, 291
- Pectin, determination of the pattern of methyl esterifications in. Distributions of contiguous nonesterified residues, 21
- Pentoses or hexuronic acids, formation of 2,3-dihydroxyacetophenone from, 211
- Pentoses or hexuronic acids, formation of reductive acid from, 217
- Phospholipase A₂, bee venom, characterization of the oligosaccharide structures on, 291
- Phosphorylase, a cellobiose, from *Cellvibrio gilvus* recognizes only the β -D-form of 5a-carba-glucopyranose, 455
- Polygalacturonic acid, purification and characterization of biologically active (1 \rightarrow 4)-linked α -D-galacturonides after partial digestion of, with endopolygalacturonase, 9
- Polysaccharide from *Escherichia coli* O127, structural studies of the O-antigenic, 255
- Polysaccharide, *Shigella flexneri* Y, synthesis of chlorodeoxy trisaccharides related to the, 195
- Potato starch granule, internal structure revealed by chemical gelatinization, 279
- Proton–proton distances in 2-acetamido-2-deoxy monosaccharides, determination from ¹H NMR relaxation measurements in solution, 129
- Pyrene-1-carboxylic acid–heptakis (2,6-di-O-methyl)cyclomaltoheptaose complex in water, revised structure of the, 337
- Reductive acid, formation from pentoses or hexuronic acids, 217
- β -L-Rhamnopyranosyl and β -L-fucopyranosyl isothiocyanate, synthesis of 2,3,4-tri-O-benzoyl-, 165
- β -Rhamnopyranosylenamines, synthesis of partially protected, 165
- O-Specific polymer from *Xanthomonas maltophilia* O19, structure of the, 249
- Starch granule, internal structure of potato, revealed by chemical gelatinization, 279
- Structure of glycosylamines and diglycosylamines in the arabinose, mannose, and rhamnose series, 329
- Syntheses of branched-chain sugars with methinium salts, 315
- Synthesis of chlorodeoxy trisaccharides related to the *Shigella flexneri* Y polysaccharide, 195
- Tetrasaccharide nystose, computer modeling, 51
- Tetrasaccharide nystose trihydrate: crystal structure analysis and hydrogen bonding, 37
- Threitol, the crystal and molecular structure of, 119
- Tobramycin, an NMR and pH study of microscopic protonation constants with the aid of partially N-acetylated derivatives, 99
- 2,3,4-Tri-O-acetyl- β -D-arabinopyranosyl azide, crystal and molecular structure, 305
- Trisaccharide unit of the lipo-oligosaccharide from *Mycobacterium linda*, synthesis of the terminal, 307
- Trisaccharides, chlorodeoxy, related to the *Shigella flexneri* Y polysaccharide, synthesis of, 195
- Xanthomonas maltophilia*, structure of the O19 antigen of, 249