

SUBJECT INDEX

- Acarviosin, synthesis and inhibitory activity of eight analogues of methyl, containing a 1,6-anhydro bridge, 177
- 2-Acetamido-1,4-imino-1,2,4-trideoxy-D-galactitol as competitive inhibitor of the human lysosomal β -hexosaminidase A, 19
- Affinity labels for a β -glycosidase and a cellulase, *N*-bromoacetyl-glycopyranosylamines as, 195
- Affinity purification and characterisation of invertase from yeast using 2,5-dideoxy-2,5-imino-D-mannitol and -D-glucitol as substrate analogs, 67
- D-Aldopentoses, unprotected, Wittig reaction of with stabilized ylides: metal-ion effects, 231
- D-Allose, D-altrose, D-gulose, and D-idose, NMR studies of molybdate complexes of, 203
- Aminosubstituted α -D-glucosylmethylbenzenes and an *N*-(C- α -D-glucosylmethyl)aniline, novel α -D-glucosidase inhibitors, 161
- Amylase from *Streptomyces griseus*, transglycosylation reaction of maltotriose-forming, 57
- Amyloglucosidase substrate recognition as evaluated from its interaction with conformationally biased isomaltosides, 87
- 1,2-Anhydro hexopyranoses, preparation from hexopyranose 2-tosylates by internal displacement, 327
- N*-Bromoacetyl- β -D-glucopyranosylamine and *N*-bromoacetyl- β -cellobiosylamine as affinity labels for a β -glucosidase and a cellulase, 195
- Cell-wall polysaccharides isolated from *Alphanoascus mephitalus* and related species, chemical structure and conformational features of, 289
- Chemoenzymic synthesis of analogues of the sialyl Lewis^x and sialyl^a tetrasaccharides, 145
- Chemoenzymic synthesis of sialylated and fucosylated oligosaccharides having an *N*-acetyl-lactosaminyl core, 129
- Chitosan samples, high-performance gel-permeation chromatography of, 301
- Conformational features of cell-wall polysaccharides isolated from *Alphanoascus mephitalus* and related species, 289
- Conformationally biased isomaltosides as substrate analogs for amyloglucosidase, 87
- 1,3-Diamino-1,3-dideoxy-tetritols and their cyclic carbonic and thiocarbonic amides as competitive inhibitors for glycoside hydrolases, 31
- Diazomethyl β -D-galactopyranosyl ketone, suicide-substrate inactivation of β -galactosidase, 101
- 2,5-Dideoxy-2,5-imino-D-mannitol and -D-glucitol as glucosidase inhibitors, 67
- 2,5-Dideoxy-2,5-imino-D-mannitol and D-glucitol from 5-azido-5-deoxy-D-glucofuranose and -L-idofuranose, a chemo-enzymic synthesis, 67
- Diels–Alder cycloaddition to unsaturated sugars, stereocontrol in: reactivities of *cis* dienophiles with cyclopentadiene, 261
- Diels–Alder cycloaddition to unsaturated sugars, stereocontrol in: reactivities of acyclic seven-carbon *trans* dienophiles derived from aldopentoses, 249
- Fucosylated and sialylated oligosaccharides having an *N*-acetyl-lactosaminyl core, chemoenzymic synthesis, 129
- Fucosyltransferase of human milk, use in the chemoenzymic synthesis of analogues of the sialyl Lewis^x and sialyl Lewis^a tetrasaccharides, 145
- (β -D-Galactopyranosylmethyl)amines, *N*-substituted, as reversible inhibitors of β -D-galactosidase, 93
- β -D-Galactosidase, *N*-substituted (β -D-galactopyranosylmethyl)amines as reversible inhibitors of, 93

- β -Galactosidase, suicide-substrate inactivation of, by diazomethyl β -D-galactopyranosyl ketone, 101
- β -Galactosidases of *Escherichia coli* with substitutions for GLU-461 can be activated, by nucleophiles and can form β -D-galactosyl adducts, 9
- Galactosyl adducts, β -galactosidases of *Escherichia coli* with substitutions for GLU-461 can be activated by nucleophiles and can form β -D-, 9
- Gel-permeation chromatography of chitosan samples, high-performance, 301
- D-Glucopyranoside, synthesis of 2,3-di-O-glycosyl derivatives of methyl α - and β -, 211
- β -Glucosidase and cellulase affinity labels, 195
- α -D-Glucosidase inhibitors, aminosubstituted α -D-glucosylmethylbenzenes and an *N*-(C- α -D-glucosylmethyl)aniline as novel, 161
- C-(D-Glycopyranosyl)methylamines, synthesis and inhibition studies of, 185
- Glycosidase catalysis, hydrolysis of α - and β -D-glucosyl fluoride by individual glucosidases: new evidence for separately controlled "plastic" and "conserved" phases in, 45
- Glycosidases, hydrolysis of glycosylpyridinium ions by anomeric-configuration-inverting, 79
- Glycoside hydrolase inhibition by 1,3-diamino-1,3-dideoxy-tetritols and their cyclic carbonic and thiocarbonic amides, 31
- Glycosyl derivatives of methyl α - and β -D-glucopyranoside, synthesis of 2,3-di-O-, 211
- Glycosylpyridinium ions, hydrolysis by anomeric-configuration-inverting glycosidases, 79
- Hexopyranose 2-tosylates and their facile conversion into 1,2-anhydro hexopyranoses by internal displacement, 327
- High-performance gel-permeation chromatography of chitosan samples, 301
- Hydrolysis of glycosylpyridinium ions by anomeric-configuration-inverting glycosidases, 79
- Hydrolysis of α - and β -D-glucosyl fluoride by individual glucosidases: new evidence for separately controlled "plastic" and "conserved" phases in glycosidase catalysis, 45
- Inhibition studies and synthesis of C-(D-glycopyranosyl)methylamines, 185
- Inhibitors of α -D-glucosidase, synthesis of aminosubstituted α -D-glucosylmethylbenzenes and an *N*-(C- α -D-glucosylmethyl)aniline as, 161
- Inositol 3-(hydrogenphosphate), synthesis of the 3-C-(phosphonomethyl) analogue of 1d-*myo*-, 315
- β -Lactosides, a facile and regioselective synthesis of partially benzoylated 3,4-O-isopropylidene-, as standardized key intermediates for sialyl Lewis X (sLe^x) analogues, c1
- Lipopolysaccharide, primary structure of the polysaccharide chain of virulent *Pseudomonas solanacearum* biotype II, 335
- Lipopolysaccharides of *Pseudomonas solanacearum*, studies of O-specific polysaccharide chains consisting of structurally different repeating units, 275
- Maltotriose-forming amylase from *Streptomyces griseus*, transglycosylation reaction of, 57
- Metal-ion effects: Wittig reaction of unprotected D-aldopentoses with stabilized ylides, 231
- Methyl acarviosin, synthesis and inhibitory activity of eight analogues containing a 1,6-anhydro bridge, 177
- Methyl-5-(D-*arabino*-tetrahydroxybutyl)-3-furoylhydrazine, reactions of 2-, 323
- Methyl α - and β -D-glucopyranoside, synthesis of 2,3-di-O-glycosyl derivatives of, 211
- Molybdate complexes of D-allose, D-altrose, D-gulose, and D-idose, NMR studies of, 203
- NMR studies of molybdate complexes of D-allose, D-altrose, D-gulose, and D-idose, 203
- Nojiritetrazoles, a new class of glycosidase inhibitors, 113
- Nojiritetrazoles, selective transition state analogue inhibitors of glycosidases, 113
- (Phosphonomethyl) analogue of 1d-*myo*-inositol 3-(dihydrogenophosphate), synthesis of the 3-C-, 315
- "Plastic" and "conserved" phases in glycosidase catalysis, hydrolysis of α - and β -D-glucosyl fluoride by individual glucosidases: new evidence for separately controlled, 45
- Polysaccharide chain of virulent *Pseudomonas solanacearum* biotype II lipopolysaccharide, primary structure, 335
- Polysaccharide chains of *Pseudomonas solanacearum* lipopolysaccharide consisting of structurally different repeating units, studies of O-specific, 275
- Polysaccharides isolated from *Aphanoascus mephitalus* and related species, chemical structure and conformational features of cell-wall, 289
- Pseudo-disaccharides as potential affinity reagents for trehalase, synthesis and evaluation of diastereoisomeric alkylating, 1

- Pseudomonas solanacearum* biotype II lipopolysaccharide, primary structure of the polysaccharide chain of virulent, 335
- Pseudomonas solanacearum* lipopolysaccharides, studies of O-specific polysaccharide chains consisting of structurally different repeating units, 275
- Reactions of 2-methyl-5-(D-arabino-tetrahydroxybutyl)-3-furoylhydrazine, 323
- Sialylated and fucosylated oligosaccharides having an N-acetylglucosaminyl core, chemoenzymic synthesis, 129
- Sialyl Lewis X (sLe^x) analogues, a facile and regioselective synthesis of partially benzoylated 3,4-O-isopropylidened β -lactosides as standardized key intermediates for, c1
- Sialyl Lewis^x and sialyl Lewis^a tetrasaccharides, use of human milk fucosyltransferase in the chemoenzymic synthesis of analogues, 145
- Substitutions for GLU-461 can be activated by nucleophiles and can form β -D-galactosyl adducts, β -galactosidases of *Escherichia coli* with, 9
- Sugar moiety of methyl acarviosin, synthesis and inhibitory activity of eight analogues containing a 1,6-anhydro bridge obtained by chemical modification of the, 177
- Synthesis and evaluation of diastereoisomeric alkylating pseudo-disaccharides as potential affinity reagents for trehalase, 1
- Synthesis and inhibition studies of C-(D-glycopyranosyl)methylamines, 185
- Synthesis of 2-actamido-1,4-imino-1,2,4-trideoxy-D-galactitol and competitive inhibition of the human lysosomal β -hexosaminidase A, 19
- Synthesis of the 3-C-(phosphonomethyl) analogue of 1d-myo-inositol 3-(dihydrogenophosphate), 315
- 2-Tosylates of hexopyranoses, facile conversion into 1,2-anhydro hexopyranoses by internal displacement, 327
- Transglycosylation reaction of maltotriose-forming amylase from *Streptomyces griseus*, 57
- Transition state analogue inhibitors of glycosidases, 113
- Trehalase, synthesis and evaluation of diastereoisomeric alkylating pseudo-disaccharides as potential affinity reagents for, 1
- Unsaturated sugars, stereocontrol in Diels–Alder cycloaddition to: reactivities of acyclic seven-carbon *trans* dienophiles derived from aldopentoses, 249
- Unsaturated sugars, stereocontrol in Diels–Alder cycloaddition to: reactivities of *cis* dienophiles with cyclopentadiene, 261
- Wittig reaction of unprotected D-aldopentoses with stabilized ylides: metal-ion effects, 231