

CONTENTS

Authentic standards for the reductive-cleavage method. The positional isomers of partially methylated and acetylated or benzoylated 1,5-anhydro-D-galactitol L.E. Elvebak II, C. Abbott, S. Wall and G.R. Gray (Minneapolis, MN, USA)	1
Discrimination of the two diastereoisomeric glycosides heterodendrin and epi-heterodendrin by the combined use of NOE and molecular mechanics P.P. Lankhorst, J.W.H. Smeets (Delft, Netherlands) and C.A.G. Haasnoot (Oss, Netherlands).	17
Crystal and molecular structure of <i>N</i> -(<i>n</i> -octyl)-6-deoxy-D-gluconamide: a novel packing of amphiphilic molecules R. Herbst, T. Steiner (Berlin, Germany), B. Pfannemüller (Freiburg i. Br., Germany) and W. Saenger (Berlin, Germany).	29
Crystal structures of methyl 6- <i>O</i> - <i>n</i> -alkanoyl- β -D-glucopyranosides Y. Abe, K. Harata (Ibaraki, Japan), M. Fujiwara and K. Ohbu (Tokyo, Japan)	43
Synthesis of carbocyclic analogues of 3-deoxy-D- <i>manno</i> -2-octulosonic acid and <i>N</i> -acetylneuraminic acid S. Ogawa, M. Yoshikawa, T. Taki, S. Yokoi and N. Chida (Yokohama, Japan)	53
A new synthesis of 6- <i>O</i> -acylsucroses and of mixed 6,6'-di- <i>O</i> -acylsucroses K. Baczko, C. Nugier-Chauvin (Rennes, France), J. Banoub (Newfoundland, Canada), P. Thibault (Nova Scotia, Canada) and D. Plusquellec (Rennes, France)	79
Highly selective nitroxyl radical-mediated oxidation of primary alcohol groups in water-soluble glucans A.E.J. De Nooy, A.C. Besemer (Zeist, Netherlands) and H. Van Bakkum (Delft, Netherlands).	89
Bromination of sugar enones and enonolactones C. Di Nardo, O. Varela, R.M. De Lederkremer, R.F. Baggio, D.R. Vega (Buenos Aires, Argentina) and M.T. Garland (Santiago de Chile, Chile)	99
TLC-LSIMS of neoglycolipids of glycosaminoglycan disaccharides and of oxymercuration cleavage products of heparin fragments that contain unsaturated uronic acid W. Chai, J.R. Rosankiewicz and A.M. Lawson (Harrow, United Kingdom)	111
Structural studies of the O-specific chain and a core hexasaccharide of <i>Hafnia alvei</i> strain 1192 lipopolysaccharide W. Jachymek (Wrocław, Poland), C. Petersson (Uppsala, Sweden), A. Helander (Stockholm, Sweden), L. Kenne (Uppsala, Sweden), C. Lugowski and T. Niedziela (Wrocław, Poland) . . .	125

Characterisation by LSI-MS and ^1H NMR spectroscopy of tetra-, hexa-, and octa-saccharides of porcine intestinal heparin W. Chai (Harrow, United Kingdom), E.F. Hounsell, C.J. Bauer (London, United Kingdom) and A.M. Lawson (Harrow, United Kingdom)	139
The structure of the O-specific polysaccharide from <i>Thiobacillus</i> sp. IFO 14570, with three different diaminopyranoses forming the repeating unit A.S. Shashkov (Moscow, Russian Federation), S. Campos-Portuguez, H. Kochanowski (Freiburg i.Br.-Zähringen, Germany), A. Yokota (Osaka, Japan) and H. Mayer (Freiburg i.Br.-Zähringen, Germany)	157
Analysis of positions of substitution of <i>O</i> -acetyl groups in partially <i>O</i> -acetylated cellulose by the reductive-cleavage method C.K. Lee and G.R. Gray (Minneapolis, MN, USA)	167
<i>Notes</i>	
Full assignment of the NMR spectrum of the capsular polysaccharide from <i>Streptococcus pneumoniae</i> serotype 10A C. Jones (South Mimms, Herts, United Kingdom)	175
The structural analysis of disaccharides from red algal galactans by methylation and reductive partial-hydrolysis R. Falshaw and R.H. Furneaux (Lower Hutt, New Zealand)	183
<i>Preliminary communication</i>	
Structural investigation on the carbohydrate backbone of the lipopolysaccharide from <i>Klebsiella pneumoniae</i> rough mutant R20/O1 ⁻ M. Süsskind (Borstel, Germany), S. Müller-Loennies (Valby, Denmark), W. Nimmich (Rostock, Germany), H. Brade and O. Holst (Borstel, Germany)	C1
<i>Regular papers</i>	
Synthesis of 3,4-di- <i>O</i> -acetyl-2,5-anhydro-1,6-dideoxy-1,6-diiodo-D-mannitol. Comparison of NMR spectral results for the solid state and solution with those of the X-ray structural determination M.A. Shalaby, F.R. Fronczek, Y. Lee and E.S. Younathan (Baton Rouge, LA, USA)	191
Cyclomaltoheptaose (β -cyclodextrin) and hydroxyethyl-substituted β -cyclodextrin inclusion complex formation with chlorogenic acid: solvent effects on inclusion complex stability P.L. Irwin, J.N. Brouillette and K.B. Hicks (Philadelphia, PA, USA)	201
Solid-phase synthesis of a glycosylated hexapeptide of human sialophorin, using the trichloroacetimidate method J. Rademann and R.R. Schmidt (Konstanz, Germany)	217
Stereoselective synthesis of a blood group A type glycopeptide present in human blood mucin W.M. Macindoe (Tokyo, Japan), H. Ijima, Y. Nakahara (Saitama, Japan) and T. Ogawa (Saitama, Tokyo, Japan)	227
Synthesis of <i>N</i> - β -D-glucopyranosyluronate derivatives of barbital, phenobarbital, metharbital, and mephobarbital S.M. Neighbors, W.H. Soine and S.G. Paibir (Richmond, VA, USA)	259
Enzymic transfer of 6-modified D-galactosyl residues: synthesis of biantennary penta- and hepta-saccharides having two 6-deoxy-D-galactose residues at the nonreducing end and evaluation of 6-deoxy-D-galactosyl transfer to glycoprotein using bovine β -(1 $\rightarrow$ 4)-galactosyltransferase and UDP-6-deoxy-D-galactose Y. Kajihara, T. Endo (Nagatsuta, Japan), H. Ogasawara, H. Kodama (Umeaoka, Japan) and H. Hashimoto (Nagatsuta, Japan)	273

The structure of the capsular polysaccharide of <i>Escherichia coli</i> O8:K43:H11 Y.M. Choy (Shatin, Hong Kong), G.G.S. Dutton (Vancouver, Canada), M.R. Leslie, H. Parolis and L.A.S. Parolis (Grahamstown, South Africa)	295
The structure of a novel polysaccharide produced by <i>Bradyrhizobium</i> species within soybean nodules J. An, R.W. Carlson, J. Glushka (Athens, GA, USA) and J.G. Streeter (Wooster, OH, USA) .	303
NMR Studies of acetan and the related bacterial polysaccharide, CR1/4, produced by a mutant strain of <i>Acetobacter xylinum</i> I.J. Colquhoun, M. Defernez and V.J. Morris (Colney, United Kingdom)	319
<i>Notes</i>	
Separation and quantitation of enantiomeric galactoses and their mono- <i>O</i> -methylethers as their diastereomeric acetylated 1-deoxy-1-(2-hydroxypropylamino)alditols M.R. Cases, A.S. Cerezo and C.A. Stortz (Buenos Aires, Argentina)	333
An alternative synthesis of (+)- <i>epiallo</i> -muscarine from D-glucose V. Popsavin, O. Berić, M. Popsavin, J. Csanádi and D. Miljković (Novi Sad, Yugoslavia)	343
Isopropylidenation of aldose bis(phenylhydrazones) E.S.H. El Ashry, N. Rashed, A. Mousaad and M. El Habrouk (Alexandria, Egypt)	349
The preparation of sucrose monophosphates from dried mixtures of sucrose and sodium phosphate E. Tarelli and S.F. Wheeler (Potters Bar, United Kingdom)	359
Isolation, purification, and characterization of cyclomaltododecaose (η -cyclodextrin) T. Endo, H. Ueda (Tokyo, Japan), S. Kobayashi (Ibaraki, Japan) and T. Nagai (Tokyo, Japan)	369
<i>Author index</i>	C9
<i>Subject index</i>	C11
<i>Contents</i>	C15