

Contents

Regular papers

Structure of the saponin adjuvant QS-21 and its base-catalyzed isomerization product by ^1H and natural abundance ^{13}C NMR spectroscopy N.E. Jacobsen, W.J. Fairbrother (South San Francisco, CA, USA), C.R. Kensil (Worcester, MA, USA), A. Lim (South San Francisco, CA, USA), D.A. Wheeler (Worcester, MA, USA) and M.F. Powell (South San Francisco, CA, USA)	1
Vacuum ultraviolet circular dichroism of gellan-family polymer films from water and dimethyl sulfoxide E.R. Arndt and E.S. Stevens (Binghamton, NY, USA)	15
Alkaline degradation of glucose: effect of initial concentration of reactants B.Y. Yang and R. Montgomery (Iowa City, IA, USA)	27
Alkaline degradation of fructofuranosides B.Y. Yang and R. Montgomery (Iowa City, IA, USA)	47
Transglycosylation reactions of permethylated methyl D-glucopyranosides with partially methylated 1,5-anhydroalditols C.K. Lee and E.J. Kim (Chuncheon, Korea)	59
Rationally designed syntheses of high-mannose and complex type undecasaccharides Y. Nakahara, S. Shibayama, Y. Nakahara (Saitama, Japan) and T. Ogawa (Tokyo, Japan)	67
Degradation of double-stranded xanthan by hydrogen peroxide in the presence of ferrous ions: comparison to acid hydrolysis B.E. Christensen, M.H. Myhr and O. Smidsrød (Trondheim, Norway)	85
Chemoenzymatic synthesis of neuraminic acid analogs structurally varied at C-5 and C-9 as potential inhibitors of the sialidase from influenza virus M. Murakami, K. Ikeda and K. Achiwa (Shizuoka, Japan)	101
Structural investigation of the polysaccharide fraction from the mucilage of <i>Dicerocaryum zanguebaricum</i> Merr. G. Barone, M.M. Corsaro (Naples, Italy), M. Giannattasio (Portici, Italy), R. Lanzetta, M. Moscardiello and M. Parrilli (Naples, Italy)	111
Notes	
Capillary electrophoresis separation of mucin motif MUC 5AC glycopeptides from in vitro O-glycosylation reactions S. Hennebicq-Reig, B. Soudan, D. Tetaert, G. Briand, C. Richet, D. Demeyer and P. Degand (Lille, France)	121

1-Aryl-6,7-dichloro-3- β -D-erythrofuranosylpyrazolo[3,4- <i>b</i>]quinoxaline C-nucleoside analogues M.A.E. Sallam, H.M. El Nahas, S.M.E. Abdel Megid (Alexandria, Egypt) and T. Anthonsen (Trondheim, Norway)	127
Facile preparation of phenyl 1-thioglycosides of partially methylated maltooligosaccharides by re- stricted thiolysis of fully methylated cyclodextrins N. Sakairi (Sapporo, Japan) and H. Kuzuhara (Urawa, Japan)	139
Thioanhydro sugar derivatives. Action of base on methyl 6-thio- α -D-galactopyranoside derivatives I. Izquierdo Cubero, M.T. Plaza Lopez-Espinosa and A. Saenz de Buruaga Molina (Granada, Spain).	145
Synthesis of some methyl 6- <i>O</i> -alkyl- α -D-glycopyranosides F. Wijnbergen, H. Regeling, B. Zwanenburg, G.J.F. Chittenden (Nijmegen, The Netherlands) and N. Rehnberg (Perstorp, Sweden).	151
Effect of <i>n</i> -alkyl glucosides on waxy maize and wheat starch retrogradation C.G. Biliaderis, D.J. Prokopowich (Winnipeg, Canada), M.R. Jacobson and J.N. BeMiller (West Lafayette, IN, USA)	157
Aldo-enol transition in periodate-oxidized dextrans S.N. Drobchenko, L.S. Isaeva-Ivanova, A.R. Kleiner and E.V. Eneyskaya (St. Petersburg, Russian Federation)	171
<i>Instructions to authors</i>	C1
<i>Regular papers</i>	
Two-bond ^{13}C - ^{13}C spin-coupling constants in carbohydrates: effect of structure on coupling magni- tude and sign T. Church, I. Carmichael and A.S. Serianni (Notre Dame, IN, USA)	177
Conformation of 4-thio-L-lyxono-1,4-lactone in solution and in the crystalline state O. Varela, P.A. Zunszain, D.O. Cicero, R.F. Baggio, D.R. Vega (Buenos Aires, Argentina) and M.T. Garland (Santiago de Chile, Chile)	187
Identification of the acidic degradation products of hexenuronic acid and characterisation of hex- enuronic acid-substituted xylooligosaccharides by NMR spectroscopy A. Teleman, T. Hausalo, M. Tenkanen and T. Vuorinen (Espoo, Finland).	197
Synthesis of methyl 2- <i>O</i> -allyl-(and 3- <i>O</i> -allyl)-5- <i>O</i> -benzyl- β -D-ribofuranoside T. Desai, J. Gigg and R. Gigg (London, UK)	209
Synthesis of [$6''$ - ^3H]-, ($6''$ - ^2H)- and (2 - ^2H)-maltotriose B. Sauerbrei, J. Niggemann, S. Gröger, S. Lee and H.G. Floss (Seattle, WA, USA)	223
Migration of secondary <i>tert</i> -butyldimethylsilyl groups in cyclomalto-heptaose and -octaose derivatives D. Icheln, B. Gehrcke, Y. Piprek, P. Mischnick, W.A. König (Hamburg, Germany), M.A. Dessoy and A.F. Morel (Santa Maria, Brazil)	237
Glycosylation with <i>N</i> -Troc-protected glycosyl donors U. Ellervik and G. Magnusson (Lund, Sweden).	251
Synthesis of precursors for the dimeric 3- <i>O</i> -SO ₃ Na Lewis X and Lewis A structures G.V. Reddy, R.K. Jain, R.D. Locke and K.L. Matta (Buffalo, NY, USA)	261

A method for the synthesis of <i>C</i> -(2-deoxy- β -glycosyl)arenes A.T. Khan, W. Ahmed and R.R. Schmidt (Konstanz, Germany)	277
A facile synthesis of 5-thio-L-fucose and 5-thio-D-arabinose from D-arabinose M. Izumi, O. Tsuruta and H. Hashimoto (Yokohama, Japan).	287
Structure of the phosphopeptidomannans from flocculent and non-flocculent yeast <i>Kluyveromyces lactis</i> M. Bilang, F. Attioui, V. Loppinet (Nancy, France), J.-C. Michalski (Villeneuve d'Ascq, France) and R. Bonaly (Nancy, France)	303
<i>Notes</i>	
Heptakis[6- <i>S</i> -(2,3-dihydroxypropyl)-6-thio]cyclomaltoheptaose and its sulfone: water-soluble β -cyclo-dextrin derivatives having modified polarity H.H. Baer (Ottawa, Canada) and F. Santoyo-González (Granada, Spain)	315
Synthesis of α - and β -L-Fuc <i>p</i> -(1 $\rightarrow$ 2)- α -L-Fuc <i>p</i> -(1 $\rightarrow$ 3)- β -D-Gal <i>p</i> NAc glycosides identified as termini of <i>Schistosoma mansoni</i> oligosaccharides V.P. Kamath and O. Hindsgaul (Edmonton, Canada)	323
Synthesis of <i>N</i> -acetyl-4,8-dideoxyneuraminic acid-containing ganglioside GM ₃ M. Yoshida, H. Ishida, M. Kiso and A. Hasegawa (Gifu, Japan)	331
Chlorodeoxy derivatives from D-galactochloralose Ö. Makinabakan, Y.G. Salman and L. Yüceer (İzmir, Turkey)	339
A one-pot procedure for the quantitative conversion of glycosides into acetylated glycosyl fluorides B.A. Bergamaschi and J.I. Hedges (Seattle, WA, USA)	345
Synthesis of partially protected 1,5-anhydroalditols by hydroboration of glycals R. Murali and M. Nagarajan (Hyderabad, India)	351
Synthesis of the 4-, 6-deoxy, and 4,6-dideoxy derivatives of D-mannose T. Nishio, Y. Miyake, K. Kubota, M. Yamai, S. Miki, T. Ito and T. Oku (Tokyo, Japan).	357
<i>Author Index</i>	C11
<i>Subject Index</i>	C13
<i>Cumulative author index, Vols. 261–280</i>	C19
<i>Contents</i>	C35