

Contents

iii Graphical Abstracts

Regular papers

- | | | |
|--|-----|--|
| H.-J. Zhong, M.A.K. Williams, D.M. Goodall (York, UK) and M.E. Hansen (Lille Skensved, Denmark) | 1 | Capillary electrophoresis studies of pectins |
| K. Mernissi-Arifi, G. Schlewer and B. Spiess (Illkirch, France) | 9 | Intramolecular acid-base properties of <i>myo</i> -inositol 1,2,6-trisphosphate analogues: influence of the hydroxyl groups, phosphate configuration and intracyclic atom substitution |
| B. Aguilera, J. Jiménez-Barbero and A. Fernández-Mayoralas (Madrid, Spain) | 19 | Conformational differences between Fuc (α 1–3) GlcNAc and its thioglycoside analogue |
| D. Le Botlan, Y. Rugraff, C. Martin and P. Colonna (Nantes, France) | 29 | Quantitative determination of bound water in wheat starch by time domain NMR spectroscopy |
| G. Crini, C. Cosentino, S. Bertini, A. Naggi, G. Torri (Milan, Italy), C. Vecchi (San Donato Milanese, Italy), L. Janus and M. Morcellet (Villeneuve d'Ascq, France) | 37 | Solid state NMR spectroscopy study of molecular motion in cyclomaltoheptaose (β -cyclodextrin) crosslinked with epichlorohydrin |
| B. Zipser, J.J. Bradford and R.I. Hollingsworth (East Lansing, USA) | 47 | Structural analysis of leech galactocerebrosides using 1D and 2D NMR spectroscopy, gas chromatography–mass spectrometry, and FAB mass spectrometry |
| F. Inglesias-Guerra, I. Romero, F. Alcudia and J.M. Vega-Pérez (Sevilla, Spain) | 57 | Alkylating agents from sugars. Cyclophosphamides derived from 2-amino-2-deoxy-D-allose. |
| G. Hodosi and P. Kováč (Bethesda, USA) | 63 | Glycosylation via locked anomeric configuration: stereospecific synthesis of oligosaccharides containing the β -D-mannopyranosyl and β -L-rhamnopyranosyl linkage |
| B. Becker and J. Thiem (Hamburg, Germany) | 77 | Regio- and stereoselective introduction of ether-linked carboxylic side chains into carbohydrates by conjugate addition reactions |
| D. Horton and O. Varela (Columbus, USA) | 85 | Synthesis of 4,6-diamino-5-(polyhydroxyalkylamido)pyrimidines: conformation of the sugar chain |
| A. Malleron and S. David (Orsay, France) | 93 | A preparation of protected 2-deoxy-2-hydroxymethyl-D-mannose and -D-glucose derivatives not involving organometallic reagents |
| A. Kardošová, M. Matulová and A. Malovíková (Bratislava, Slovak Republic) | 99 | (4-O-Methyl- α -D-glucurono)-D-xylan from <i>Rudbeckia fulgida</i> var. <i>sullivantii</i> (Boynton et Beadle) |
| R. Falshaw, R.H. Furneaux and D.E. Stevenson (Lower Hutt, New Zealand) | 107 | Agars from nine species of red seaweed in the genus <i>Curdiea</i> (Gracilariaceae, Rhodophyta) |
| T. Nishimura, M. Ishihara, T. Ishii and A. Kato (Ibaraki, Japan) | 117 | Structure of neutral branched xylooligosaccharides produced by xylanase from <i>in situ</i> reduced hardwood xylan |
| N. Wellner (Norwich, UK), M. Kačuráková, A. Malovíková (Bratislava, Slovakia), R.H. Wilson and P.S. Belton (Norwich, UK) | 123 | FT-IR study of pectate and pectinate gels formed by divalent cations |
| P.J. Jenkins and A.M. Donald (Cambridge, UK) | 133 | Gelatinisation of starch: a combined SAXS/WAXS/DSC and SANS study |

W.G.T. Willats, S.E. Marcus and J.P. Knox (Leeds, UK)	149	Generation of a monoclonal antibody specific to (1→5)- α -L-arabinan
Notes		
T. Oshikawa, M. Yamashita (Hamamatsu, Japan), K. Seo and Y. Hamazu (Numazu, Japan)	153	X-ray single crystal structure analyses of 5-deoxy-5-C-(alkylphosphinyl)-glucopyranose
Y. Du and R.J. Linhardt (Iowa City, USA)	161	Stereospecific synthesis of α -C-glycosyl derivatives (" α -C-glycosides") of <i>N</i> -acetylneuraminic acid by samarium-mediated reductive desulfonylation of a glycosyl phenylsulfone
T. Das and M.S. Shashidhar (Pune, India)	165	Racemic 2,4-di- <i>O</i> -benzoyl- <i>myo</i> -inositol 1,3,5-orthoformate: a versatile intermediate for the preparation of <i>myo</i> -inositol phosphates
M. Dromowicz and P. Köll (Oldenburg, Germany)	169	A convenient synthesis of D-idose.
B.M. Smith, J.L. Owens, C.N. Bowman and P. Todd (Boulder, USA)	173	Thermodynamics of borate ester formation by three readily grafted carbohydrates
P. Borrachero-Moya, F. Cabrera-Escribano, M. Gómez-Guillén and M. Del Rocío Paredes-León (Seville, Spain)	181	Synthesis of 4-(4,6-di- <i>O</i> -benzyl-2,3-dideoxy- β -D- <i>erythro</i> -hex-2-enopyranosyl)pyrazoles from 3,4,6-tri- <i>O</i> -acetyl-D-glucal
J. Wang, P. Kováč (Bethesda, USA), P. Sinaÿ (Paris, France) and C.P.J. Glaudemans (Bethesda, USA)	191	Synthetic C-oligosaccharides mimic their natural, analogous immunodeterminants in binding to three monoclonal immunoglobulins
S. Freimund and S. Köpper (Oldenburg, Germany)	195	Dimeric structures of 1,5-anhydro-D-fructose
M. Ohnishi (Kyoto, Japan), G. Okada (Shizuoka, Japan) and T. Yazaki (Kyoto, Japan)	201	Characterization of the subunit structure of the β -glucosidase from <i>Aspergillus niger</i> , an aspect of the mechanism of carbohydrate recognition
A.K. Choudhury and N. Roy (Calcutta, India)	207	Synthesis of some galactofuranosyl disaccharides using a galactofuranosyl trichloroacetimidate as donor
T. Kumazawa, N. Asahi, S. Matsuba, S. Sato (Yamagata, Japan), K. Furuhashi (Tokyo, Japan) and J.-I. Onodera (Yamagata, Japan)	213	Conversion of β -D-C-glucopyranosyl phloroacetophenone to a spiroketal compound
I. Izquierdo Cubero, M.T. Plaza López-Espinosa, M. Rodríguez Alonso, R. Asenjo Asenjo and A. Ramírez Fernández (Granada, Spain)	217	The stereoselective synthesis of novel 4-octulose derivatives
G.R. Smith, J.J. Finley IV and R.M. Giuliano (Villanova, USA)	223	Synthesis of methyl α -L-callipeltoside
V. Pozsgay (Bethesda, USA), N. Sari and B. Coxon (Gaithersburg, USA)	229	Measurement of interglycosidic $^3J_{CH}$ coupling constants of selectivity ^{13}C labeled oligosaccharides by 2D <i>J</i> -resolved 1H NMR spectroscopy
H.P. Ramesh and R.N. Tharanathan (Mysore, India)	239	Structural characteristics of a mixed linkage β -D-glucan from sorghum (<i>Sorghum bicolor</i>)
	245	Apology
	iii	Graphical Abstracts
Regular papers		
I. Bajza, K.E. Kövér and A. Lipták (Debrecen, Hungary)	247	Synthesis of <i>p</i> -trifluoroacetamidophenyl (4,6-dideoxy-4-formamido-3-C-methyl-2- <i>O</i> -methyl- α -L-mannopyranosyl)-(1→3)-(2- <i>O</i> -methyl- α -D-rhamnopyranosyl)-(1→3)-(2- <i>O</i> -methyl- α -L-fucopyranosyl)-(1→3)-(α -L-rhamnopyranosyl)-(1→2)-6-deoxy- α -L-talopyranoside: a spacer-armed pentasaccharide glycopeptidolipid antigen of <i>Mycobacterium avium</i> serovar 14
J. Bartek, R. Müller and P. Kosma (Vienna, Austria)	259	Synthesis of a neoglycoprotein containing the Lewis X analogous trisaccharide β -D-GalpNAc-(1→4)[α -L-Fucp-(1→3)]- β -D-GlcpNAc
M. Bukowska, M. Maciejewski and J. Prejzner (Warszawa, Poland)	275	Trimethylsilylation of cyclodextrins with <i>N</i> -(trimethylsilyl)acetamide in <i>N,N</i> -dimethylformamide

S.-Q. Zhang, Z.-J. Li, A.-B. Wang, M.-S. Cai and R. Feng (Beijing, P.R. China)	281	Synthesis of a phenylpropanoid glycoside, Osmanthuside B6
C. Krog-Jensen and S. Oscarson (Stockholm, Sweden)	287	Efficient synthesis of differently protected methyl (ethyl 1-thio- β -D-glucopyranosid)uronates and their evaluation as glucuronic acid donors and acceptors
É. Bozó, S. Boros, J. Kuzsman, E. Gács-Baitz and L. Párkányi (Budapest, Hungary)	297	An economic synthesis of 1,2,3,4-tetra-O-acetyl-5-thio-D-xylopyranose and its transformation into 4-substituted-phenyl 1,5-dithio-D-xylopyranosides possessing antithrombotic activity
M. Pagliaro (Palermo, Italy)	311	Autocatalytic oxidations of primary hydroxyl groups of cellulose in phosphoric acid with halogen oxides
B.-G. Huang and M. Bobek (Buffalo, USA)	319	Synthesis and in vitro antitumor activity of some amino-deoxy 7-hexofuranosyl-pyrrolo[2,3- <i>d</i>] pyrimidines
K.M. Halkes, D.J. Lefeber, C.T.M. Fransen, J.P. Kamerling and J.F.G. Vliegthart (Utrecht, The Netherlands)	329	Synthesis of the spacer-containing β -D-GalpNAc-(1 $\rightarrow$ 4)- β -D-GlcpNAc-(1 $\rightarrow$ 3)- α -D-Galp moiety, representing the non-fucosylated backbone trisaccharide of the glycocalyx glycan of the parasite <i>Schistosoma mansoni</i>
Q. Ding (Wuhan, China), S. Jiang (Shatin, Hong Kong), L. Zhang (Wuhan, China) and C. Wu (Shatin, Hong Kong)	339	Laser light-scattering studies of pachyman
S.J. Gebbie, I. Gosney (Edinburgh, UK), P.R. Harrison (Widnes, UK), I.M.F. Lacan (Edinburgh, UK), W.R. Sanderson and J.P. Sankey (Widnes, UK)	345	A convenient large-scale synthesis of methyl α -maltoside: a simple model for amylose
M.M. Corsaro, C. De Castro (Napoli, Italy), A. Evidente (Portici, Italy), R. Lanzetta, A. Molinaro (Napoli, Italy), L. Mugnai (Firenze, Italy), M. Parrilli (Napoli, Italy) and G. Surico (Firenze, Italy)	349	Chemical structure of two phytotoxic exopolysaccharides produced by <i>Phomopsis foeniculi</i>
H. Jacobs, R.C. Eerlingen, N. Rouseu (Heverlee, Belgium), P. Colonna (Nantes, France) and J.A. Delcour (Heverlee, Belgium)	359	Acid hydrolysis of native and annealed wheat, potato and pea starches—DSC melting features and chain length distributions of lintnerised starches
D. Zhan, P. Janssen and A.J. Mort (Stillwater, USA)	373	Scarcity or complete lack of single rhamnose residues interspersed within the homogalacturonan regions of citrus pectin
H. Baumann, H. Scheen, B. Huppertz and R. Keller (Aachen, Germany)	381	Novel regio- and stereoselective O-6-desulfation of the glucosamine moiety of heparin with <i>N</i> -methylpyrrolidinone–water or <i>N,N</i> -dimethylformamide–water mixtures
T. Mishima, M. Hisamatsu (Tsu, Japan), W.S. York (Athens, USA), K. Teranishi and T. Yamada (Tsu, Japan)	389	Adhesion of β -D-glucans to cellulose
W.B. Severn (Upper Hutt, New Zealand), R.H. Furneaux, R. Falshaw (Lower Hutt, New Zealand) and P.H. Atkinson (Upper Hutt, New Zealand)	397	Chemical and spectroscopic characterisation of the phosphatidylinositol manno-oligosaccharides from <i>Mycobacterium bovis</i> AN5 and WAg201 and <i>Mycobacterium smegmatis</i> mc ² 155
F. Chavagnat (Villeurbanne, France), A. Heyraud, P. Colin-Morel (Grenoble, France), M. Guinand and J. Wallach (Villeurbanne, France)	409	Catalytic properties and specificity of a recombinant, overexpressed D-mannuronate lyase
A. Heyraud, P. Colin-Morel, C. Gey (Grenoble, France), F. Chavagnat, M. Guinand and J. Wallach (Villeurbanne, France)	417	An enzymatic method for preparation of homopolymannuronate blocks and strictly alternating sequences of mannuronic and guluronic units
A. Percy, H. Ono and K. Hayashi (Ibaraki, Japan)	423	Acceptor specificity of cellobiose phosphorylase from <i>Cellvibrio gilvus</i> : synthesis of three branched trisaccharides
Notes		
J. Madaj, A. Trynda, A. Konitz, A. Nowacki and A. Wiśniewski (Gdańsk, Poland)	431	X-ray diffraction and high resolution NMR spectroscopy analysis of methyl β -D-glucofuranosidurono-6,3-lactone

A. Maras, M. Erden, H. Seçen and Y. Sütbeyaz (Erzurum, Turkey)	435	One-pot synthesis from 1,4-cyclohexadiene of ($\pm$)-1,4/2,5-cyclohexanetetrol, a naturally occurring cyclitol derivative
S. Tsujihata and F. Nakatsubo (Kyoto, Japan)	439	A novel synthetic method for α -D-galactofuranose 1,2,5-orthopivalate
	445	Book review
	I	Author index
	III	Subject index
	IX	Volume contents